


Bodleian Libraries

UNIVERSITY OF OXFORD

This book is part of the collection held by the Bodleian Libraries and scanned by Google, Inc. for the Google Books Library Project.

For more information see:

http://www.bodleian.ox.ac.uk/dbooks



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 2.0 UK: England & Wales (CC BY-NC-SA 2.0) licence.



ent to Bodleian
1817.

8vo BS.
I. 182

HORTUS BENGALENSIS,

OR

A CATALOGUE

OF THE

PLANTS

GROWING IN

THE HONOURABLE EAST INDIA COMPANY's

Botanic Garden

AT

CALCUTTA.

Serampore

PRINTED AT THE MISSION PRESS.

1814.


BIBLIOTHECA · BODLEIANA ·

EXPLANATORY REMARKS.

IN the following Catalogue, I have chosen the *Linnæan* Sexual mode of arrangement, in preference to the alphabetical, or any other; because it appears the most convenient for those who have a knowledge of that system; and for those who have not, an Index will obviate every difficulty.

That we ought to adopt the *Linnæan* names only, must appear evident to every one on the slightest reflection; particularly here in *India*, where the number of well established, or applicable English names bears so very small a proportion to the whole. Many of the former are become, by long use, perfectly familiar to our ears; such as *Geranium, Narcissus, Convolvulus, Ranunculus*, &c. &c. and no doubt practice will in time make the whole equally so.

All the nations of *Europe* know well the advantages they derive from the general knowledge of the same figures, or characters in Arithmetic, Chemistry, &c. &c. Botanists, particularly those who are not masters of the science, must derive equal advantages from a general knowledge of one universal nomenclature; I will therefore endeavour to contribute my mite, to render a more general communication of Botanic knowledge less difficult, by omitting, in this Catalogue, the usual column for English names, and consign the few used, to that appropriated for the Asiatic names; such as the *Sanskrit, Hindoostani, Bungali, Tamul, Telinga*, &c. &c. provincial, or vernacular names, whenever they can be ascertained with any tolerable degree of certainty. I say with any tolerable degree of certainty; for the natives of *India*, even the more learned, particularly in *Bengal*, do not always agree, except when the plant is particularly conspicuous for some peculiar quality, use, or habit.

With the view of compressing this catalogue into as small a size as could conveniently be done; and at the same time conveying as much information as can well be expected from a catalogue; more abbreviations, or characters, have been employed than are usual in Botanical publications.

In the column of synonyma they are,

1. S. for *Sanskrit.*

2. A. *Arabic.*

3. P. *Persian.*

4. H. *Hindoostani, Hindi,* or *Hindooee,* the language used in every part of *Hindoostan* proper, and the *Dukhin.*

5. B. *Gouree,* or as it is commonly called *Bungala,* or *Bungali,* the language spoken in all the provinces of *Bengal.*

6. U. *Uriya,* the language in use over *Orissa,* often called *Orixa ;* extending from *Medini poora,* to between *Ganjam,* and *Manakapatam ;* and inland to *Cammal-poora.*

7. T. *Tailanga, Telunga,* or *Telinga,* the language of a nation of the same name; extending along the coast of *Coromandel,* from between *Ganjam* and *Manakapatam,* to between *Ungoli* and *Nelore.*

8. Tam. *Tamel, Tamul,* &c. often called *Malabar.* The language of the inhabitants of the east, and south end of the Peninsula of *India ;* its northern limits between *Nelore,* and *Ungoli.* This country is called *Dravira* by Mr. Colebrooke, see *Asiat. Researches,* 7. p. 226.

9. Can. *Canara.* The language, or Dialect of *Mysore,* &c. intermediate mountainous tracts, between both coasts of the peninsula.

10. M. *Mahratta,* or *Marhatta,* the language of a numerous, well known people of the same name.

11. C. *Cingalese, Cinghalese,* or *Ceylonese,* the language of the inhabitants of the Island of Ceylon.

12. M-n. *Malayan,* the language of the Malays.

13. Ch. *Chinese.*

14. N. *Nawar,* or the language of *Napaul.*

15. E. *English.*

Abbreviations, or characters found in the column *Duration and Habit,* (by the latter term, I mean external appearance only,) are,

⊙ Annual.

H. Herbaceous, perishing down to the root annually.

♂ Biennial.

♃ Perennial.

♄ Shrubby.

T. Tree.

Any one of the following letters coming after any of the above characters, denote the habit of the plant to be as follows, viz.

L. Large. e. g. L. T. a large tree.

S. Small. S. ♄ a small shrub.

E. Erect.

D. Diffuse, or spreading much.

C. Cespitose, spreading close on the ground; and includes the Linnæan term *prostratus*, or prostrate.

Sar. Sarmentose, such as produce long slender shoots, or runners; with bunches of leaves, and roots from their joints; like the Strawberry, &c.

Cr. Creeping, or running along the ground, &c. and sending forth roots as they extend themselves.

R. Radicans, clinging to any other body for support, by means of fibres which do not imbibe nourishment; as *Ivy*, &c. *Smith's Introduction to Botany*, p. 118.

Sc. Scandent, or climbing; such as cannot support themselves, but ascend, or climb by means of tendrils, or otherwise.

T. Twining (commonly called creepers,) ascending spirally round their prop.

P. Parasitical, growing on some other plant, not on the ground.

A. Aquatic.

The two last columns, *Time of Flowering (Estivation) and ripening their seed, (Maturation;)* to save room are also marked with the number of the month, or the letters.

C. S. The *cool season;* from the beginning of November, to the end of February.

H. S. *Hot season;* from the beginning of March, to the middle or end of June.

D. S. *Dry season;* from the beginning of November, to the middle or end of June.

R. S. *Rainy season;* from the middle of June, or beginning of July, to the end of October.

O. The whole year round.

W. ROXBURGH.

INTRODUCTION.

IT having been necessary for the superintendant of the Honourable Company's Botanical Garden to make a voyage to sea, for the recovery of his health, I embrace the present occasion of his absence to bear testimony to those abilities, that zeal and success, which have been so eminently displayed in bringing the Botanic Garden to its present state of perfection. This testimony, though so justly due to him, he himself could not have published, nor is it probable that he would have permitted any one else to have done it. I hope, however, that he will pardon a friend for attempting it in his absence; and in doing it I shall almost exclusively confine myself to those facts which are furnished by the catalogue itself.

The utility of Botanical Gardens is so universally acknowledged, that there is now scarcely a civilized state in the world which has not one or more of them, and in many instances royal munificence has been called forth to encourage or support them. Many public bodies too have thought such an institution a necessary appendage. In Europe indeed almost every private nursery is a botanical garden, stored with plants from the different parts of the world; and, expensive as these establishments are, and great as the capitals are which are employed in them, the public demand for their productions has been found sufficient not only to cover the expenses, but also to secure a competency of wealth to their industrious owners.

By means of these Gardens, encouragement is given to the curious in all parts of the world to transmit seeds or plants to them, without fear of their being lost. All the treasures collected by travellers with so much labour, and at such great hazard and expense, are thereby preserved, and a laudable ambition to search with diligence for the hitherto unobserved or undescribed productions of nature is excited and maintained; a stronger proof of the truth of this observation can scarcely be desir-

ed than the unparalleled accession to our stock of plants from all parts of the globe within the last hundred years.

The Botanical Garden at Calcutta was begun by Colonel R. Kydd in March, 1786,* and collections of plants from different parts of the east were soon introduced into it with so much success, that, notwithstanding all the labour and difficulties which necessarily attend a first undertaking of this kind, the number of plants brought into it in the short space of eight years,† amounted to somewhat more than Three Hundred; all these are mentioned in the catalogue as introduced before the year 1794.

The unremitted attention paid by Dr. Roxburgh to the improvement of the garden, and his eminent abilities as a botanist, are far more fully exhibited in the following Catalogues than they could possibly be by any eulogium from a friend. The number of described species now in the garden amounts to about Three Thousand Five Hundred; for the knowledge of no fewer than One Thousand Five Hundred and Ten of which, as named and described in this catalogue, we are indebted to his indefatigable and discriminating researches. Among these are a great number of new genera, some of which have already a considerable number of species ascertained. To these should be added, those plants contained in the second catalogue, which, though not yet in the Botanical Garden, have been described, and many of them accurately drawn by him; these amount to Four Hundred and Fifty-Three. This, indeed, forms as important a part of his valuable labours, as the description of those already in that rich repository of Indian plants. The total number of plants drawn and described by him is One Thousand Nine Hundred and Sixty-Three.

The extensive correspondence which he has maintained, not only with gentlemen in all parts of India, but in most parts of the world, will be witnessed by the list of donors to the garden which he superintends; and the liberality with which he has al-

* A part of it had been previously cultivated as a private garden by Col. Kydd, but it was augmented, and became a public establishment at the time above-mentioned.

† Dr. Roxburgh came to it in the autumn of 1793, so that the time was not more than seven years and a half. Most of the grasses, and other indigenous plants which appear all over the country as weeds, it is presumed were in the Garden from its commencement: Dr. Roxburgh afterwards described them.

ways communicated to those who have applied to him, requires no other witness than the universal voice of the inhabitants of India, and even of many in Europe and America, to whom he has constantly and liberally sent supplies. In both these countries, several learned bodies have testified the high sense they entertain of his abilities by electing him one of their members.*

By the formation of Botanical Gardens the labours of scientific men have been called forth, and greatly aided. The great number of genera, the number of species belonging to each genus, and the numerous varieties of some species, thereby brought together, and placed in one point of view, afford facilities for the improvement of Botanical science which might have been sought in vain without the aid of such institutions. The labours of Tournefort, Linnæus, Smith, Jussien, Cavanilles, D'Herritier, Gærtner, Barton, and many others equally deserving of notice, would either never have made their appearance, or would have appeared in a form very different, and probably much inferior to that in which they are now published, had such institutions never existed.

But the Botanical Gardens and other noble collections of plants in Europe could never have been brought to the perfection in which they now appear, had not public or private repositories of plants been formed in the different settlements in Asia, Africa, and America. Hence their importance in all our foreign settlements is clearly established: for though it is freely confessed, that much has been accomplished by the ardour and enterprize of travellers and collectors; it will be obvious, on the slightest reflection, that the utmost which collectors of plants can do, must necessarily be very limited, unless there be some place in the countries where these researches are made, in which their collections may be deposited, and afterwards, in many instances, multiplied. A traveller, or a collector of plants, however diligent,

* Dr. Roxburgh has been chosen
1. Fellow of the American Philosophical Society of Philadelphia.
2. Corresponding Member of the Society for the Encouragement of Arts.
3. Fellow of the Royal Society of Physicians at Edinburgh.
4. Fellow of the Linnæan Society of London.
5. Fellow of the Linnæan Society of Philadelphia.
6. F. R. S. E.
7. Honorary Member of the Society of the Friends of Natural History at Berlin.

would collect plants in vain, unless he could keep them in safety till he had an opportunity of dispatching them to the place he wished; and seeds, bulbs, and other roots, would soon perish unless they could be sown or planted so as to furnish him with a stock when he had an opportunity of sending them to Europe. But to obviate these, and many other inconveniences, Botanical Gardens in our foreign possessions offer an easy remedy.

This, however, is not the only advantage arising from these institutions: timber trees, however valuable and important, can scarcely be propagated to advantage in countries differing much in temperature from those where they are indigenous: the Quercus robur will not grow in an Indian forest; nor can the Tectona grandis be brought to perfection in an English stove. Yet the cultivation of timber is in every country an object of the first importance. Here, then, is another instance of the utility of public gardens where these trees may be introduced and cultivated; their seeds perfected, and their plants multiplied, and where experiments may be made upon the soil most congenial to their nature, and the mode of culture which suits them best, and, also, upon the diseases and other causes of miscarriage to which they are incident.

The same reasoning will apply to fruit trees, and indeed to grain of every description. Grain, it is true, is only sown as an article of botanical curiosity in countries unfriendly to its growth; but where particular grains are cultivated, they are articles of the first importance, not only supplying nutriment for man and beast, but frequently furnishing a superabundance sufficient to ward off entirely, or greatly to alleviate, the miseries of other countries, and at all times furnishing a very important article of export trade.

It is readily granted, that Agricultural Societies would more effectually accomplish the object of improving the culture of grain, and that the premiums bestowed by them upon successful candidates in the various branches of husbandry, would operate with greater success, where these operations must necessarily be carried forward on a large scale, and a comparative view made of the advantage attending different modes of culture, and of the produce of different kinds of soil. This is, however, impracticable in many of our foreign possessions, and must necessarily be at-

tended with many defects in them all, which it is not here necessary to mention; the best substitute for such a society seems to be, the attaching of an Experimental Farm to a Botanical Garden, a measure which the liberal views of the government of British India have induced them to adopt, and which has been well repaid by the experiments made there on the cultivation and produce of different kinds of grain, on the strength and durability of the fibres of the bark of various plants, and upon various other subjects highly important to the agricultural interests of the country.

It is well known, that the virtues and excellence of Medicinal Drugs depend much upon the soil where they grow, and the manner of preparing them for the market. A sufficient quantity of medicinal plants for a full trial of their virtues, and for experiments on their culture and preparation, could scarcely be preserved any where except in a Botanical garden, where a large number of species may be collected, and carefully examined in all their different states. Some species, especially of the scitamineæ, are so much alike, that it is to be feared valuable drugs have fallen into disrepute through persons ignorantly confounding distinct species with one another and that fatal accidents may have been occasioned by the want of that discrimination, which botanical knowledge alone can supply. It is also highly probable, that many drugs of acknowledged value may be produced from a great number of plants differing widely from each other. This is known to be the case with starch, which is produced from a great number of farinaceous seeds, and even from some roots. Arrow root, formerly supposed to be obtainable from no plant but the West Indian Maranta arundinacea, is now known to be produced in great abundance from the tubers of several species of Indian Curcuma. The same may be observed of Caoutchouc or Indian rubber, which can be obtained from more East Indian plants than one, and very abundantly from the Ficus elastica.

A Botanical garden gives the same facilities for experiments upon the various Colouring Plants, and upon others employed in dying as Mordants; or upon such as are used in the arts in any other way. I forbear to mention the great value of large collections of esculent plants, and of others which are employed in the domestic

œconomy of the inhabitants of a country, or which by their beauty contribute to the enjoyment and pleasure of mankind.

But another very important advantage of an institution of this kind is, the propagating of useful plants, natives of other countries, and the consequent enriching of the country into which they are introduced. This has been done to a very large extent in England, which is now adorned with, and reaps the advantage of, a very large stock of vegetables originally imported from foreign countries. But many of these, which are mere objects of curiosity in England, are of the greatest utility when transported to a country congenial to their constitutions. The Mahogany tree, for instance, which but a few years ago was brought from Jamaica to this country, thrives so luxuriantly in Bengal, that many thousand trees of it are growing here, and even small pieces of furniture have been already made of the wood. The Pimento and Coffee prosper now as well in Bengal as in their native soil, and the Nutmeg, notwithstanding the climate is somewhat too cold for it, already produces fruit every year.

It is a most gratifying circumstance that our present Catalogue of Indian plants contains eight hundred and fifty-eight species of trees, some of which produce the best kinds of Timber in the world. The Teak (Tectona grandis) the Shal*a*, (Shorea robusta) and the Sissoo, (Dalbergia Sissoo) have long been well known, and deservedly esteemed. The wood of the jack tree (Artocarpus integrifolius) and the Toon, (Cedrela Toona) are but little inferior to mahogany. A good judge indeed would scarcely distinguish between the first of these and the finest mahogany, after both have been a little time in use. The Sit-shal*a* is universally employed to make furniture, and there are but few kinds of woods better adapted for that purpose.

The present catalogues furnish us with no fewer than eighteen species of Indian oak, some of which appear to be nearly or quite equal to English oak; ten species of Dalbergia, which are useful timber trees. Several mimosas also, and a great number of other species, appear for the first time among the number of prime timber trees, furnishing good materials for Ship building, Architecture, Furniture, and Domestic convenience.

The number of Medicinal Plants which it contains, is also very large. Most of the scitamineæ, by their sensible qualities, encourage us to expect from them considerable efficacy as medicines, not to mention that some of them, as Ginger, Turmeric, and Cardamoms, have been long in use both for medicinal and culinary purposes. It will appear that nearly all the plants of this description mentioned in the catalogue, have been but lately discovered, and placed in the Botanical garden; it must still be the labour of years to ascertain all their useful properties, and to apply them to the various purposes for which they are fitted. Had they never been collected, however, their virtues and uses would have remained wholly unknown, and all the notice we could have obtained of them must have depended wholly upon the uncertain, and often erroneous, information which might be accidentally obtained from the natives, whose total want of inquisitiveness, and discrimination, frequently occasion their confounding species with each other, which are entirely distinct.

The facility with which either public or private gardens in Europe may be abundantly supplied with plants, is obvious. The nurseries furnish every plant in abundance which has been cultivated for any length of time; and hence a single month would enable a person of property to procure a very extensive collection. But in this country the case is widely different: when plants or seeds, usually cultivated in Europe, are sent to this country, scarcely one in ten of them, on an average, arrives in a living or vegetating state, and of those which grow, perhaps scarcely one in ten succeeds, often because the whole stock is confined to a single plant, the loss of which, if it dies, cannot be repaired; and sometimes, because the seeds arrive at a time improper for sowing them. In other instances, the climate is unfavourable to them, and very frequently the mode of culture proper for these plants when removed to a hot climate, is unknown: little dependance can therefore be placed on supplies from England. But among the Natives of the East, there are few who ever think of the productions of nature: the whole number of plants enumerated in their most approved vocabularies, and books on the Materia Medica, does not exceed four or five hundred. Indeed, it often appears a matter of surprize to them, that any person should employ himself in what is to them so useless and unentertaining a pursuit as the study of Nature. Europeans are in general so circumstanced however, that little can be expected from them unless they have a strong inclina-

tion to Botanical studies. But they are still the only persons who are likely to contribute to an institution of this nature; and when it is considered, that to collect plants, and send them to the distance of several hundred miles, is very expensive, we shall cease to wonder that the number contributed from this quarter has not been greater. Indeed a total of three thousand two hundred species contributed to the Botanical Garden in the space of twenty years, three thousand of which at least have been given by the Europeans resident in India, not only vindicates them from the charge of indifference to this object, but evinces a degree of attention to it scarcely paralleled in an equal population in any other country.

A long series of years must, however, elapse before the catalogue of a Botanical Garden which depends on such resources alone, can be very extensive, however liberal the supplies may be. Still it will have the merit of presenting to the public a list of plants the far greater part of which were unknown to European Botanists, and therefore, though not so rich in the number of genera and species as those in a situation where plants can be easily purchased, it will far more extensively serve the cause of science, and thus more effectually answer the end of such an institution, than a large number of gardens, which, however rich, could furnish the Botanist with little of what is new.

Those who think superficially upon this subject, may suppose, that the stock of Asiatic productions is now nearly exhausted. This is so far from being the fact, that it may be considered as but just opened. The eastern frontier of Bengal has, in the short space of four years, produced one hundred and ninety species, all collected by one man, Mr. M. R. Smith, of Silhet, and he far from being in such opulent circumstances as will permit him to expend much in these pursuits. The success with which he has led the way, is sufficient to warrant very enlarged expectations when the whole of that extensive region shall be carefully examined. When to this we add, the whole of our northern frontier, reaching from the Garrow mountains to Cashmere, and reflect, that a very great part of it borders on the most elevated range of mountains in the world, we may surely indulge sanguine expectations of large additions to our botanical treasures without danger of disappointment.

To the extensive regions of forest-land on our borders, we must add the vast tracts of forest and mountain in every part of Hindoost'han; and with these were we to unite the countries east of Bengal, and the Islands scattered about the Indian Ocean, if the number of plants yet undiscovered bears any proportion to those discovered within the last twenty years upon any given area of country in India, and indeed actually conveyed to the Botanical Garden at Calcutta, the sum total will almost exceed the bounds of credibility.

We are not, however, to conclude that all the plants even of those parts of Bengal which are the best known, are as yet described, or have even been observed. To effect this in any place, a man must be resident on the spot, and be in the habit of constantly examining the fields, the woods, and the different receptacles of water: he must let no opportunity escape him, but collect under his own eye every thing he finds: this however requires such persevering diligence, and attention, that it can never be expected to be universal. A very great part therefore of this large field must remain partially explored for a long time to come: although a more complete knowledge will not cease to be matter of earnest desire to every lover of Botanical science.

Were Gentlemen who reside in the different parts of India to appropriate a small spot of ground for the reception of the plants which grow wild in their neighbourhood, and in their occasional excursions to collect those which they find, without overlooking even the most common or despised, and were they then to transmit them, as opportunity offered, to the Botanical Garden, or even a share of them, the collective total furnished by this simple expedient, would soon be very great. But as it would be folly to expect that every European, who resides in India, or even the greater part of them, should have such a taste for the study of nature as to adopt this plan, the utmost we can hope is, that those who are desirous of contributing to the progress of one of the most rational of studies, may adopt some regular method, which should make their labours useful to the Botanical world.

Opportunities of conveying plants from the distant parts of India to Calcutta can occur but seldom, and when they offer, even to those most inclined to communi-

cate of their vegetable treasures, the number of plants sent must be very few if they are then to be sought for in the woods; and it is probable that of these scarcely one in ten would arrive in a growing state; not to mention that of the numerous species to be found in any given district, a very few only could be in a state sufficiently conspicuous to be found at any one season of the year; plants not in flower being generally overlooked: to which may be added, that the labour of searching through an extensive district for plants, is too great to be carried on to any advantage upon the mere spur of the moment, and if undertaken, would be very imperfectly done by the natives of Hindoost'han, who must be the persons employed to collect and bring home the plants.

But on the other hand, were any gentleman to appropriate, as above suggested, a small spot for the reception of plants growing in his neighbourhood, it would be easy, when riding about the country, to make some particular mark, or set up some sign by any plants he might discover, or to point them out to his attendant; and on his return to send a man or two to fetch and plant them in his nursery. By this means the number would soon be very considerable; and they would be always at hand when an opportunity of sending them might occur; and as the person who makes such a collection would naturally bring several plants of the same species into this nursery, he would always have it in his power to replace any that might perish on the passage, or even after being received at the Botanical Garden.

It might render this mode more effectual, and at the same time more agreeable to the collector, were he to make a catalogue of the plants collected, without giving them names (unless he were a scientific botanist) only numbering them No. 1, 2, 3, 4, &c.; and writing opposite to the number the vernacular name, which in most instances should be done by a native in the character proper to the language. The name of the place where the plant was discovered might make another column, and the kind of soil in which it grows a third; to which might be added, remarks upon the plant, either collected from the natives, or made by the person himself from his own observation.

The more effectually to secure success, these plants should be placed in boxes,

or large earthen vessels, some time before-hand, that they might have time to strike root firmly; which would in many instances prevent their perishing on their passage. Seeds, especially of annual plants, should be collected when ripe, and the season of the year specified when they spring up, and when they flower. Seeds may always be sent by post at a small expence.

Many of the plants which are indigenous in the northern and more elevated parts of Hindoost'han will not grow if immediately removed to Bengal: this would probably be the case with most of those which are indigenous to the alpine regions of the Himalaya mountains; the Grape vine does not grow freely in Bengal, though it prospers well about Patna, the different species of Melastoma have not yet been preserved above two years in the neighbourhood of Calcutta, though they abound in the northern and eastern parts of Bengal. The Rubia munjistha grows plentifully among the northern mountains, but it can scarcely be kept alive near Calcutta. This is also the case with the Valeriana Jatamansi, and with many other valuable plants; even the Pines, and other timber trees of the northern mountains, are kept alive with difficulty, and are always small and stunted near Calcutta; but were those collected by any Gentleman of property and munificence and placed in a small garden near Hurdwar, or even as far south as Delhi; and were those nearest the eastern frontier, in a similar manner, collected and placed in a garden at Sylhet, or even at Dhaka, it is probable they would prosper, and that they might in a moderate time be inured to the climate of the lower parts of Bengal.

Though desirable, it is not absolutely necessary, that the gentleman who might form such a garden should be a scientific botanist; he might constantly communicate every thing to the superintendant of the Botanical Garden at Calcutta, either in a living state, or, where that was difficult, by good dried specimens, or when practicable, both ways. This would ensure the propagation of all those which could be naturalized there, while these Gardens would be large nurseries for the preservation of those plants which will not thrive in the lower parts of Bengal, and for the supplying and enriching of the Company's Botanical Garden, and even the private collections of Gentlemen desirous to promote their culture.

Something of the same nature in each of the Islands would be desirable, to secure collections there, and to preserve the plants collected, that duplicates, and even triplicates might be sent in succession to the Botanical Garden at Calcutta, without which they cannot be expected, in general, to succeed; but which would secure their naturalization in Bengal if either so useful or beautiful as to make that desirable; and would greatly promote Botanical knowledge, by adding to our present catalogues the greatest part of the undiscovered riches of the vegetatable kingdom in the Eastern part of the World.

In examining the catalogue, several errors in the terminations of the specific names have been detected. These arose from their having been removed from one genus to another, without the requisite change of the Gender being made. The copy from which the catalogue was printed was made out by a native writer, and these alterations were forgotten. The greatest part of it was printed off before this was discovered; these errors therefore, and a few other Typographical ones which have been observed, the scientific reader is requested to correct.

June 4, 1814.

W. CAREY.

CLASSIS I.

MONANDRIA MONOGYNIA.

Systematic names.	*Synonyma & Remarks.*	*Native place.*	*Donors and time, &c.*	*Duration & habit.*	*Time of flowering.*	*Time of seed.*
CANNA.						
Indica red.	B. *Surba-juya*.1		Before 1794.	H. E.		
yellow.	B. *Surba-juya*.2		Before 1794.	H. E.		
glauca.		America.	W. Hamilton, Esq. 1799.	H. E.	R. S.	C. S.
PHRYNIUM.						
dichotomum, R.	H. Seetul-patee.3	India, &c.	1795.	♄ E.	H. S.	7---8
virgatum, R.		Malabar.	Dr. J. Anderson, 1807.	♃ E.	R. S.	C. S.
spicatum, R.		Pegu.	Mr. F. Carey, 1809.	♃ E.	R. S.	
imbricatum, R.		Chittagong.	Mr. J. R. 1810.	♃ E.	R. S.	R. S.
parviflorum, R.		Bengal.	Dr. F. Buchanan, 1809.	♃ E.	R. S.	
capitatum.	B. *Kudalee*.4	Bengal.	Dr. F. Buchanan, 1796.	♃ E.	R. S.	C. S.
HEDYCHIUM.						
coronarium.	B. *Doolala-chumpa*.5	India.	Before 1794.	H. E.	R. S.	C. S.
flavum, R.	B. *Kattea-tilook-seer*.	Silhet.	Mr. M. R. Smith, 1810.	H. E.	R. S.	
coccineum.		Bengal.	Dr. F. Buchanan, 1809.	H. E.		
angustifolium, R.	B. *Bhooi-ada*.	Silhet.	Mr. M. R. Smith, 1810.	H. E.	5---6	C. S.
gracile, R.	B. *Kattea* laffear.	Silhet.	Mr. M. R. Smith, 1811.	H. E.	R. S.	
KÆMPFERIA.						
Galanga.	B. *Chundra-moola*.6	India.	Before 1794.	H. E.	R. S.	C. S.
rotunda.	B. *Bhooi-chumpa*.7	India.	Before 1794.	H. E.	H. S.	
angustifolia, R.	B. *Madum nirbish*.8	Bengal.	Kurrim-khan, 1796.	H.	R. S.	
pandurata, R.9		Sumatra.	Mr. W. R. 1803.	H. E.	R. S.	
ovalifolia, R.		Malacca.	Major Farquhar, 1810.	H. E.	R. S.	
CURCUMA.						
SECT. I.						
Spikes lateral.						
Zedoaria, R.	H. & B. *Bun-huldi*.10	India.	Before 1794.	H. E.	H. S.	6
Zerumbet, R.	H. *Kakhur*.11	India.	Dr. F. Buchanan, 1798.	H. E.	H. S.	
zanthorhiza, R.		Amboyna.	Mr. C. Smith, 1798.	H. E.	R. S.	
elata, R.				H. E.	R. S.	
cæsia, R.	B. *Neel-kanth'ha*.12	Bengal.	Dr. W. Carey, 1808.	H. E.	H. S.	
aeruginosa, R.		Pegu.	Mr. F. Carey, 1808.	H. E.	H. S.	
ferruginea, R.		Bengal.		H. E.	H. S.	
rubescens, R.		Bengal.		H. E.	H. S.	
comosa, R.		Pegu.	Mr. F. Carey, 1809.	H. E.	H. S.	
leucorhiza, R.13		Bengal.	I. Glass, Esq. 1804.	H. E.	R. S.	
angustifolia, R.	H. Teekor.14	Hindoostan.	H. Colebrooke, Esq. 1802.	H. E.	R. S.	8---9
SECT. II.						
Spikes central.						
longa.	B. H. *Huldi*.15	S. of Asia.	Before 1794.	H. E.	R. S.	
Amada, R.	B. *Amada*.	Bengal.	Dr. W. Carey, 1803.	H. E.	R. S.	
viridiflora, R.	M-n. Tommon-giring.	Sumatra.	Dr. C. Campbell, 1801.	H. E.	R. S.	
montana, R.		Coromandel.		H. E.	R. S.	
reclinata, R.	H. B. *Amba-huldi*.	Hindoostan.	H. Colebrooke, Esq. 1802.	H. E.	R. S.	
tordifolia, R.		Pegu.	Mr. F. Carey, 1809.	H. E.	R. S.	
AMOMUM.						
cardamomum.16						
angustifolium.		Madagascar.	Captain Tennant, 1802.	♃ E.	C. S.	
aculeatum, R.	M-n. Globba-durioen.	Moluccas.	Mr. W. R. 1803.	♃ E.	R. S.	9---10
maximum, R.	M-n. Globba.	Moluccas.	Before 1794.	♃ E.	H. S.	
dealbatum, R.		Chittagong.	Mr. J. R. 1810.	♃ E.	H. S.	C. S.
subulatum, R.17	B. *Bungali-elachee*.	Bengal.	B. Smith, Esq. 1810.	♃ E.	H. S.	
aromaticum, R.	H. *Morung-elachee*.	Morung, &c.	Mr. M. R. Smith, 1811.	♃ E.	H. S.	9
sericeum, R.	B. *Dooi-kesha*.	Garrow hills.	Mr. M. R. Smith, 1812.	♃ E.	R. S.	
ZINGIBER.						
I. *Spikes radical.*						
officinalis.18	H. & B. Ada.	India, &c.	Before 1794.	H. E.	R. S.	
Zerumbet.19	H. Buch.	India.	Before 1794.	H. E.	R. S.	10

1 S. *Vana-keli*. H. Lal-*subjuya*. T. *Kristmatamara*. 2 H. *Subjuya*. 3 S. *Seetula-putrika*. H. A. 4. t. 7. 4 H. M. 11. t. 34. 5. 3 var. 6 S. *Chundra-moolika*. 7 S. *Bhoomi-chumpuka*. 8 H. *Kanjan-boora*. 9 H. A. 5. t. 69. f. 1. H. M. 11. t. 10. 10 A. & P. Judwar. 11 S. *Karkhura*. B. Shoothee. A. Zerambad. 12 or *Kala-huldi*. 13 *From the tubers of this and the next following the edible fecula called Tikur (Arrow-root) is prepared.* 14 or Ticor. 15 S. *Huridra*. A. Urukussup. 16 H. A. 5. t. 65. f. 1. 17 Bengal, or Morung Cardamum. 18 S. *Ardruka*. B. Adruk, &c. A. Zenjabil. 19 or Muhaburee-buch. H. M. 11. t. 13.

Cassumunar,1 R.	H. & B. Bun-ada.	Hindoostan.	Before 1794.	H. E.	R. S.	9, 10, 11
ligulatum, R.		Hindoostan.	Colonel Hardwick, 1803.	H. E.	R. S.	
rubens, R.		Bengal.	Dr. F. Buchanan, 1808.	H. E.	R. S.	
squarrosum, R.		Pegu.	Mr. F. Carey, 1803.	H. E.	R. S.	
panduratum, R.		Pegu.	Mr. F. Carey, 1808.	H. E.	R. S.	
II. *Spikes terminal.*						
capitatum, R.		Hindoostan.	Colonel Hardwick, 1802.	H. E.	R. S.	9---10
elatum, R.		Bengal.	Captain B. Blake, 1810.	H. E.	7---8	9---10
COSTUS.						
speciosus.2	H. & B. Keoo.	India, &c.	Before 1794.	H. E.	R. S.	9---10
ALPINIA.						
I. *Inflorescence terminal.*						
Galanga.3	H. Kulinjan.	Malay Isl.	Dr. C. Campbell, 1800.	♂ or ☿ E.	H. S.	11
Allughas.4	B. Taruka.	Bengal.	Before 1794.	♂ or ☿ E.	R. S.	10---11
bracteata, R.		Bengal.	Mr. W. R. 1802.	♂ E.	H. S.	
malaccensis.5	[chumpa.	Chittagong.	Mr. W. Maddern, 1804.	♂ or ☿ E.	H. S.	8---10
nutans.	B. H. Poora-nag-	Bengal.	Dr. W. Carey, 1800.	☿ E.	H. S.	9---10
mutica, R.		P. of W. Isl.	Mr. W. R. 1802.	☿ E.	H. S.	R. S.
spicata, R.		Sumatra.	Mr. W. R. 1803.	♂ E.	H. S.	
calcarata.		China.	Captain Garnault, 1799.	☿ E.	H. & R. S.	
II. *Inflorescence radical.*						
Cardamomum,6 R.	H. & B. Elachee.	Malabar.	Mr. W. Dyer, 1809.	☿ E.		
C. medium, R.		Garrow hills.	Mr. M. R. Smith, 1811.	☿ E.	H. S.	
punicea, R.		Sumatra.	Dr. C. Campbell, 1804.	☿ E.	R. S.	
linguiforme, R.		Bengal.	1807.	☿ E.	R. S.	
GLOBBA.						
marantina.7		Moluccas.	Accident, 1798.	H. E.	R. S.	
orixensis, R.		India.	Dr. F. Buchanan, 1808.	H. E.	R. S.	
Careyana, R.		Pegu.	Mr. F. Carey, 1810.	H. E.	R. S.	
subulata, R.		Chittagong.	Mr. W. R. 1802.	H. E.	H. S.	H. S.
spatulata, R.		Silhet.	Mr. M. R. Smith, 1811.	H. E.	H. S.	H. S.
LOPEZIA.						
mexicana.		S. America.	W. Hamilton, Esq. 1809.	⊙ E.	C. S.	C. S.
BOERHAAVIA.						
procumbens.8	B. Gadha-poorna.	India.	Indigenous.	H. D.	○	○
repanda.		St. Helena.	Mr. Burchell, 1809.	☿ D.	C. S.	H. S.
MITHRIDATEA.						
quadrifida.		Mauritius.	Monsieur Jannet, 1810.	T.		

MONANDRIA DIGYNIA.

BLITUM.						
virgatum.		Europe.	Dr. Hornmann, 1807.	⊙	H. S.	

1 T. Kar-pooshpa. 2 S. Kemboo. H. M. 11. t. 8. 3 S. Koolanjuna. A. Kholinjan.
4 S. Taruka. H. M. 11. t. 14. 5 H. A. 5. t. 71. f. 1. 6 S. Ela. Can. Elum-chedi. 7 H. A. 5. t. 61. f. 2.
8 S. Shothughnee. The red variety.

CLASSIS II.

DIANDRIA MONOGYNIA.

NYCTANTHES.						
arbor-tristis.	B Shiooli.1	India.	Before 1794.	S. T.	9---10	C. S.
JASMINUM.						
I. *Leaves simple.*						
Zambac, sing. or wild.	H. Bun-mullika.2	India.	Before 1794.	♄ Sc.	H. S.	R. S.
double.3	S. B. Mullika.	India.	Before 1794.		H. S.	R. S.
rose.	B. Moota-bela.4	India.	Before 1794.		H. S.	R. S.
scandens.		India.	Mr. Maddern, 1803.	♄ Sc.	C. & H.S.	R. S.
elongatum.		Hidgelee.	Mr. Chapman.	♄ Sc.	D. S.	R. S.
lanceofolium, R.		Silhet.	Mr. M. R. Smith, 1810.	♄ Sc.		
pubescens.	S. B. Koonda.5		Before 1794.	♄ Sc.	☋	
bracteatum, R.		Sumatra.	Dr. C. Campbell, 1803.	♄ Sc.	C. S.	
latifolium.	T. Udivi-mulli.	India.	1799.	S. Tw.	H. S.	
arborescens.	B. Bura-koonda.	Hindoostan.	Col. A. Kyd, 1799.	S. Tw.	H. S.	
angustifolium.6						
simplicifolium.		S. Sea Isl.	1799.			
trinerve.		Silhet.	Mr. M. R. Smith, 1810.	♄ Sc.	H. S.	R. S.
coarctatum, R.		Chittagong.	Mr. J. R. 1811.	♄	H. S.	
laurifolium, R.		Silhet.	Mr. M. R. Smith, 1811.	♄ Sc.	3---4	6---7
attenuatum, R.	Bun-nowaree.	Silhet.	Mr. M. R. Smith, 1811.	♄ Sc.		
II. *Leaves compound.*						
heterophyllum, R.		Napaul.	Dr. F. Buchanan, 1801.	♄ E.		
auriculatum.7	H. & B. Joothi.	India.	1796.	♄ D.	H. S.	
lanceolaria, R.	[pooshpika.	Silhet.	Mr. M. R. Smith, 1812.	♄ E.	R. S.	
chrysanthemum, R.	S. H. & B. Hema-	Napaul.	Mr. Gibbons, 1799.	♄ E.	☋	
grandiflorum.8	B. H. Jatee.	India.	Before 1794.	♄ Sc.	☋	
paniculatum, R.		China.	Mr. W. Kerr, 1812.	♄ E.	R. S.	
LIGUSTRUM.						
laurifolium, R.		Chittagong.	Mr. J. R. 1811.	♄ Sc.	4---5	
CHIONANTHUS.						
ramiflora, R.		Moluccas.	H. C. 1798.	T.	H. S.	10
dichotoma, R.		Coromandel.	Uncertain, 1803.	♄ E.	H. S.	8
PHILLYREA.						
robusta, R.	H. Bhooi-moora.	Silhet.	Mr. M. R. Smith, 1811.	L. T.	6---7	1---2
paniculata, R.		China.	Before 1794.	T.	H. S.	
SCHREBERA.						
swietenioides.9	T. Mookadi.	Coromandel.	1801.	T.	H. S.	
OLEA.						
dioeca, R.	B. Atta-jam.	Silhet.	Mr. M. R. Smith, 1811.	L. T.	3	7
europea.		S. of Europe.	Mr. W. Dick, 1800.	S. T.		
fragrans white.	Ch. Quai-fa.	China.	Before 1794.	♄	C. S.	
red.	Ch. Tan-quai-fa.	China.	Mr. W. Kerr, 1811.	♄		
MILLINGTONIA.						
simplicifolia, R.	B. Dunta-rungee.	Silhet.	Mr. M. R. Smith, 1811.	T.	2---3	7---8
pinnata, R.	B. Batiwa.	Silhet.	Mr. M. R. Smith, 1811.	T.	4	9
ERANTHEMUM.						
pulchellum, R.10	B. Kala-jatee.	India.	Before 1794.	♄ E.	C. S.	
barleriordes, R.		Andamans.	Before 1794.	S. ♄ E.	H. S.	
diantherum, R.		Moluccas.	H. C. 1798.	S. ♄ E.	R. & C. S.	
racemosum, R.		Moluccas.	H. C. 1798.	S. ♄ E.	C. S.	
suffruticosum, R.						
JUSTICIA.						
1. *Anthers simple.*						
Ecbolium.11	H. Oodoo-jatee.	India.	Before 1794.	S. ♄	☋	
vitellina, R.		Chittagong.	Mr. J. R. 1811.	S. ♄	2---3	6
thyrsiflora, R.		Bengal.	Dr. W. Carey, 1800.	♄	C. S.	4---5
alba, R.		Andamans.	Lieut. Stokoe, 1796.	S. ♄	H. S.	
picta.12		Asia.	Before 1794.			
paniculata.13	B. Kala-megh.	Coromandel.	Dr. Klien, 1803.	♂ E.	R. S.	C. S.
echioides.14		India.	Dr. Klien, 1803.	☉ E.	☋	☋
polysperma, R.	B. Choota-mecheta.	Bengal.	Indigenous.	☉ A.	C. S.	H. S.
acaulis.	T. Nella-meera.	Coromandel.	1800.	H. S.	H. & C. S.	

1 S. B. Shephalika. H. Har-Singahar. 2 S. Asphota. B. Moorga, & Mullika. H. M. 6. t. 55.
3 H. M. 6. t. 50. 4 or Patna-mullika. S. Suptula. H. M. 6. t. 51. 5 S. Maghya. B. Kath-bel.
H. M. 6. t. 54. 6 H. M. 6. t. 53. 7 or Jooi. S. Yoothika. 8 S. Malutee.
H. Chumbeli. H. M. 6. t. 52. 9 Tam. Mogalinga. 10 H. Gool-sham.
11 H. M. 2. t. 20. 12 H. M. 6. t. 60. 13 or Kulpa-nath. Muha-titta. T. Nella-vemoo.
Tam. Kreata. H. M. 9. t. 56. 14 H. M. 9. t. 46.

II. *Anthers double, corol bilabiate.*						
nasuta.1	B. Jooi-pana.	Silhet.	Before 1794.	♄		
lanceolaria, R.	H. Toogee.	Silhet.	Mr. M. R. Smith, 1812.	♄ E.	12---1	
speciosa, R.		Bengal.	Dr. W. Carey, 1803.	☿	C. S.	
Careyana, R.			Dr. W. Carey, 1812.	♄ E.	C. S.	H. S.
tinctoria, R.	B. Beta-rung.		1795.	♄	C. S.	
chinensis.		China.	Mr. W. Kerr, 1809.		R. & C. S.	
bicalyculata.	Nasa-bhaga.	Coromandel.	H. C. 1802.	☉ E.	D. S.	
III. *Anthers double, corols ringent.*						
Adhatoda.2	B. & H. Bakus.	India.	Indigenous.	E. ♄	C. S.	
decussata, R.		Pegu.	Mr. F. Carey, 1810.	♄ E.	H. S.	
gandarussa.3	B. Jugut-mudun.	India.	Before 1794.	♄ D.	C. S.	
betonica.4		Coromandel.	1803.	♄ E.	D. S.	D. S.
ramosissima, R.				♄ Cr.	D. S.	D. S.
tranquebariensis.			Dr. Rottler, 1802.	H. D.	☾	○
glabra, R.		India.	Nella-mandu.	H. E.	R. S.	
tomentosa, R.			Indigenous.	H.	C. S.	
peruviana.		Peru.	W. Hamilton, Esq. 1805.	S. ♄ E.	C. S.	
repens.		India.	Indigenous.	H. D.	○	○
procumbens.				H. D.	○	☾
pectinata.				H. D.	C. S.	
lamifolia, R.		Bengal.	Indigenous.	☉ E.	C. S.	
GRATIOLA.						
amara, R.5		Moluccas.	H. C. 1800.	C. S. ☉ D.	C. S.	
grandiflora.		India.	Indigenous.	S. ☉ Cr.	R. S.	C. S.
integrifolia, R.6		Bengal.		S. ☉ D.	R. S.	C. S.
lucida.7		India.		S. ☉ D.	R. S.	C. S.
veronicifolia.8		Bengal.		S. ☉ Cr.	R. S.	C. S.
juncea, R.		India.		S. ☉ E.	R. S.	C. S.
oppositifolia.				S. ☉ E.	R. S.	C. S.
Monnieria.9	H. Shwet-chamni.			☉ Cr.	R. S.	C. S.
reptans, R.		Moluccas.	Accident.	S. ☉ Cr.	R. S.	C. S.
serrata, R.		Bengal.	Indigenous.	S. ☉ E.	R. S.	C. S.
parviflora, R.		India.		S. ☉ E.	R. S.	C. S.
hyssopioides.				S. ☉ E.	R. S.	C. S.
racemosa, R.		Bengal.		S. ☉ E.	R. S.	C. S.
CALCEOLARIA.						
pinnata.		Peru.	Accident.	S. ☉ E.	C. S.	
UTRICULARIA.						
stellaria.		India.	Indigenous.	☉ A.	R. S.	C. S.
fasciculata, R.10	H. Jhanji.	Bengal.		☉ A.	R. S.	C. S.
diflora, R.	H. Chota-jhanji.			☉ A.	R. S.	C. S.
VERBENA.						
nodiflora.11	H. Bhooi-Okra.	India.		H. Cr.	☽	○
jamaicensis.12		W. Indies.	W. Hamilton, Esq.	♂ D.	☽	○
bonariensis.		Buenos ayres.	1798.	♂ E.	☽	○
officinalis.		Napaul.	Dr. F. Buchanan, 1802.	☉ E.	☾	☾
LYCOPUS.						
dianthera, R.		Napaul.	Dr. F. Buchanan, 1802.	☉ E.	H. S.	H. S.
MONARDA.						
punctata.		N. America.	W. Hamilton, Esq. 1802.	☉ E.	D. S.	D. S.
SALVIA.						
bengalensis, R.		Bengal.	Before 1793.	♄ E.	R. S.	C. S.
coccinea.13		N. America.	W. Hamilton, Esq. 1797.	☉ E.	☾	☾
hispanica.		Europe.	A. B. Lambert, Esq.	☉ E.	C. S.	C. S.
brachiata, R.14	B. Bhoo-toolsi.	Bengal.	Indigenous.	☉ E.	C. S.	C. S.
lyrata.		America.	W. Hamilton, Esq. 1790.	H.		
syriaca,		Syria.	W. Hamilton, Esq. 1790.	H.	C. S.	
officinalis.	Sage.		Mr. Mc Mahon, 1809.			
parviflora, R.	B. Koka-buradi.	Silhet.	Mr. M. R. Smith, 1811.	☉ E.	R. S.	C. S
FRAXINUS.						
chinensis.		China.	Before 1794.	S. T.	H. S.	

DIANDRIA TRIGYNIA.

PIPER.						
nigrum.15	H. Gol-mirch.	Malabar, &c.	Before 1793.	♄ R.		
trioecum, R.	T. Mirial-tiga.	Circars.	1793.	♄ R.	R. & C. S.	
longum.16	H. Pippul.	Bengal.	Before 1793.	♄ R.	R. & C. S.	
Betle.17	H. & B. Pan.	India.	Before 1793.	♄ R.	R. & C. S.	
reptans.		Moluccas.	H. C. 1802.	♄ R.	R. & C. S.	
sarmentosum, R.	M-n. Gadu.	P. Pinang.	H. C. 1802.	♄ Sar.	R. S.	
Chuvya, R.18	H. & B. Chuee.	Backergunge.	Juy narayun Ghosal, 1808.	♄ R.		
peepuloides, R.19		Silhet.	Mr. M. R. Smith, 1811.	♄ R.		
silvaticum.20	H. Puhari pippul.		Mr. M. R. Smith, 1811.	♄ R.	R. S.	R. S.

1 T. & Tam. Naga-mulli. H. M. 9. t. 69. 2 S. B. Vasooka. 3 H. M. 9. t. 42. 4 H. M. 2. t. 21. 5 Karanga amara. Vahl. H. A. 5. t. 170. f. 1. 6 H. M. 9. t. 57. 7 H. A. 5. t. 170. f. 3. 8 H. A. 5. t. 170. f. 2. 9 B. Ada-beerna. 10 T. Natsoo. 11 S. Unkota. T. Nella-pippala. 12 Stachy tarpheta. Vahl. 13 S. Rosea. Vahl. 14 H. Bun-baburee. 15 S. B. Muricha. M-n. Ladda. 16 or Pilpul. S. B. Pippulee. T. Peepal. Pippulee-mool the root. 17 S. Tamboolee, Purna, &c. M-n. Siri. H. A. 5. t. 116. f. 2. 18 S. Chuvya 19 Silhet long Pepper. 20 i. e. mountain pippul.

CLASSIS III.

TRIANDRIA MONOGYNIA.

VALERIANA.	[shee.					
Jatamansa.	S. B. H. *Jutamang-*	Tibet, &c.	Hon. C. A. Bruce, 1796.	H. E.		
OLAX.						
scandens, R.	T. Turka-vepa.	Coromandel.		C. ♄ Sc.	H. S.	C. S.
imbricata, R.		Silhet.	Mr. J. R. 1810.	L. Sc. ♄	H. S.	
phœnicarpa, B.	B. *Moicha.*	Silhet.	Mr. M. R. Smith, 1812.	S. T.		5
LOEFLINGIA.						
indica.		India.	Indigenous.	☉ D.	D. S.	D. S.
HIPPOCRATEA.						
indica.		Coromandel.	1804.	C. Sc. ♄	H. S.	3
obtusifolia, R.			Dr. Berry, 1802.	Sc. ♄	H. S.	H. S.
arborea, R.	H. *Kotta-pahari*a.	Hindoostan.	Dr. W. Hunter, 1797.	S. T. D.	H. & R. S.	3---4
JOHNIA.						
salacioides, R.	B. *Goluk-laroo.*	Tipperah.	Mr. S. Harris, 1800.	♄	H. S.	R. S.
coromandeliana, R.		Coromandel.	Mr. Heshusius, 1796.	♄	C. S.	H. S.
CROCUS.						
vernus.		Europe.	Mr. M'Mahon, 1810.	H.		
GLADIOLUS.						
marginatus.		Cape of G. H.	H. Hope, Esq. 1810.	♃ E.	H. S.	
IRIS.						
florentina.		Europe.	From the Cape, 1799.	H.	3---4	
chinensis.1		China.	1806.	H.	2---3	
susiana.		Levant.	Dr. W. Carey, 1811.	H.	H. S.	
Xiphium.		Europe.	Dr. W. Carey, 1812.	H.		
MOREA.						
chinensis.	H. *Dosbi-chundi.*	India.	Before 1794.	H. E.	R. S.	R. S.
MARICA.						
paludosa.		Guiana.	W. Hamilton, Esq. 1801.	H.	R. S.	R. S.
COMMELINA.						
communis.	B. *Juta-kanchura.*	India, &c.	Indigenous.	☉ Cr.	R. S.	R. S.
bengalensis.3	B. *Kanchura.*			☉ Cr.	R. S.	R. S.
salicifolia, R.4	B. *Panee-kanchura.*	Bengal.		☉ Cr.	R. S.	R. S.
nana, R.				☉ Cr.	R. S.	C. S.
nudiflora.5	B. *Kanduli.*	India.		☉ Cr.	R. S.	R. S.
cespitosa, R.		Chittagong.	Mr. W. R. 1801.	☉ Cr.	R. S.	C. S.
scapiflora, R.	H. *Kurelli.*	Hindoostan.	Col. Hardwick, 1802.	H. E.	R. S.	R. S.
herbacea, R.		Sumatra.	Dr. C. Campbell, 1804.	H. E.	☾	☾
erecta.		N. America.	W. Hamilton, Esq. 1805.		R. S.	
virginica.		N. America.	W. Hamilton, Esq. 1804.		R. S.	C. S.
SONERILLA.						
maculata, R.	B. *Swot.*	Garrow hills.	Mr. M. R. Smith, 1811.	S. ♂	R. S.	C. S.
emaculata, R.						
angustifolia, R.						
XYRIS.						
indica.6		India.	Indigenous.	☉ E.	C. S.	H. S.
KYLLINGA.						
monocephala.7	B. *Shwet gothoobi.*	India.	Indigenous.		R. S.	
umellata.8	B. *Bura gothoobi.*				R. S.	
triceps.9					R. S	
CYPERUS.						
semi-nudus, R.10	H. *Goola-meth'hi.*					
dubius.	B. *chota gothoobi.*					
monocephalus, R.	B. gotoobi.					
punctatus, R.						
compressus.	B. H. *chooncha.*11					
diformis.	B. *Behoora.*					
cruentus.						
incurvatus, R.						
tortuosus, R.						
rotundus.	B. & H. *Mootha.*12					
pertenuis, R.	B. *Nagur-mootha.*					
tuberosus.						
tenuiflorus.	T. *Ghelaba-tunga.*					
Iria.	B. & H. *Bura-choocha.*					

1 B. M. 373. 2 H. M. 11. t. 37. B. *Dusha-bahoo-chundi.* 3 H. *Kanuraca.* 4 S. Langulee.
5 B. *Kendru.* H. M. 9. t. 63. 6 H. M. 9. t. 71. 7 B. *Nirbisee.* 8 H. M. 12. t. 63.
9 H. M. 12. t. 52. 10 T. *Ghœdoo-tunga.* 11 *Salli-tunga.* 12 A. *Sud.* S. *Moosta.*

inundatus, R.	B. & H. Patee.
Pangorii.	B. & H. Chumati-patee.
procerus.1	
spinulosus, R.	
elatus.	B. Gol-mulunga.2
verticillatus.	B. Burethi.
distans.	B. Panee-mulunga.
tegetus, R.	B. & H. Madoor-kati.
tegetiformis, R.	
alopecuroides.3	
Haspan.	

SCIRPUS.

I. *Leafless.*

tuberosus, R.	
plantagineus.	B. & H. chechka.
tumidus, R.	
incurvatus, R.	
articulatus.	H. Put-puti-chetchka.
mucronatus.	
supinus.	
pentagonus, R.	

II. *Leafy.*

monostachyos.	
strobelinus, R.	
bispicatus, R.	B. Kesuri-mulunga.4
schœnoides, R.	H. Moor-moori.
tristachyos.	
squarrosus.5	H. Goori.
arvensis.	
glomeratus.	B. Bura-nirbisi.
diphyllus.	B. Tal-nuroo.
miliaceus.	
complanatus.	B. Koratiya Juvani.
tetragonus, R.	B. Bura Juvani.
quinqueangularis.	B. Juvani.
anceps, R.	B. Joopi.
Kysoor, R.	B. & H. Kesooriya.6
grossus.	

TRIANDRIA DIGYNIA.

SACCHARUM.						
cylindricum.	B. Ooloo.7	India.	Indigenous.	H. E.	R. S.	C. S.
spontaneum.	B. Kashiya.8			H. E.	R.S.	C. S.
fuscum, R.	H. & B. Pata-khuree.			H. E.	4---5	R. S.
semidecumbens, R.	B. & H. Khuree.			♃	R. S.	C. S.
canaliculatum, R.		Bengal.		♃ E.	R. S.	C. S.
officinarum.9		India.	Before 1794.	♃ E.	R. S.	
chinensis, R.10		China.	H. C. 1795.		R. S.	
procerum, R.	B. Teng.	Bengal.	1796.	♃ E.	R. S.	
Sara, R.	S. B. & H. Shura.	Bengal.	Indigenous.		R. S.	C. S.
Munja, R.	H. Munja.11	Hindoostan.	Mr. Arbuthnot.	H. E.	R. S.	C. S.
ANTHISTIRIA.						
ciliata.		India.	Indigenous.	⊙ E.	C. S.	H. S.
polystachya, R.		Bengal.		⊙ E.	R. S.	C. S.
scandens, R.				♃ Sc.	C. S.	C. S.
heteroclita, R.				⊙ Cr.	C. S.	C. S.
prostrata, R.12		India.		⊙ Cr.	C. S.	C. S.
cymbaria, R.13				⊙ E.	C. S.	C S.
arundinacea, R.		Bengal.	Kurrim-khan, 1802.	♄ E.	C. S.	C. S.
ANDROPOGON.						
I. *Spikes terminal.*						
serratum.	B. Detara.14	Bengal.	Indigenous.	⊙ E.	C. S.	H. S.
contortum.	T. Yedda.	Coromandel.	Accident.	⊙ E.	C. S.	H. S.
tenellum, R.		Bengal.	Indigenous.	⊙ E.	C. S.	H. S.
binatum, R.				⊙ D.	C. S.	H. S.
conjugatum, R.		India.	Indigenous.	⊙ E.	C. S.	H. S.
filliformis, R.		Bengal.		⊙ E.	C. S.	H. S.
tristachyum, R.				⊙ E.	10---11	C. S.
lanceolatum, R.		Coromandel.		⊙ E.	C. S.	H. S.

1 H. M. 12. t. 50. 2 T. Rakesha. 3 H. M. 12. t. 42. 4 T. Kushi-tunga. 5 H. M. 12. t. 36. 6 Roots esculent. 7 T. Barom-hiss. H. A. 6. t. 7. f. 2. A. 8 H. Khugura. T. Billoo-gaddi. S. Kasha. 9 S. Ikshoo. B. Ik, Ak, Ookh, Pooree or straw-coloured cane. P. Nie-shukir. H. Kulooa a pale variety, and Kajooli a large red sort, S. Poondru. 10 Reckoned a richer cane than officinarum. 11 S. Munja. 12 Andropogon prostratus, Linn. mant. 304. 13 Andropogon cymbarius, Linn. mant. 303. 14 or Detto. S. Deva tara.

scandens, R.	H. Looari.	India.	Indigenous.	♂ Sc.	C. S.	H. S.
pertusum.				☉ E.	C. S.	H. S.
Bladhii.				☉ E.	C. S.	H. S.
II. *Panicles terminal, naked.*						
aciculatum.	H. & B. Chor-kanta.1			♂ Cr.	R. S.	10---11
punctatum, R.	B. Goaliya.			☉ E.	C. S.	H. S.
muricatum.	H. & B. Bena.2	India.	Indigenous.	H. E.	R.S.	C. S.
glabrum, R.	B. Gundha-goorana.	Bengal.		☉ E.	C. S.	H. S.
bicolor, R.3	B. Kala-debdhanya.	India.	Cultivated.	♂ E.	C. S.	H. S.
Sorghum, R. 3 var.4	H. Jooar.	India.	Cultivated.	♂ E.	C. S.	H. S.
saccharatum, R.5	H. Dedhan.			☉ E.	C. S.	H. S.
laxum.	H. Kula-moocha.6		Indigenous.	♂ E	C. S.	H. S.
miliaceum, R.		Hurdwar.	General Martin.	☉ E.	C. S.	H. S.
III. *Panicles foliaceous.*						
Schœnanthus.	H. Gundha-bela.7	India.	Cultivated.	♃ E.		
Nardus.	B. Korunkoosha.8	N. of Bengal.	General Martin.	♃ E.	R. S.	C. S.
Martinii, R.		Coromandel.	General Martin.	♃ E.	R. S.	C. S.
PASPALUM.						
scrobiculatum.9	T. Aruga.	India.	1802.	☉ E.	R. S.	R. S.
Kora.	H. Koda.10		Indigenous.	☉ P.	R. S.	R. S.
longiflorum.11				♃ Cr.	R. S.	R. S.
longifolium, R.		Sumatra.	Dr. C. Campbell, 1806.	☉ E.	R. S.	R. S.
stoloniferum.		Peru.	A. B. Lambert, Esq. 1802.		D, S.	R. S.
PANICUM.						
I. *Spikes simple.*						
barbatum, R.12		Malay Islands.	Accident, 1800.	♃ E.	R. S.	C. S.
spicatum, R.13	H. Bujura.	India.		☉ E.	R. S.	9---10
involucratum, R.			Indigenous.	☉ E.	R. S.	C. S.
glaucum.	H. Pingi-nachi.14			☉ E.	R. S.	C. S.
holcoides, R.	B. Swati.			☉ E.	R. S.	C. S.
interruptum.	H. Nardul.15			☉ A.	R. S.	C. S.
curvatum.				☉ E.	R. S.	C. S.
II. *Spikes fascicled.*		[&c.				
Dactylon.16	B. Doorba.	Asia, Europe,	Indigenous.	☉ Cr.	○	○
ægyptiacum.		India.		☉	R. S.	
ciliare.	B. & H. Makur-jali.17			☉ Cr.	R. S.	
filiforme.		China.	Mr. W. Kerr, 1809.	☉ Cr.	R. S.	
lineare.		Cape of G. H.	1799.	☉ Cr.	R. S.	
cimicinum.	[gaddi.	India.	Indigenous.	☉ E.	R. S.	
corymbosum, R.	T. Isheepa-bora	India.	Indigenous.	☉ A.	R. S.	
III. *Spikes alternate secund.*						
brizoides.	T. Woada.18	India.	Indigenous.	☉	R. S.	C S.
fluitans.	H. Peenar.			☉ A.	R. S.	C. S.
lanceolatum.		Bengal.		☉ Cr.	R. S.	C. S.
Burmanni.19				☉ Cr.	R. S.	
stagninum.	H. Dula.20	India.		☉ E.	R. S.	
Crus-corvi.				☉ Cr.	R. S.	
colonum, red.	H. Lal-shama.21			☉	R. S.	
green.	H. Shama.			☉	R. S.	
cuspidatum, R.		Bengal.		☉ D.	R. S.	
IV. *Spikes alternate, scattered.*						
setigerum.	H. Bura Jal-ganti.	India.		♃ Cr.	R.S.	
repens.				☉ Cr.	R.S.	
hirsutum, K.	H. Jal-ganti.22					
tomentosum, R.		[dia.		☉	R. S.	
verticillatum.	H. Dora biaroo.23	Europe & In-		☉	R. S.	
italicum.24*	H. & B. Kangni.		Before 1794.	☉ E.	R. S.	9
strictum, R.	B. Gang-bena.	Bengal.	1800.	☉ E	R. S.	
hispidulum.	B. Bura-shama.		Indigenous.	☉ E.	R.S.	
frumentaceum, R.25*	B. Shama.	India.		☉ E.	R.S.	
V. *Panicled.*						
tenellum, R.	H. Joopi-kunka.	India.		☉ D.	R. S.	

1 S. Chora-pushpa. H. M. 12. t. 43. 2 and the root kuskus. S. Veeruna. 3 H. Mukoo, Junar. Dedhan, or Kala-jooar. Holcus. Linn. and cultivated over various parts of India for its grain, which the natives eat. 4 T. Tella-janoo. B. Deb-dhanya. Holcus. Linn, &c. Next to rice, this may be said to be the most extensively cultivated of all the culmiferous tribe, and forms a very considerable part of the diet of the natives of the countries where it is reared. 5 Holcus. Linn. is also extensively cultivated in many parts. 6 T. Gadi-janoo. 7 M. Sirek. S. Bhoostrina. 8 H. Iwarankoosa. 9 H. Kodaka-chaul. S. Kudruva. In elevated tracts this is also cultivated for its grain, which the poor eat. 10 T. Neer-aruga. 11 H. M. 12. t. 44. 12 Compare with P. Polystachion. 13 T. Peddi-ganti. Holcus. spicatus. Linn, &c. four varieties are cultivated by the native farmers, who eat the grain, and feed their cattle with the straw. 14 T. Naka-kora. 15 T. Wolam. 16 S. Doorva. H. Doob. Tam. Arugam-pillee, &c. T. Gherka. Agrostis Linearis Linn. sp. pl. ed. Willd. 1. 375. Retz. obs. 4. no. 51. Asiat. res. 4. 249. It is reckoned the best grass in India for cattle. 17 T. Shangalli-gaddi. 18 P. flavidum and flaccidum, I consider varieties of this. 19 H. A. 6. t. 5. f. 3. 20 T. Pedda-woondoo. 21 S. Shamaka. T. Woondoo-gaddy. 22 T. Woondoo. 23 T. Chicklenta. 24 or Kangnee. T. Kora. S. Kungoo. 25 T. Bonta-shama.

* Of these there are several varieties, all cultivated by the native farmers for the grain, which are articles of diet, particularly amongst the poor.

serratutum, R.				⊙ E.	R. S.	
paludosum, R.	H. & B. *Kulusnar*.1			⊙ E.	R. S.	
uliginosum, R.	B. *Burunda*.2			⊙ A.	R. S.	
sarmentosum, R.		Sumatra.	Dr. C. Campbell, 1804.	♃ Cr.	R. S.	C. S.
miliare, 3. var.*	T. Nella-*shama*.	India.	Before 1794.	⊙ E.	R. S.	C. S.
miliaceum.3*	B. & H. Cheena.		1794.	⊙ E.	R. S.	C. S.
miliaceoides, R.				⊙ E.	R. S.	C. S.
plicatum, R.		Sumatra.	Dr. C. Campbell.	⊙ E.	R. S.	C. S.
nervosum, R.		Napaul.	Dr. F. Buchanan, 1802.	⊙ E.	R. S.	C. S.
costatum, R.		Mauritius.	Captain Tennant, 1802.	⊙ Cr.	R. S.	C. S.
PHLEUM.		[Napaul, &c.				
crinutum.		Hindoostan.	Col. Hardwicke, 1802.	⊙ D.	C. S.	H. S.
MILIUM.						
filiforme, R.	H. Kan-*koojooriya*.	Bengal.	Indigenous.	⊙ Cr.	R. S.	
sanguinale, R.	H. *Chiri-chira*.	India.	1802.	⊙ Cr.		
ramosum.		Bengal.		⊙ E.		
adscendens.		Napaul, &c.	Dr. F. Buchanan, 1802.		3	
pulcher, R.		China.	Mr. W. Kerr, 1809.			
AGROSTIS.						
linearis, K.4						
tenacissima.	B. Panee *Doorba*.5	India.	Indigenous.	⊙ Cr.	R. S.	
diandra.	B. Bena-*jon*i.	India.	Indigenous.	⊙ E.	R. S.	
ISCHŒMUM.						
aristatum.	B. Melancha.6	Bengal.	Gardeners.	⊙	C. S.	C. S.
rugosum.		Bengal.	Indigenous.	⊙	C. S.	C. S.
semisagittatum, R.		Bengal.	Indigenous.	⊙	C. S.	C. S.
conjugatum, R.		Bengal.	Indigenous.	⊙ Cr.	C. S.	C. S.
geniculatum, R.		Bengal.	Indigenous.	⊙ Cr.	C. S.	C. S.
cuspidatum, R.		Bengal.	Kurrim-khan.	♂ A.	C. S.	C. S.
APLUDA.						
aristata.	H. Goruma.7	India.	Indigenous.	♃ Sc.	C. S.	
geniculata, R.		Bengal.	Indigenous.	♃ Sc.	C. S.	
AIRA.						
filiforme, K.	B. Choota *pini-nuti*.	India.	Indigenous.	⊙ E.		
MELICA.						
refracta, R.		Amboyna.	Mr. C. Smith, 1798.	E.	C. S.	
diandra, R.		Bengal.	Indigenous.	⊙ E.	R. S.	
latifolia, R.		Garrow hills.	R. Kyd, Esq. 1810.	E.	3---4	R. S.
CHLORIS.						
barbata.		India.	Indigenous.	⊙	R. S.	C. S.
POA.						
chinensis.	B. *Bura-pini-nuti*.	Bengal.	Indigenous.	⊙ E.	R. S.	C.S.
cynosuriodes.	H. & B. *Koosha*.8	India.	Indigenous.	♃ E.	R. S.	C. S,
diandra, R.		Bengal.	Indigenous.	⊙ E.	C. S.	C. S.
nutans.	T. Urenka.	India.	Indigenous.	♂ E.	R. S.	C. S.
plumosa.9	H. *Puluka*.	India.	Indigenous.	⊙ E.	R. S.	C. S.
multiflora, R.		Bengal.	Indigenous.	⊙ E.	R. S.	C. S.
unioloides.	H. *Koni*.	Bengal.	Indigenous.	⊙ D.	C. S.	C. S.
gangetica, R.		Bengal.	Indigenous.	⊙ D.	C. S.	C. S.
annua.		England.	Accident, 1808---9.	⊙ D.	2---3	C. S.
angustifolia.		Europe.	Accident, 1808---9.	⊙ E.		
cylindrica, R.		China.	Mr. W. Kerr, 1811.	⊙ E.	R. & C.S.	C. S.
ELEUSINE.						
corocana.†	H. & B. *Marooa*.10	Asia.	Before 1794.	⊙ E.	R. & C.S.	C. S.
stricta, R.†	T. Pedasodi.11	Asia.		⊙ E.	R. & C.S.	C. S.
ægyptiaca.12	B. *Makur-jali*.	India.	Indigenous.	⊙ E.	○	○
indica.	H. *Mal-ankuri*.13	India.	Indigenous.	⊙ E.	○	○
verticillata, R.				⊙ E.		
calycina, R.		India.		⊙ E.		
ARUNDO.						
bifaria.	T. *Potoo-ederoo*.	Bengal, &c.	1800.	♃ E.	R. & C.S.	C. S.
karka.	H. & B. *Nul*.14	India.	1796.	♃ E.	R. & C.S.	C. S.
bengalensis.	B. *Gaba-nul*.	Bengal.	Indigenous.	♃ E.	R. & C.S.	C. S.
gigantea			W. Hamilton, Esq. 1809.	♃ E.	R. & C.S.	C. S.
MANISURUS.						
granularis.		Coromandel.	1807.	⊙ E.	R. S.	C. S.
ROTTBOELLA.						
glabra, R.	B. *Baksha*.	Bengal.	Indigenous.	Cr.		
compressa.	H. *Pan-shiru*.15	India.		D.		
exaltata.	B. *Bura Sooti*.16		Kurrim-khan.	⊙ E.		
perforata, R.	B. *Kurki*.17					

1 T. Soda. 2 T. Gundra. 3 S. *Unoo*. T. Worga. 4 See Panicum Dactylon.
5 T. Sema-gherka. 6 H. Munta. 7 B. *Choon khuriki*. 8 S. *Koosha*, Doorva, &c.
T. Durba or *Durpa*. 9 H. M. 12. t. 41. 10 T. *Sodi*. Can. *Raggi*. 11 or *Muddi-ruba*.
12 H. M. 12. t. 95. 13 T. *Kar-sodi*. 14 S. *Nula*. T. Naga-sara-ma-dantoo. 15 T. Sheroo-*panuku*.
16 T. *Kunda-punookoo*. 17 T. *Punookoo*.

* See note in the preceding page.

† Cynosurus Linn. Several varieties of these very useful plants are very generally cultivated during the rainy and cool seasons, on lands too much elevated for rice. In Malwa it is called Maud, or Mal. In the Carnatic Natcheney, see H. M. 12. t. 78.

AVENA.					
sativa.	Oats.		⊙ E.		
HORDEUM.					
hexastichon.1	B. Jub.		⊙ E.	C. S.	2---3
vulgare.2		A. B. Lambert, Esq.	⊙ E.	C. S.	2---3
TRITICUM.					
æstivum.3	B. Gom.		⊙ E.	C. S.	2---3
hybernum.					

TRIANDRIA TRIGYNIA.

MOLLUGO.					
pentaphylla.	B. Khet-papara.4	India.	Indigenous.	⊙ D. D. S.	H. S.
triphylla.	B. Jul-papara.			⊙ D. D. S.	H. S.
verticillata.				⊙D. D. S.	H. S.

1. S. Yuva. H. Jow. N. Tosa. A. She-eer. Very generally cultivated over Hindoostan during the cool season. 2 Spring Barley. 3 S. Godhooma. P. Gundum. H. Gow. A. Bur. Three or more varieties are very generally cultivated over the interior of Hindoostan between October and March. 4 T. Beri-shadrosi. S. Kshetra purputee.

CLASSIS IV.

TETRANDRIA MONOGYNIA.

EMBOTHRIUM.						
speciocissimum.	Waratah.	N. S. Wales.	Mrs. Blaxland, 1811.			
HEDYOTES.						
hispida.		Moluccas.	Accident, 1798.	⊙ D.	R. S.	
lineata, R.		Chittagong.	Mr. J. R. 1810.	⊙ D.	R. S.	R. S.
scandens, R.	B. *Guji.*	Silhet.	Mr. M. R. Smith, 1811.	♃ Sc.	H. S.	R. S.
SPERMACOCE.						
teres, R.		Mysore.	Dr. F. Buchanan, 1801.	♃ E.	R. S.	C. S.
exerta, R.		Coromandel.		♃ E.	9---10	12.
lævis, R.		Bengal.			R. S.	C. S.
costata, R.		Moluccas.	Accident.	♂ D.	○	○
semierecta, R.		Sumatra.		♂ D.	R. S.	C. S.
scabra.1		Coromandel.	Dr. Rottler, 1803.	⊙ D.	○	○
articularis.			1803.	⊙ D.	○	○
hispida.	T. Madana-burta-kada.		1803.	⊙ D.	○	○
HYDROPHYLAX.						
maritima.		Coromandel.	Dr. Klein, 1800.	♃ Cr.	○	○
RUBIA.						
Munjista, R.	H. *Munjit.*2	Boutan, &c.	Mr. Arbuthnot, 1801.	Sc.		
IXORA.						
coccinea.3		India.	1800.	♄ E.	○	○
Bandhuca, R.	B. *Rungun.*4	Bengal.	Before 1794.	♄ D.	○	C. S.
fulgens, R.5		Moluccas.	H. C. 1800.			
stricta, R.6 var. 1		Moluccas.	H. C. 1798.	♄ E.		
var. 2	Ch. Kang-mou-tang.	China, &c.	Before 1794.	♄ E.		
alba.	Ch. Pa-mou-tang.		1794.	♄ E.		
cuneifolia, R.		Dacca.	Col. P. Murray, 1801.	♄ E.	3---4.	6---7.
brachiata, R.		Bengal.	Accident, 1804.	♄ E.	3.	5---6.
parviflora.	B. *Gundhul-rungun.*7	India.	Before 1794.	S. T.	H. S.	R. S.
barbata, R.8			1794.	S. T.	3---5.	6---7.
undulata, R.	B. & H. *Paluk-jooi.*		1794.	S. T.	3---4.	6.
Pavetta, R.9	Tam. Pavatti.			L. ♄	5.	4.
tomentosa.	H. & B. *Jooi.*10			L. ♄	H. S.	R. S.
villosa, R.	H. *Choonaree.*	Silhet.	Mr. M. R. Smith, 1812.	♄ E.	5---6.	9---10.
acuminata, R.			1812.	♄ E.	4---5.	9.
CALLICARPA.						
incana, R.	B. Mattura.11	India.	Indigenous.	L. ♄ E.	H. & R. S.	C. S.
acuminata, R.		Silhet.	Mr. M. R. Smith, 1810.	♄ E.	4---6.	C. S.
Cana.		India.	Mr. W. R. 1803.	♄ E.	○	○
villosa.		Silhet.	Mr. M. R. Smith, 1810.	S. T.		
macrophylla.		Silhet, &c.	Mr. J. R. 1811.	♄ E.	R. S.	C. S.
arborea, R.		Chittagong.	Mr. W. R. 1801.	S. T.	○	○
lanceolaria, R.	B. Koamoor.	Silhet.	Mr. M. R. Smith, 1811.	♄	○	○
BUDDLEIA.						
Neemda.	B. *Nimda.*	Chittagong.	Dr. F. Buchanan, 1798.	S. T.	H. S.	R. S.
gujanense.		S. America.	Mr. W. Lockhead, 1808.			
EXACUM.						
tetragonum, R.	B. *Koochuri.*	Bengal.	Indigenous.	⊙ E.	R. S.	C. S.
carinatum, R.				⊙ E.		
sulcatum, R.				⊙ E.	C. S.	H. S.
hyssopifolium.12						
PLADERA.						
decussata, R.	B. *Dunkoni.*	India.	Indigenous.	S. ⊙ E.	R. & C. S.	C. S.
virgata, R.13				S. ⊙ E.	R. & C. S.	C. S.
pusilla, R.14		Coromandel.		S. ⊙ E.	R. S.	C. S.
PLANTAGO.						
major.	E. Great Plantain.	Britain.	Accident, 1795.	⊙	R. S.	C. S.
virginica.		Virginia.		⊙	R. S.	
lanceolata.		Europe.		⊙	R. S.	
Lagopus.		Spain, &c.	A. B. Lambert, Esq. 1802.		C. S.	2---3
Cynops.		Europe.	1802.	⊙	C. S.	1
littoralis.		Cape of G. H.	1799.			

1 H. M. 9. t. 76. 2 S. B. Munjistha. 3 H. M. 2. t. 12. 4 S. Bandhuka. H. Ranjun. 5 H. A. 4. t. 46. 6 Two varieties H. A. 4. t. 47. is the first, and B. M. 169. or Hang-mou-tong of the Chinese is the second. 7 T. Tadda-palla. 8 H. M. 2. t. 13. 9 H. M. 5. t. 10. T. Nuni-papata. 10 T. Naum-papata. 11 S. Mashandar. 12 See Gentiana verticillata. 13 Exacum diffusum Wild. 14 Hopea dichotoma Vahl. Exacum sessile Willd.

stricta.			A. B. Lambert, Esq. 1802.		C. S.	2—3.
Ispagula, R.	H. & B. Ispagool.1	India.	Dr. Fleming, 1808.	⊙ E.	C. S.	C. S.
CISSUS.						
sicyoides.		Chittagong.	Mr. J. R. 1810.	♄ Sc.	R. S.	C. S.
cordata, R.		Amboyna.	H. C. 1799.	♄ Sc.	R. S.	
vitiginea.	B. Gwali-lata.2	India.	Indigenous.	♄ Sc.	R. S.	
glauca, R.	B. Goor-gwaliya.			♄ Sc.	H. S.	R. S.
adnata, R.		Dacca.	Col. P. Murray, 1801.	♄ Sc.	H. S.	
quadrangularis.	H. & B. Har-jora.	India.	Indigenous.	♄ Sc.	H. S.	R S.
pentagona, R.		Chittagong.	Mr. J. R. 1810.	♄ Sc.		
angustifolia, R.		Chittagong.	1805.	♄ Sc.	2	R. S.
carnosa.3	T. Canapa-tiga.	India.	Indigenous.	♄ Sc.	5	
crenata.		Chittagong.	Mr. J. R. 1810.	♄ Sc.		
femina, R.		India.		♄ Sc.		
elongata, R.	B. Dhena.	Coromandel.	Before 1794.			
auriculata, R.		Mysore.	Mr. Heyne, 1802.	♄ Sc.	R. S.	C. S.
latifolia.		Chittagong.	Mr. J. R. 1810.	♄ Sc.	R. S.	C. S.
villosa, R.			1810.	♄ Sc.		
lanceolaria, R.		Dacca.	Col. P. Murray, 1801.	♄ Sc.	2.	
pedata.	B. Gwaliya.4	India.	Indigenous.	♄ Sc.	R. S.	R. S.
glandulosa, R.		Chittagong.	Mr. J. R. 1810.	♄ Sc.	R. S.	C. S.
serrulata, R.			1810.	♄ Sc.	4—5	
FAGARA.						
Rhetsa, R.	T. Rhetsa-maun.	Circars.	Palla-nundee, 1803.			
triphylla.5		Sumatra.	Dr. C. Campbell.	L. T.		
Ariscorina.		Bengal.	Dr. F. Buchanan, 1809.			
Budrunga.	B. Budrunga.	Silhet.	Mr. M. R. Smith, 1811.	L. T.	3—4	7—8
nitida.	Ch. Tcheun-tseu.	China.	Mr. W. Kerr, 1812.	♄ E.		
LUDWIGIA.						
parviflora, R.	H. & B. Bun lubunga.	India.	Indigenous.	⊙ D.	R. S.	R. S.
prostrata, R.		Pegu.	Mr. F. Carey, 1809.		R. S.	R. S.
OLDENLANDIA.						
alata.6	B. Gundha-bhadalee.	India.	Indigenous.	⊙ E.	○	○
paniculata.7		Moluccas.	Accidentally.	♂ D.	○	○
crystallina, R.8	B. Poonkka.	India.	Indigenous.	⊙ D.	○	○
diffusa, R.9				⊙ D.	R. S.	R. & C. S.
biflora.	B. Khet-papura.10			⊙ D.	R. S.	R. & C. S.
herbacea, R.11	T. Veri-nella-vemos.			⊙	R. S.	R. & C. S.
ramosa, R.		Pegu.	Mr. F. Carey, 1809.	⊙ D.	R. & C. S.	R. & C. S.
triflora, R.		Bengal.	Indigenous.	⊙ D.	R. & C. S.	C. S.
AMMANNIA.						
vesicatoria, R.	B. Bun-murich.12	India.	Indigenous.	⊙ E.	R. S.	R. S.
rotundifolia.		Chittagong.	Dr. F. Buchanan, 1796.	⊙ D.	R. S.	R. S.
octandra.		Coromandel.	1800.	⊙ E.	R. S.	R. S.
multiflora, R.		India.	Indigenous.	⊙ E.	R. S.	R. S.
nana, R.				⊙ Cr.	R. S.	R. S.
TRAPA.						
bispinosa, R.	H. Singara.13	India.	Indigenous.	⊙ A.	5—6	C. S.
tetraspinosa, R.		Silhet.	Mr. M. R. Smith, 1811.	⊙ A.	R. S	C. S.
POTHOS.						
cordata.		America.	H. C. 1805.	♃		
scandens.14		India.	Mr. J. R. 1810.	♄ R.	4—5	R. S.
officinalis, R.	B. Guj-pippul.	Bengal.	Kurrim khan, 1807.	♄ R.	4—5	R. S.
Peepla, R.	B. Pipul.	Silhet.	Mr. M. R. Smith, 1810.	♄ R.	4—5	R. S.
gigantea, R.		Moluccas.	Mr. W. R. 1803.	♄ R.	9—10	
decursiva, R.		Silhet.	Mr. M. R. Smith, 1810.	♄ R.	4—5	R. S.
heterophylla, R.	H. & B. Kanta-man.	Bengal.	Kurrim khan, 1801.	♄ Cr.	C. S.	
Læsia, R.	H. & B. Kanta-cuchoo.	India.	Indigenous.	♃	C. S.	H. S.
CHLORANTHUS.						
inconspicuus.	Ch. Tcheu-lan.	China.	Mr. W. Kerr, 1812.	♄ E.	R. S.	
ELÆAGNUS.						
dulcis, R.	B. Shooshuna.	Silhet.	Mr. M. R. Smith, 1811.	T.	C. S.	H. S.
triflora, R.		Sumatra.	Dr. C. Campbell, 1804.	L. ♄ Sc.	C. S.	H. S.
conferta, R.	B. Gwara.	India.	1795.	L. ♄ Sc.	C. S.	H. S.
SANTALUM.						
album.	S. B. H. Chunduna.15	India.	Dr. Berry, 1796.	T.	○	○
GYROCARPUS.						
Jacquini.	T. Tanacoo.	India.	Gen. Macdowall, 1802.	S. T.	H. S.	
RIVINA.						
lævis.		West Indies.	W. Hamilton, Esq. 1800.	♃ E.	○	○

TETRANDRIA DIGYNIA.

CUSCUTA.						
reflexa, R.	B. Huldi-algosa.	India.	Indigenous.	⊙ P.		

1 A. Burirki-toona. 2 G. Marata-tiga. 3 H. M. 7. t. 9. 4 S. Godha-pudee.
5 H. A. 2. t. 62. 6 Hedyotes racemosa, Willd. S. Gundha bhudra. H. Thel-koodi. 7 Burm. ind. 98. t. 15.
8 Hedyotes pumila, Willd. 9 Hedyotes diffusa, Willd. 10 S. Kshetra purputea. 11 Hedyotes herbacea, Willd.
12 H. Dad-mari. G. Aghundra-pacoo. 13 S. Sringata, or Sringataka. B. Panee-phul. 14 H. M. 7. t. 40.
15 A. Sandul-abiyar. P. Sundul-sufed.

aggregata, R.		Persia.	Accident, 1803.	⊙ P.	3.	
capitata, R.	H. & B. Algoa.	Bengal.	Indigenous.	⊙ P.	3.	
sulcata, R.				⊙ P.	3.	

TETRANDRIA TETRAGYNIA.

COLDENIA.						
procumbens.		India.	Indigenous.	⊙ D.	C. S.	H. S.
BRUCEA.	[rowang.					
sumatrana, R.1	Ma. Ampaaoo-kar-	Sumatra.	Mr. Ewer, 1802.	S. T.	○	C. S.
MYRIOPHYLLUM.						
verticillatum.	Tam. Meelapa.	India.	Indigenous.	⊙ A.	R. & C. S.	C. S.
tuberculatum.		Bengal.		⊙ A.	R. & C. S.	C. S.
POTAMOGETON.						
indicum, R.		Bengal.	Indigenous.	⊙ A. Cr.	R. & C. S.	C. S.
tuberosum, R.				⊙ A.	2—3.	4—5.

1 H. A. 7. t. 15.

CLASSIS V.

PENTANDRIA MONOGYNIA.

HELIOTROPIUM.						
peruvianum.		Peru.	Mr. J. R. 1801.			
indicum.	B. Hatee-shoora.1	India.	Indigenous.	☉ E.		
coromandelianum.				☉		
curassavicum.		America.	H. C. 1807.		R. S.	
paniculatum, R.		Chittagong.				
CYNOGLOSSUM.						
racemosum, R.		India.	Indigenous.	☉ E.	R. S.	C. S.
diffusum, R.		Bengal.		☉ D.	9---2	
marifolium, R.		Chittagong.	Mr. W. R. 1801.	☉ D.		
BORAGO.						
officinalis.		Europe.	Accident, 1799.	☉		
indica.	B. Chota-kulpa.	India.	Indigenous.	☉ D.	C. S.	
zeylanica.		Malabar.	Dr. Buchanan, 1801.	☉ E.	C. S.	
spinulosa, R.		Bengal.	Indigenous.			
LITHOSPERMUM.						
viridiflorum.		Chittagong.	Mr. J. R. 1810.	♄ E.	☉	
ANCHUSA.						
italica.		Europe.	Dr. Hornman, 1807.			
ECHIUM.						
plantagineum.		Italy.	A. B. Lambert, Esq. 1802.	☉		
vulgare.		Europe.	W. Hamilton, Esq. 1810.	☉ D.	C. S.	
NOLANA.						
prostrata.		Peru.	A. B. Lambert, Esq. 1802.	☉ D.	1---2	
MENYANTHES.						
indica.	H. & B. Bura-chooli.2	India.	Indigenous.	☉ A.	R. S.	
cristata, R.3	H. Chooli.			☉ A.	C. & R. S.	
HOTTONIA.						
serrata.4						
indica.5						
ANAGALLIS.						
arvensis.		Europe.	Accident.	☉ D.	C. S.	
PLUMBAGO.						
zeylanica.	B. Chitra.6	India.	Before 1794.	♃	R. & C. S.	
rosea.	H. B. Lall-chita.		Before 1794.	♃	R. & C. S.	
PORANA.						
volubilis.		Bengal.	Uncertain.	♄ Tw.	C. S.	H. S.
paniculata, R.		Bengal.	H. Creighton, Esq. 1799.	♄ Tw.	C. S.	H. S.
racemosa, R.		Napaul.	Dr. F. Buchanan, 1802.	☉ Tw.	C. S.	C. S.
LETTSOMIA, R.						
splendens, R.		Chittagong.	Mr. J. R. 1811.	♄ Tw.		
aggregata, R.	T. Yerra-katha.	Coromandel.	Dr. Rottler, 1803.	♄ Tw.	C. S.	3
nervosa, R.	H. & B. Bish-tarak.7	India.	Indigenous.	♄ Tw.	R. S.	C. S.
speciosa.8	H. & B. Chota-bish-		Hon. C. Greville, 1807.	♄ Tw.	R. S.	C. S.
argentea, R.	[taruk.9	Bengal.	Mr. W. R. 1801.	♄ Tw.	R. S.	C. S.
setosa, R.		Circars.	H. C. 1804.	♄ Tw.	C. S.	D. S.
strigosa, R.		Bengal.	Dr. F. Buchanan, 1796.	♄ Tw.	10	
cuneata, R.		Mysore.	General Martin, 1796.	♄ Tw.	C. S.	
cymosa, R.		Malabar.	Captain Dickenson, 1802.	♄ Tw.	C. S.	H. S.
pomacea, R.		Mysore.	Dr. F. Buchanan, 1800.	♄ Tw.	5---11	C. S.
bona nox, R.		India.	1798.	♄ Tw.	R. S.	C. S.
ornata, R.		Hindoostan.	Col. Hardwicke, 1802.	♄ Tw.	R. S.	C. S.
CONVOLVULUS.						
I. *Twining, leaves undivided.*						
gangeticus, R.		Bengal.	Indigenous.	♄ Tw.	C. S.	H. S.
semidigynus, R.		Sirinagur.	Col. Hardwicke, 1796.	♄ Tw.	C. S.	H. S.
fastigiatus, R.		Bengal.	Dr. W. Carey.	♃ Tw.	○	
malabaricus.		Mysore.	Dr. B. Heyne, 1805.	♄ Tw.	C. S.	H. S.
blandus.		Coromandel.	H. C. 1803.	♃ Tw.	D. S	H. S.
laurifolius, R.		Mysore.	Dr. B. Heyne, 1805.	♃ Tw.	C. S.	H. S.
parviflorus.		Coromandel.	Mr. P. Heshusius, 1802.	☉ Tw.	C. S.	H. S.
calycinus, R.		Bengal.	Col. Hardwicke, 1802.	♃ Tw.	R. S.	R. S.
obscurus.		India.	Dr. F. Buchanan, 1802.	☉ Tw.	C. S.	H. S.
purpureus.	Convolvulus major.	America.	General Martin, 1800.	☉ Tw.	C. S.	H. S.

1 S. Shree-hustinee. 2 S. Koomooda. 3 H. M. 11. t. 29. 4 See Serpicula.
5 See Columnea. 6 or Chita. S. Chitruka. T. Citra. 7 T. Koakaiti. 8 Does not differ specifically from C. nervosus. 9 or Sumoodra-shoka.

discolor.			Dr. Jacquin, 1806.	⊙ Tw.	C. S.	H. S.
umbellatus.			W. Hamilton, Esq. 1805.	♄ Tw.	C. S.	H. S.
sphœrocephalus, R.						
red flowered.1		Bengal.	Dr. W. Carey, 1801.	⊙ Tw.	C. S.	H. S.
white flowered.		Bengal.	Mr. W. R. 1801.	⊙ Tw.	C. S.	H. S.
pilosus, R		Mysore.	Dr. B. Heyne, 1802.	⊙ Tw.	C. S.	H. S.
arvensis.		Europe.	W. Hamilton, Esq. 1804.	H. Tw.	C. S.	
Malcolmii, R.		Persia.	Major J. Malcolm.	H. Tw.	D. S.	
medium.		India.	Accident.	♂ Tw.	R.&C.S.	
tridentatus.			Dr. F. Buchanan.	⊙ Tw.	R. S.	
bicolor.	H. Jurud-kulmee.		Indigenous.	♃ Tw.	R.&C.S.	H. S.
Turpethum.	H. B. Doodh-kulmee.2			♃ Tw.	R.&C.S.	H. S.
II. *Twining, leaves lobate, or compound.*						
hederaceus.		America.	Dr. Klien, 1804.	⊙ Tw.	C. S.	R. S.
carolinus.		Carolina.	W. Hamilton, Esq. 1805.	⊙ Tw.	D. S.	D. S.
Nil.		America.	1797.	⊙ Tw.	C. S.	H. S.
vitifolius.		Coromandel.	H. C. 1802.	♃ Tw.	R.&C.S.	H. S.
dentatus.		India.	Indigenous.	♂ Tw.	C. S.	H. S.
copticus.					C. S.	C. S.
paniculatus.	B. & H. Bhooi-koomra.3			H. Tw.	C. S.	H. S.
digitatus, R.		Coromandel.	H. C. 1804.	♂ Tw.	C. S.	H. S.
pedatus, R.			Accident, 1812.	Tw.	C. S.	H. S.
hirsutus, R.		Hindoostan.	Colonel Hardwicke, 1804.	⊙ Tw.	R. S.	C. S.
macrorhizos.		America.	W. Hamilton, Esq. 1801.			
viscidus, R.		West Indies.	Mr. W. Lockhead, 1807--8.	Tw.	C. S.	
pentaphyllus.			1807--8.	Tw.	C. S.	
heptaphyllus.		India.	Indigenous.	♃ Tw.	C. S.	
dissectus.		America.	W. Hamilton, Esq. 1801.	Tw.	C. S.	H. S.
III. *Prostrate or creeping (not twining).*						
siculus.		Europe.	H. C. 1807.	⊙ D.	D. S.	
reniforme, R.		Bengal.	Indigenous.	♃ Cr.	C. S.	C. S.
flagelliformis, R.		Mysore.	Dr. B. Heyne.	♃ Cr.	R.&C.S.	
reptans.	B. Kulmee-shak.4	India.	Indigenous.	♂ Cr.	C. S.	
Batatas.5	B. Sukur-kunda.					
red.6			Before 1793.	H. Cr. T.	D. S.	
white.7		Malayana.	Mr. W. R. 1802.	H. Cr.	D. S.	
cespitosus, R.		Bengal.	Dr. W. Carey, 1800.	⊙ Cr.	C. S.	H. S.
stipulaceus, R.		Chittagong.	Mr. W. R. 1802.	♃ Cr.	H. S.	
pentagonus, R.		Moluccas.	H. C. 1800.	♃ Cr.	D. S.	
bilobatus, R.			H. C. 1800.	♃ Cr.	R. S.	
pes-capræ.	B. Dobutee-luta.8	India.	Dr. Berry, 1800.	♃ Cr.	⊙	D. S.
brasiliensis.		Brazil, &c.	Hon. C. Greville, 1807.	♃ P.		
IPOMOEA.						
grandiflora, R.	H. Dhol-kulmee. 9	India.	1795.	♃ Tw.	☌	☌
salicifolia, R.		Bengal.	Dr. F. Buchanan, 1809.	⊙ Tw.	R. S.	C. S.
muricata, R.10		Malabar.	Mr. Robertson, 1794.	⊙ Tw.	R. S.	C. S.
sepiaria, K.	G. Metta-tooti.11	India.	Indigenous.	♄ ♃ Tw.	☋	☋
cœrulescens.			Dr. Rottler, 1806.	♃ Tw.	C. S.	H. S.
cœrulea, K.	H. Neel-kulmee.12		Indigenous.	⊙ Tw.	D. S.	D. S.
phœnicea, R.13		Dindigul.	Dr. A. Berry, 1799.	⊙ Tw.	C. S.	H. S.
coccinea.		West Indies.	Dr. W. Carey, 1796.	⊙ Tw.	C. S.	H. S.
lacunosa.		N. America.	W. Hamilton. Esq.		C. S.	
carnea.		America.	Hon. C. Greville, 1807.	♄ Tw.	C. S.	H. S.
ventricosa.		Furrukabad.	Mr. Robertson.	Tw.	10---12	H. S.
tuberosa.		West Indies.	1794.	L. ♄ Tw.	C. S.	H. S.
Pes-tigrides.	B. Langulee-luta.14	India.	Indigenous.	⊙ Tw.	R.&C.S.	
Quamoclit.	B. Turoo-luta.15			⊙ Tw.	R. S.	C. S.
red.	B. H. Lal-turoo-luta.			⊙ Tw.	R. S.	C. S.
white.	B. H. Suffet-turoo-luta.					
CAMPANULA.						
americana.		N. America.	W. Hamilton, Esq. 1802.	⊙		
SPHENOCLEA.						
zeylanica.	H. Jeel-murich.16	India.	Indigenous.	⊙ E.	R. & C.S.	C.S.
NAUCLEA.						
Cadamba, R.	H. & B. Kudum.17	India.	Indigenous.	L. T.	R. S.	10--12
macrophylla, R.		Amboyna.	H. C. 1797.	L. T.	6---7	
undulata, R.			H. C. 1798.	T.	R. S.	10--12
cordata, R.		Ceylon.	General Macdowall, 1802.	T.	5---6	C. S.
parvifolia, R.	T. Bota-kudumee.	Coromandel.	1800.	T.	H.&R.S.	C.S.
cordifolia.	B. Keli-kudum.18	India.	Mr. W. Gwilt, 1801.	L. T.	H. S.	H. S.
tetrandra, R.	B. Shwet kudum.	Silhet.	Mr. M. R. Smith, 1811.	S. T.	H. S.	C. S.
ovalifolia, R.	B. Shala.		Mr. M. R. Smith, 1811.	T.		
RONDELETIA.						
exerta, R.		Bengal.	H. Creighton, Esq. 1801.	S. T.	H. S.	

1 Compare with Ipomœa tamnifolia. 2 T. Tella-tagada. 3 S. Vidaree. T. Matta-pal-tiga. 4 S. Kulumbee. T. Tooti-koora. 5 S. Shurkura-kunda. 6 H. & B. Lal-shukurkunda-aloo. 7 H. & B. Suffet-shukurkunda-aloo. 8 also Chagul-khoori. 9 S. Munda-vulee. H. M. 11. t. 50. 10 Convol. muricatus Linn. 11 H. M. 11. t. 53. 12 The seeds miri, and Kala-dana. 13 Is nearly allied to sanguinea. Vahl. 14 H. M. 11. t. 59. 15 S. B. H. Kama-luta. 16 T. Neeroo-pippul. 17 S. Kudumba. 18 T. Daduga.

paniculata, R.		**Moluccas.**	**Captain Anderson, 1802.**	**L. T.**	**3,4,5,6.**	
tinctoria, R.		Midnapore.	S. Swinton, Esq. 1809.	S. T.	12.	5.
RANDIA.						
racemosa, R.	B. Peetunga.	Chittagong.	Dr. F. Buchanan, 1796.	L. ♄ E.	○	○
stricta, R.	H. Shwet-buruna.	Napaul, &c.	Before 1794.	L. ♄ E.	○	○
polysperma, R.		Chittagong.	Mr. J. R. 1811.	♄ E.	5.	8---9.
CINCHONA.						
thyrsiflora, R.		Bengal.	Dr. W. Carey, 1799.	T.	6---7.	10--12.
PSYCHOTRIA.						
undata.		Bahamas.	W. Hamilton, Esq. 1804.	♄ E.	4---5.	
herbacea.	B. Kudi-munkoon.	India.	Mr. M. R. Smith, 1812.	H. Cr.	5---6.	
COFFEA.						
arabica.	E. Coffee.1	Arabia.	Before 1794.	L. ♄ E.	2---5.	11--12--1.
bengalensis, R.	B. Bun Kuwa.	Bengal.	1794.	♄ E.	H. S.	C. S.
myrtifolia, R.		Mauritius.	Monsieur Jannet, 1811.	♄ E.		
PÆDERIA.						
fœtida.	H. Ghundalee. 2	India.	1795.	♄ Tw.	C. S.	H. S.
recurva, R.		Chittagong.	Mr. J. R. 1810.	L. E. ♄		
erecta, R.		Silhet.	Mr. M. R. Smith, 1810.	♄ E.	R. S.	C. S.
CANTHIUM.						
parviflorum.	T. Ballusoo-kura.3	India.	1800.	♄ E.	H. S.	9.
parvifolium, R.		Chittagong.		♄ E.		
WEBERA.						
scandens.	B. Guja-kuta.	Silhet.	Mr. M. R. Smith, 1812.	♄ Sc.	4---5	
corymbosa.	H. & B. Kanura,4	India.	Mr. W. R. 1801.	♄ E.	H. S.	
odorata, R.	B. Patagruja.	Silhet.	Mr. M. R. Smith, 1812.	♄ E.	H. S.	
MUSENDA.						
corymbosa.		India.		♄ E.	8---11.	
frondosa.	H. Bebina.5	India.	H. Creighton, Esq. 1797.	♄ E.	H.&R.S.	C. S.
DENTELLA.						
repens.	B. Bhoomi-pat.	India.	Indigenous.	⊙ D.	D. S.	D. S.
HAMILTONIA.						
suaveolens, R.		Rajemahl.	Mr. W. R. 1801.	L. ♄	C. S.	H. S.
CHIOCOCCA.						
racemosa.		Jamaica.	W. Hamilton, Esq.	♄ D.	8---9	
VANGUERIA.						
spinosa.	H. & B. Muyuna.6	Bengal, China.	Indigenous.	L. ♄ E.	H. S.	R. S.
GARDENIA.						
I. *Unarmed.*						
florida, single.	H. & B. Gundha-raj.7	China.	Before 1794.	♄ E.	H. S.	C. S.
double.8			1794.	♄ E.	H. S.	
costata, R.		India.	1797.	S. T.	H. S.	C. S.
latifolia, R.	H. Papura.9		1800.	S. T.	H. S.	C. S.
arborea, R.	T. Garga.10	Coromandel.	1801.	S. T.	H. S.	C. S.
lucida, R.		Chittagong.	Dr. F. Buchanan, 1798.	S. T.	H. S.	C. S.
Thunbergia.11		Cape of G. H.	1799.	S. T.	6.	
Rothmanica.		Cape of G. H.	1799.	♄ E.		
obtusifolia, R.		Pegue.	Mr. F. Carey, 1809.	♄ E.		
II. *Armed.*		[nore.				
turgida, R.		Boutan, Bed-	Dr. W. Carey, 1799.	S. T.	H. S.	
uliginosa.	B. Piraloo.12	India.	Dr. W. Carey, 1799.	S. T.	H. S.	C. S.
dumetorum.	T. Manga.	Coromandel.	Mr. B. Heyne, 1796.	L. ♄ E.	H. S.	C. S.
nutans, R.			1802.	♄ E.	H. S.	C. S.
longispina, R.		India.	1799.	S. T.	H. S.	C. S.
spinosa.			1803.	♄ E.	H. S.	C. S.
campanulata, R.		Chittagong.	Mr. Maddern, 1802.	♄ E.	H. S.	C. S.
Randia.		West Indies.	Sir G. Younge, 1800.	♄ E.	H. S.	C. S.
floribunda.		Coromandel.	1800.	♄ E.	H. S.	C. S.
fasciculata, R.		Silhet.	Mr. M. R. Smith, 1812.	♄ E.	H. S.	
MORINDA.						
tinctoria, R.	H. & B. Ach.13	India.	Before 1794.	S. T.	○	○
bracteata, R.14		Ganjam.	Mr. Snodgrass, 1797.	S. T.	R.&C.S.	C. S.
exerta, R.	B. Bun-ach.15	Bengal.	Indigenous.	S. T.	H.&R.S.	C. S.
multiflora, R.	H. Aal.	Nagpore.	H. Colebrooke, Esq. 1701.	L. ♄ E.	R. S.	C. S.
angustifolia, R.		Chittagong.	D. F. Buchanan, 1797.	L. ♄ E.	3	5
citrifolia.16		India.	Before 1794.	S. T.	H.&R.S.	R. S.
LONICERA.						
Periclymenum.17		England.	H. C. 1806.	♄ Tw.	H. & R. S.	
sempervirens.18		N. America.	Mr. Aiton, 1807.	♄ Tw.	3--.4.	
TRIOSTEUM.						
perfoliatum.		N. America.	W. Hamilton, Esq.			
angustifolia.		Chittagong.	Dr. F. Buchanan.			
CŒVOLA.						
Taccada, R.19		India.	Mr. W. R. 1803.	L. ♄.	○	○

1 A. Bun. B. Kuwa. B. Bila murich. 2 Gundha-bhudra. 3 H. M. 5. t. 36. & 37. 4 or Kankra. S. Kurnikara. T. Boma-papata. 5 H. M. 2. t. 18. 6 The fruit Muyun-phul. S. Muduna.
7 S. Gundha-raja. 8 The same synonyma. 9 T. Karinga. 10 Tam. Karinga.
11 When thirteen years old 7---8 feet high, and flowered for the first time.marum. 12 T. Nella-kakisha.
Can. Bou-gu-næ. Tam. Wagattæ. 13 or Aal. S. Uchyoota. T. Tagaree. 14 H. A. 3. t. 98.
15 or Aal. 16 H. M. 1. t. 52. 17 Woodbine, or Honey-suckle. 18 Trumpet Honey-suckle.
19 H. M. 4. t. 59.

LOBELIA.						
radicans.		China.	Mr. W. Kerr, 1812.	⊙ Cr.	R.&C.S.	C. S.
triangulata, R.		India.	Indigenous.	⊙	R.&C.S.	C. S.
BÆOBOTRYS.						
indica, R.		Chittagong.	Dr. F. Buchanan, 1797.	L. ♄	C. S.	H. S.
ramentacea, R.	B. *Mulmooriya.*	Silhet.	Mr. M. R. Smith, 1812.	♄ E.	3---4	10---11
MIRABILIS.						
Jalapa, white.						
red.						
purple.						
yellow with varieties of all these mixed.	B. *Krishna-keli.*1	Indies.	Before 1794.	H. D.	R. S.	C. S.
ongiflora.		Mexico.	1807.			
ARDISIA.						
glandulosa, R.	H. *Bun Nurukalee.*	Silhet.	Mr. M. R. Smith, 1811.	♄.	R. S.	
paniculata, R.		Chittagong.	Mr. J. R. 1810.	♄ E.		
umbellata, R.		Sumatra.	Dr. C. Campbell, 1800.	♁ ♄ E.	4	9---10.
solanacea.2		India.	1799.	⊙ ♄ E.		
colorata, R.	B. *Umur-kalee.*	Silhet.	Mr. M. R. Smith, 1811.	♄ E.	♁	♁
ERYCIBE.						
paniculata, R.		Coromandel, &c.	Mr. J. R. 1810.	♄ Sc.	9---10	
EMBELIA.						
robusta, R.		Bengal.	Mr. W. R. 1801.	S. T.	R. S.	C. S.
ribes.	B. *Baberung.*	Silhet. &c.	Mr. M. R. Smith, 1811.	L. Sc. ♄	2---3	8---9
CHIRONIA.						
centauroides.	B. *Geema.*	Bengal.	Indigenous.	⊙ E.	D. S.	D. S.
VERBASCUM.						
Thapsus.		Asia & Europe.		⊙ E.	C. S.	H. S.
phlomoides.		Europe.	W. Hamilton, Esq. 1802.			
SOLANDRA.						
grandiflora.		Jamaica.	W. Hamilton, Esq. 1800.	♄ D.	12---1	4---5
DATURA.						
Stramonium.		America.	W. Hamilton, Esq.	⊙ E.		
Tatula.		Cape of G. H.	1799.	⊙ E.		
fastuosa double.	B. & H. Kala dhoo-	India.	Indigenous.	⊙ E.	♁	♁
single.	[*tura.*3					
Metel.	B. *Dhootura.*4	India, &c.		⊙ E.	♁	♁
lævis.		Abyssinia.	A. B. Lambert, Esq.	⊙ E.		
NICOTIANA.						
Tabacum.	H. & B. *Tumak.*5	America.	Before 1794.	⊙ E.	C. S.	H. S.
fruticosa.6			W. Hamilton, Esq. 1800.	⊙ E.	C. S.	H. S.
rustica.			1799.	⊙ E.	C. S.	H. S.
paniculata.		Peru.	Sir A. Hume, 1812.	♄		
glutinosa.		Peru.	1804.	⊙ E.	C. S.	H. S.
quadrivalvis.		America.	Mr. Mc Mahon, 1810.	⊙ E.	C. S.	H. S.
havanniensis.		Cuba.	Alex. Gordon, Esq. 1812.			
ATROPA.						
physaloides.		Peru.	F. Horsely, Esq. 1796.	⊙ E.	D. S.	H. S.
PHYSALIS.						
flexuosa.	H. Usgund.7	Coromandel.		S. ♄ E.	R. & C. S.	H. S.
alkekengi.		Persia.				
peruviana.	B. *Tapureea* 8	Peru.	Before 1794.	♃ D.	C. S.	H. S.
minima.	B. *Bun-tapureea.*9	India.	Indigenous.	☉ D.	R. & C. S.	H. S.
angulata.		Malacca.	Accident, 1812.	⊙	R. & C. S.	C. S.
SOLANUM.						
I. *Unarmed.*						
pubescens.	B. *Urush.*10	India.	Indigenous.	L. ♄ E.	♁	♁
auriculatum.		Mauritius.	Capt. Tennent, 1802.	L. ♄ E.	12---7	D. S.
spirale, R.	B. *Bagooa.*	Silhet.	Mr. M. R. Smith, 1810.	♄ E.	♁	♁
tuberosum.	H. *Bilati-aloo.*11	Peru.		H. D.	C. S.	
lycopersicum.		S. America.		⊙ D.	C. S.	
nodiflorum.		Mauritius.	Mons. Sere, 1810.	⊙ E.	♁	♁
rubrum.	B. & H. *Goorkhi.*	India.	Indigenous.	☉ E.	♁	♁
nigrum.		America.	W. Hamilton, Esq. 1809.	⊙ E.	♁	♁
decemdentatum, R.		China.	Mr. Kerr, 1803.	⊙ E.	♁	♁
Pseudocapsicum.		Madeira.	Col. Hardwicke, 1811.	♃ E.	♁	♁
macrocarpon.		Peru.	W. Hamilton, Esq. 1796.	♃ E.	♁	♁
æthiopicum.		Mauritius.	Col. Hardwicke, 1811.	☉	♁	♁
II. *Armed.*						
Melongena.12	H. & B. *Bagoon.*	India.	Before 1794.	♃ D.	C. S.	H. S.
longum, R.13	B. *Gooli Bagoon.*			♃ D.	C. S.	H. S.

1 H. *Goolabas* also *Sanji.* 2 B. *Kaka-thontee* and *Bhooi-jam.* T. Moryoora. S. *Dhoostoora.* 4 H. & B. *Sada-dhootura.* 5 and *Tambaca.* S. *Tamrakoota.* Broad leaved Virginian or sweet-scented Tobacco. 6 Narrow leaved tobacco. 7 T. Penneru. S. *Ushwagundha.* 8 or *Tophlu.* E. Brazil-cherry. 9 H. Toolati-pari. 10 S. *Gandheera.* T. *Russa-ghedi.* 11 Common Potatoe. 12 S. *Vartaki* or *Hingoolee.* T. Wang-kari. Of this very useful plant there are several varieties in constant cultivation by the natives, such as green, white, purple, yellow, &c. 13 Two or three varieties of this species are also cultivated, viz. yellow, purple, &c.

insanum.1		Moluccas.	Mr. C. Smith, 1798.	♂	C. S.	H. S.
diffusum, R.	T. Nalla mollunga.	India.	Indigenous.	♃ D.	R.&C.S.	C. S.
Jacquini.	B. H. Kanthakaree.2	Bengal.		♂ D.	○	○
marginatum.		Africa.	W. Hamilton, Esq.	♄ E.		
hirsutum, R.	B. Ram bagoon.	India.	Indigenous.	♄ E.	R. S.	C. S.
indicum.	B. H. Byakool.3			♄ D.	○	○
stramonifolium.		America.	W. Hamilton, Esq. 1800.	♄ E.	○	○
sodomeum.		Cape of G. H.	1798.	♄		
pentapetaloides, R.		Brazil.	Dr. Wallich, 1808.	♂ D.	○	○
trilobatum.	T. Oochinta kurra.	Coromandel.	Nalla-mandu.	♄ Sc.	○	○
æthiopicum.		Mauritius, &c.	Col. Hardwicke, 1811.	♃ D.	○	○
CAPSICUM.						
annuum.		America.	Before 1794.	♃ E.	○	○
grossum.	H. Kaffri-murich.			♃ E.	○	○
frutescens red.	B. Lall lunka murich.4	India.	Before 1794.	♃ E.	○	○
yellow.	B. Huldi-lunka murich.5		1794.	♃ E.	○	○
baccatum.		West Indies.	W. Hamilton, Esq.	♃ E.	○	○
purpureum, R.		Moluccas.	Accident, 1798.	♃ E.	○	○
minimum, R.	H. Dhan-murich.6	India.		♃ E.	○	○
SERISSA.						
fœtida.		China.	Before 1794.	♄ E.	○	
STRYCHNOS.						
Nux vomica.	B. H. Koochila.7	India.	1800.	T.	2	
Potatorum.	B. Nirmulee.8	India.	1800.	T. E.	H. S.	
colubrina.			Mr. M. R. Smith, 1811.	♄ Sc.	4	9---11
CORDIA.						
Myxa.	B. Buhooari.9	India.	Before 1794.	S. T.	4---5	
latifolia, R	B. Buhooari.10	Hindoostan.	Gen. Martin, 1801.	T.	5---6	
polygama, R.		Coromandel.	1799.	S. T.	5	
angustifolia, R.		Mysore.	Dr. F. Buchanan, 1801.	S. T.	3	5---6
monoica, R.	H. Pida.11	Coromandel.	H. C. 1799.	S. T.	R. S.	
Sebestena.		West Indies.	Sir G. Young, 1801.	S. T.	○	
campanulata.12		Moluccas, &c.	Mr. Burchell, 1809.	S. T.		
serrata, R.		Travancore.	Dr. A. Berry, 1808.	S. T.	8---9	
grandis, R.		Chittagong.	Mr. J. R. 1810.	T.	10---11	
EHRETIA.						
aspera.	T. Tella-juvi.	India.	H. C. 1804.	S. T,	R. S.	
lævis.	T. Serigada.		Dr. W. Carey, 1808.	S. T.	1---2	5---6
buxifolia.	T. Bapanaboori.]		H. C. 1802.	S. T.	5	10
internodis.		Mauritius.	Col. Hardwicke, 1811.	T.		
serrata, R.		Chittagong.	J. Glass, Esq.	T.	C. S.	6
CHRYSOPHYLLUM.						
cainito.	Star-apple.	West Indies.	H. C. 1795.	T.	H. S.	R. S.
acuminata, R.	H. Peetakara.	Silhet.	Mr. M. R. Smith, 1810.	T.	4---5	10
SIDEROXYLON.						
melanophleum.		Cape of G. H.	Dr. R. 1799.	S. T.		9---10
TECTONA.						
grandis.	H. B. Segoon.13	Pegu, Malabar.	Before 1794.	L. T.	R. S.	10---11
DALRYMPELIA.						
pomifera, R.	B. Junki-jam.	Silhet.	Mr. M. R. Smith, 1812.	T.	3---4	9---10
RHAMNUS.						
incanus, R.		Moluccas.	H. C. 1800.	T.	C. S.	H. S.
circumscissus.		India.	1801.	♄ D.	1,2,3	12---1
catharticus.		Europe.	H. C. 1807.	L. ♄		
virgatus, R.		Napaul.	Dr. F. Buchanan, 1802.	♄	H. S.	10--11
ZIZIPHUS.						
trinervéus, R.		Mysore.	Dr. F. Buchanan, 1801.	T.	H. S.	10--11
albens, R.		China.	Dr. W. Hunter, 1803.	S. T.	4	4
latifolius, R.		India.	Dr. W. Carey.	S. T.	3	R. S.
Jujuba.	B. Kool.14		Indigenous.	S. T.	R. S.	C. S.
vulgaris.		Persia.	Before 1794.	S. T.	R. S.	C. S.
nitidus, R.		China.	1794.	S. T.	H. S.	C. S.
elliptica, R.		Travancore.	Dr. A. Berry, 1808.	T.	5---6	
glaber, R.		Chittagong.	Mr. J. R. 1810.	♄ Sc.		
Lotus.		Persia.	Hon. C. A. Bruce, 1801.	S. T.	H. S.	
Xylopyrus.	T. Gotti.	India.	Col. A. Kyd, 1802.	S. T.	R. S.	C. S.
Caracatta.	Can. karakatta.	Mysore.	Dr. F. Buchanan, 1801.	S. T.	R. S.	C. S.
scandens.	B. Shya-kool.15	India.	Indigenous.	♄ Sc.	H. S.	R. S.
rotundifolius.		Mysore.	Dr. F. Buchanan, 1801.	L. ♄		
microphyllus, R.		Coromandel.	Dr. B. Heyne, 1807.	♄ D.		
tomentosus, R.		Chittagong.	Mr. J. R. 1810.	♄		
incurva, R.		Napaul.	Dr. F. Buchanan, 1801.	S. T.	R. S.	

1 H. A. 5. t. 86. f. 1. No variety of this, so far as I know, is ever cultivated on the continent of India.
2 S. Kanthakarika. 3 B. Goor-kamai. 4 S. Muricha. 5 H. Huldi-murich. 6 B. Dhan-lunka murich. 7 S. Kutaka. T. Musadi. 8 S. Kutaka. E. clearing nut. H. Nell-mall. T. Induga. 9 S. Buhoovaruka. H. Lesoora. A. Sepstan. T. Neckra. 10 H. Bura Lesoora. 11 T. Panagheri. 12 H, A. 2. t. 75. 13 T. Teek. Can. Jaadi. Mn. Jatus. 14 S. Kurkundha and Koli. H. Biur. T. Renga. 15 S. Vudura. T. Parami.

VENTILAGO.						
madraspatana.	B. *Rukta-ptita*.1	Coromandel.	H. C. 1803.	L. ♄ Sc.	C. S.	H. S.
GOUANIA.						
tiliæfolia.2		Coromandel.	Nalla-mandu, 1803.	L. Sc. ♄	9---1	H. S.
EUONYMUS.						
garcinifolius, R.	M. Moee.	Silhet.	Mr. M. R. Smith, 1812.	S. T.	○	○
atropurpurea.		Napaul.	Dr. F. Buchanan, 1802.	♄ E.	3---4.	
CELASTRUS.						
multiflorus, R.		Napaul.	Dr. F. Buchanun, 1802.	♄ Sc.	R. S.	1
montanus, R.	T. Pedda-chintu.	Coromandel.	Nalla-mandu, 1803.	♄ E.	C. S.	H. S.
emarginatus.	T. Chinny-chintu.		1804.	♄ E.	C. S.	H. S.
verticillatus, R.	N. Tibiliti.	Napaul.	Dr. F. Buchanun, 1802.	T.	1 & 6	4 & 9
nutans, R.		Mysore.	1800.	♄ Sc.	5	7---8
monospermus, R.	B. Tiara.	Silhet.	Mr. M. R. Smith, 1812.	♄ Sc.	4	10
robustus, R.	B. Sheelkoil.		1812.	L. T.	8	4
buxifolius, R.		Travancore.	Dr. A. Berry, 1809.	♄ D.		
PETTOSPERMUM.						
Tobira.3		China.	Mr. W. Kerr, 1812.	♄		
ELÆODENDRUM.						
glaucum,	Ceylon Tea.4	India.	Genl. Macdowall.	T. E.	H. S.	9---10
HOVENIA.						
dulcis.	N. Mun Kokoshi.	Napaul.	Dr. F. Buchanan, 1802.	T.	H. S.	R. S.
SCOPOLIA.						
aculeata.	T. Conda-cashina.5	India.	Mr. Chapman, 1797.	♄ Sc.	○	
CEANOTHUS.						
americanus.		America.	W. Hamilton, Esq. 1802.	T.		
BUTTNERIA.						
pilosa.		Chittagong.	1801.	♄ Sc.	R. S.	4
catalpæfolia.		America.	Mr. W. Lockhead, 1803.	♄ Sc.	9---10	
LEEA.						
sambucina.		Malay Islds.	Mr. C. Smith.	♄ E.	R. S.	C. S.
staphylea.	B. *Kookoor-jechwa*.6	India.	Indigenous.	♄ E.	R. S.	C. S.
hirta.	B. *Kaka-jungha*.7		Uncertain.	♄ E.	R. S.	C. S.
robusta, R.		Coromandel.	Nalla-mandu, 1802.	♄ E.	7---10	12--3
crispa.	B. & H. *Bun chalta*.	India.	Indigenous.	H. E.	R. S.	C. S.
macrophylla, R.	B. *Dholsumoodra*.8	Bengal.		H. E.	R. S.	C. S.
HEDERA.						
Helix.	Common Ivy.	England.	H. C. 1807.	♄ R.		
VITIS.						
indica.	H. *Amdhouka*.9	India.	Indigenous.	H. Sc.	3---4	6---8
lanata, R.			Mr. Abraham, 1800.	♄ Sc.	4---5	
latifolia, R.	B. *Goorila*.10		Indigenous.	L. ♄ Sc.	5---6	6---8
vinifera.	S. & B. *Draksha*.11		Before 1794.	♄ Sc.	2---4	6---8
parvifolia, R.		Bengal.	Col. P. Murray, 1801.	♄ Sc.	3---4	
capensis.		Cape of G. H.	1799.	L. ♄ Sc.		
CEDRELA.						
Toona, R.	B. H. *Toon*.12	India.	Before 1794.	L. T.	2---4	4---6
villosa, R.		Tipperah.	S. Harris, Esq. 1797.	L. T.		
CAMUNIUM.						
chinensis, R.	Ch. *Sam-yeip-lan*.13	China.	Mr. W. Kerr, 1812.	♄ E.	R. S.	
MILNEA.						
edulis, R.	B. *Gumee*.	Silhet.	Mr. M. R. Smith, 1811.	T.	6---7	9---10
MANGIFERA.						
indica.	B. & H. *Am*.14	India.	Indigenous.	L. T.	2---3	5---7
oppositifolia, R.	Burm. *Meriam*.15	Pegu.	Mr. F. Carey, 1809.	S. T.		
Gandaria.	Mn. Gandaria.	Moluccas.	Dr. C. Campbell, 1803.			
silvatica.16		Silhet.	Mr. M. R. Smith, 1811.	T.	9---10	2
VARECA.						
molucana, R.		Moluccas.	H. C. 1798.	S. T.	○	
heteroclita, R.		Amboyna.	H. C. 1798.	♄	○	
VIOLA.						
suffruticosa.		Coromandel.	Dr. Rottler, 1804.	S. ♃ D.	○	○
primulifolia.		Bengal, &c.	J. Addison, Esq. 1811.	⊙	○	○
apetala, R.		Mauritius.	1811.	⊙	○	○
IMPATIENS.						
balsamina.	B. *Doopatee*.17	Coromandel.	Captain Tennent.	⊙ E.	○	○
tripetala, R.		Silhet.	Mr. M. R. Smith, 1811.	⊙ E.	○	○
natans, R.		Coromandel.	Dr. B. Heyne, 1811.	⊙ A.	R. S.	C. S.
STRELITZIA.						
Reginæ.		Cape of G. H.	1799.	♃ E.		
MUSA.						
Sapientum.	H. & B. *Kula*.18	India.	Before 1794.	H.	○	○
the wild sort.	H. & B. *Ram-kula*.19	Chittagong.	Mr. Maddern.	H.	○	○

1 T. *Yerra-chirtli*. 2 Restinaria scandens. Gœrt. 3 Kœmpf. 796---7.
4 Celastrus. Vahl. 5 H. M. 5. t. 41. 6 T. Ancadoo. 7 S. Kakangee.
8 S. *Samoodruka*. 9 H. M. 7. t. 16. 10 H. M. 7. t. 7. 11 H. Angoor. 12 S. *Kooveruka*. *Toona*.
13 Two vart. 14 S. *Amra*. T. *Mamadi*. Of this, like the apple, there are numerous, but inconstant varieties. 15 Two varieties, sweet and sour. 16 and another undetermined species. 17 S. Dwiputree. H. Gool-mendi. 18 S. *Kudalee*. T. Aretti. Of this sort the cultivated varieties are infinite, some of them of exquisite taste and flavour, surpassing the finest pear. 19 Wully-kola, &c.

Paradisiaca.	H. & B. Katch-kula.1	India.	Before 1794.	H.	☋	☋
coccinea.	Ch. Cou-ang-chok-chee.	China.	1794.	H.	☋	☋
ornata, R.	B. Ramaneegee-kula.	Chittagong.	Mr. Maddern, 1801.	H.	☋	☋
superba, R.		Dindigul.	Dr. J. Anderson, 1800.	♃	R. S.	C. S.
textillis.	Manilla hemp.	Manilla.	Dr. J. Fleming, 1811.	H.		
glauca, R.		Pegu.	Mr. F. Carey, 1810.	♃	R. S.	C. S.
HELTCONIA.						
buccinata.3		Amboyna.	H. C. 1798.	♃ E.	H.S.	
ACHYRANTHES.						
aspera.3	B. *Upanga.*	India.	Indigenous.	♂ S. E.	☋	☋
Monsonia.4			Mr. H. Russell, 1803.	☉ C.	C. S.	H. S.
lappacea.5	B. cheecheereea.		Indigenous.	♃ D.	R.&C.S.	C. S.
alternifolia.	B. *Luta-muhooia.*6			☉ E.	R. S.	C. S.
prostrata.7		Moluccas.	Accident, 1798.	☉ D.		
ferruginea, R.	H. Lal-*Sirinchee.*	India.	Indigenous.	☉ D.	R. S.	
scandens, R.	H. *Nooriya.*	Bengal.		♃ Sc.	R.&C.S.	
lanata.8	H. Chaya.	India.		♂ E. ☉	C. S.	H. S.
incana, R.		Coromandel.	Mr. Julian, 1809.	☉ E.	R.&C.S.	C. S.
triandra, R.9	B. & S. Shalincha.		Indigenous.	☉ Cr.	R.&C.S.	C. S.
CELOSIA.						
argentea.	B. Shwet-*moorga.*10	India.		☉ E.	C. S.	C. S.
cristata, red.	B.& H. Lall *Moorga.*11			☉ E.	C. S.	C. S.
yellow.12	B. Huldi *Moorga.*			☉ E.	C. S.	C. S.
comosa.		Sumatra.	Dr. C. Campbell.	☉ E.	C. S.	C. S.
cernua, R.		Rajemahl.	Mr. W. R. 1801.	☉ E.	10--12	C. S.
nodiflora.		Coromandel.	Nalla-mandu, 1801.	☋	☋	☋
Monsoniæ.13						
DEERINGIA.						
celosioides.	B. Gola-mohuree.14	Bengal.	Indigenous.	♄ Sc.	C. S.	H. S.
tetragyna, R.		Moluccas.	Mr. C. Smith, 1800.		C. S.	H. S.
ILLECEBRUM.						
sessile.15						
lanatum.16						
CARISSA.						
Carandas, black.	B. *Kurumcha.*17	India.	Indigenous.	L. ♄ E.	2---5	6---9
red.	From China.	China.	Before 1794.	♄	☋	H. S.
villosa, R.		India.		♄	H. S.	R. S.
diffusa, R.	B. San *Kurumcha.*18		Mr. Chapman, 1797.	♄ D.	H.&R.S.	
OPHIOXYLON.						
serpentinum.	B. H. *Chundra.*19	India.	Indigenous.	S. ♄	☋	☋
GYNOPOGON.						
stillatum.20		Moluccas.	H. C. 1798.	♄ T.	C. S.	R. S.
Hunterii. R.		Pinang.	Dr. W. Hunter, 1802.	L. ♄ E.	☋	
CERBERA.						
odallam.21		India.	Before 1794.	T. E.	☋	☋
fruticosa, R.		Pegu.	Mr. F. Carey, 1810.	♄ E.	☋	☋
Thevetia.		West Indies.	W. Hamilton, Esq. 1801.	S. T.	☋	☋
quaternifolia, R.22		Sumatra.	W. Ewer, Esq. 1803.	T.		
ALLAMANDA.						
cathartica.		Guiana.	W. Hamilton, Esq. 1803.	L. ♄	☋	☋
VINCA.						
rosea red.	H. Gool-feringa.23	India.	Before 1794.	♃ E.	☋	☋
white.	H. Gool-feringa.24		1794.	♃ E.	☋	☋
NERIUM.						
odorum, single red.	H. & B. Lal *Kurubee.*25	Syria.	Before 1794.	♄	☋	
double red.	H.& B. *Padma Kurubee.*26		1794.	♄	☋	
single white.	H.&B. Shwet *Kurubee.*		1794.	♄	☋	
double white.				♄	☋	
single crimson.			1794.			
tinctorum, R.	T. Chit-ankaloo.	Coromandel.	1790.	T.	2---3	1---2
coccineum, R		Silhet, &c.	R. K. Dick, Esq. 1805.	T.	4---5	
grandiflorum, R.		India.	Dr. B. Heyne, 1804.	♄ T.		
reticulatum, R.	B. *Karunta.*27	Silhet.	Indigenous.	♄ T.	R. S.	
pisidium, R.			Mr. M. R. Smith, 1810.	♄ Sc.	5---6	1---2
antidysentericum.28				T.	H. S.	
coronarium.29						

1 T. Komaretti. Of the plantain, which I consider a variety of sapientum only, there are but few variations known to me, viz. three or four. 2 H. A. 5. t. 62. f. 2. 3 S. *Upamarga.* B. H. *Chichiria.* 4 Pluk. alm. t. 334. f. 4. & 357. f. 4. 5 H. M. 10. t. 59. 6 H. Gangatiah. T. Chanchelli-kura. Compare with A. polygonoides, Willd. ed. sp. pl. 1. p. 1195, and with his description of A. muricatum, p. 1193, which agrees well with this plant, particularly the corpus viride truncatum, &c. 7 H. M. 10. t. 79. 8 H. M. 10. t. 29. 9 H. M. 10. t. 11. 10 or *Chooa.* H. M. 10. t. 38 & 39. C. albida and margaritacea of Willd. I consider only varieties of this. 11 S. *Muyoora* and *Khurashwa.* 12 Willdenows castrensis, and coccinea, I can only consider varieties of the last, cristata and castrensis, I suspect, is only another deviation, by soil and culture. 13 See Achyranthes Monsoniæ. 14 H. *Lutmun.* 15 See Achyranthes triandra. 16 See Achyranthes lanata. 17 S. *Krishnapakphula.* *Avigna.* H. *Kurunda.* T. Waka. 18 T. Wakloo. 19 and Chotachand. S. *Chundruka.* 20 H. A. 5. t. 20. 21 H. M. 1. t. 39. 22 H. A. 2. t. 363. 23 E. red Periwinkle. 24 White Periwinkle. 25 S. *Kuruveera*, or *Pruteehasa.* 26 H. M. 9. t. 1. 27 T. Adivi-palla-tiga. 28 See Echites antidysenterica. 29 See Tabernæmontana coronaria.

ECHITES.						
acuminata, R.	H. *Pinguree.*	Silhet.	Mr. M. R. Smith, 1812.	♄ Sc.	4---5	1---2
antidysenterica, R.	H. *Koorchi.*1	India.	1795.	S. T.	2---5	10---11
scholaris.	B. *Chhatin.*2		Indigenous.	T.		
venenata, R.	T. Ganda-ganeroo.		Dr. C. Campbell, 1805.	♄ E.	○	○
frutescens, R.	B. & H. *Shyama-luta.*3		Indigenous.	♄ Tw.	R. S.	C. S.
dichotoma, R.	H. Hapur-malee.4	Bengal.		♄ Tw.	C.&H.S.	C. S.
coryophyllata, R.	S. & B. Malutee.5	India.	1802.	♄ Tw.	R. S.	
paniculata, R.		Silhet.	Mr. M. R. Smith, 1811.	♄ Sc.	H. S.	1---2
grandiflora, R.		Chittagong.	Mr. J. R. 1811.	L.Sc.♄	D. S.	C. S.
macrophylla, R.	B. Harkee.	Silhet.	Mr. M. R. Smith, 1812.	♄ Sc.	6	8---4
PLUMERIA.						
acuminata.	B. *Gobur-champa.*6	India, &c.	Before 1794.	T.	H. & R. S.	
alba.		West Indies.	W. Hamilton, Esq. 1801.	T.	R. S.	
TABERNÆMONTANA.						
coronaria, single.	S. B. H. *Tugura.*7	India.	Before 1794.	♄E.○	H.&R.S.	C. S.
double.8		India.	Before 1794.	♄E.○	H.&R.S.	C. S.
recurva, R.		Chittagong.	Dr. Buchanan, 1797.	♄ E.		
crispa, R.9		China.	Before 1794.	L.♄E.		
parviflora, R.		Sumatra.	Dr. C. Campbell, 1805.	S.♄E.	R. S.	
dichotoma, R.		Ceylon.	Rev. Dr. John, 1796.	S.T.E.	R. S.	
WILLUGHBEIA.						
edulis, R.	B. *Luti-am.*	Chittagong.	Mr. F. Carey, 1809.	L.Sc.♄ ○		H. S.
MELODINUS.						
monogynus, R.	H. *Sadul kuwa.*	Silhet.	Mr. M. R. Smith, 1811.	L.Sc.♄	3---4	10---11

PENTANDRIA DIGYNIA.

PERGULARIA.						
odoratissima.10						
glabra.11						
PERIPLOCA.						
esculenta.12						
emetica.13						
indica.14						
tunicata.15						
CYNANCHIUM.						
extensum.16						
viminale.17						
APOCYNUM.						
frutescens.18						
ASCLEPIAS.						
I. *Corol rotate.*						
gigantea, rosy.	B. *Akund.*19	India.	Indigenous.	L. ♄ E.	○	○
white.	B. *Shwet akund.*20			L. ♄ E.	○	○
acida.	S. B. *Soma-luta.*21		1800.	♄ Tw.	R. S.	
racemosa, R.				♄ Tw.	5---6	8--10
asthamatica.	H. *Unta-mool.*22			♄ Tw.	C. S.	
tunicata, R.	B. *Chagul-patee.*23			♄ Tw.	R. S.	
volubilis.	T. Doodi-palla.24			♄ Tw.	R. S.	
annularia, R.	T. Palla-gurgi.25		Colonel Hardwicke.	♃ Tw.	R. S.	
pseudosarsa, R.						
narrow leaved.	H. Sada-*booree.*26		Indigenous.			
broad leaved.	B. H. *Ununta-mool.*27					
rosea, R.	B. *Doodh-luta.*28			♃ Tw.	R. S.	C. S.
tenuissima, R.		Bengal.	Dr. F. Buchanan, 1796.	♃ Tw.	R. S.	C. S.
parasitica, R.		Delta of the Ganges.		P. ♃ Cr.		
curassavica.		S. America.	Mr. White.	♂ & ♃ E.	○	○
fruticosa.		Cape of G. H.	Dr. W. Carey, 1808.	♃ E.	R. S.	R.&C.S.
II. *Corol salver shaped.*						
tinctoria, R.	M. Taroom-Akkar.	Sumatra, &c.	Before 1794.	♄ Tw.	R.&C.S.	R. S.
echinata, R.	H. Sagwanee.29	India.	Indigenous.	♃ Tw.	R.&C.S.	H. S.
geminata, R.	H. Chota dood-*luta.*			♄ Tw.	R.&C.S.	
odoratissima.	H. Kunja-*lata.*30		Hon. N. C. Bruce.	♄ Tw.	R. S.	
pallida, R.			Indigenous.	♄ Tw.	R. S.	
laurifolia, R.					R. S.	3

1 S. *Kootuja.* T. Palla-cadija. 2 S. *Supta-purna.* 3 S. Syama. T. Nalla-tiga. 4 S. *Vishulya-krit.* Pergularia glabra Willd. 5 H. M. 9. t. 14. 6 H. Goolachin. 7 H. Firki-tagar. H. M. 2. t. 54. 8 H. Bura-*tugur.* H. M. 2. t. 55. 9 H. M. 1. t. 46. 10 See Asclepias odoratissima. 11 See Echites dichotoma. 12 See Asclepias rosea. 13 See Asclepias pseudosarsa. 14 See Asclepias pseudosarsa. 15 See Asclepias tunicata. 16 See Asclepias echinata. 17 See Asclepias acida. 18 See Echites frutescens. 19 S. B. *Urka.* T. Nella-jeleeroo. 20 S. *Ulurca.* T. Nella-jeeleeroo. 21 T. Tiga-chemoodoo. 22 T. Kaka-palla. 23 H. Callia-lata. 24 S. Madhu-malati. H. Teet-conga. 25 H. M. 9. t. 7. 26 T. Palla Soucundi. H. M. 10. t.34. 27 S. *Ununta.* T. Ghœdi-soucundi. These cannot even be called varieties, as both are produced from the same root, and I believe it includes Ceropegia tenuifolia. Mant. 346. Periploca indica Willd. and Periploca emetica. Retz. Its root has been used as a substitute for Sarsaparilla. 28 S. *Doogdhika.* T. Doodi-palla. 29 T. Jutuga. 30 Mn. Tunki. H. M. 7. t. 26. f. 1.

herbacea, R.		Bengal.	Dr. W. Carey.			
tenacissima, R.	H. *Jitee.*	Rajemahl.	Mr. W. R. 1801.	♄ Tw.	H. S.	
tingens, R.	Burm. Mayzing-aklay.	Pegu.	Dr. F. Buchanan.	♄ Tw.	5. 6.	
acuminata, R.		Chittagong	Mr. J. R. 1810.	♄ Tw.	5—7	
micrantha, R.		Hindoostan.	Col. Hardwicke, 1802.	♄ Tw.	R. S.	
pulchella, R.		Silhet.	Mr. M. R. Smith, 1812.	♄ Tw.	R. S.	
CEROPEGIA.						
Candelabrum.		India.	Dr. Klien & Mr. Rottler.	♄ Tw.		
tuberosa, R.	T. Batsala-mandu.		Dr. Klien & Mr. Rottler, 1801.	H. Tw.		
juncea, R.	T. Bella-mandu.	Coromandel.	Dr. Klien & Mr. Rottler.	H. Tw.		
STAPELIA.						
variegata.		Cape of G. H.	Mr. C. Smith, 1795.	P.		
umbellata, R.		Mysore.	Lady Clive.	♃ E.	5—6	
erivacaria.			Dr. F. Buchanan.			
CHENOPODIUM.						
urbicum.		India, &c.	Indigenous.	⊙ E.	C. S.	D. S.
album.	B. Beto Sag.1	Europe.	Accident.	⊙ E.	D. S.	D. S.
murale.				⊙ E.	D. S.	D. S.
viride.		India, &c.	Indigenous.	⊙ E.	D. S.	D. S.
laciniatum.		Bengal.		⊙ E.	D. S.	D. S.
botrys.		Europe.	W. Hamilton, Esq.	⊙ D.		
ambrosioides.		S. America.	Before 1794.	⊙ D.	○	○
ATRIPLEX.						
hortensis.	Garden orache.	Tartary.		⊙ E.	C. S.	
BETA.						
Ecula, White garden beet.		Europe.	Before 1794.	♂	C. S.	H. S.
Mangel worzel.			Colonel Beatson, 1811.	♂	C. S.	
vulgaris, green beet.			Before 1794.	♂	C. S.	
red beet.			Before, 1794.	♂	C. S.	
bengalensis, R.	H. & B. Palung.2	Bengal.	Indigenous.	☉ E.	C. S.	C. S.
GOMPHRENA.						
globosa, red.	H. Gool-mukhmul.3	In Gardens over India.	Before 1794.	⊙ D.	R. S.	C. S.
white.4				⊙ D.	R. S.	C. S.
MOACURRA.						
gelonioides, R.	H. Moukoora.	Silhet.	Mr. M. R. Smith, 1811.	S. T.	4—5	12
ULMUS.						
integrifolia.	T. Nauli.5	Coromandel.	Dr. Berry, 1802.	L. T.	C. S.	
virgata, R.		China.	Sir John Royds, 1797.	S. T.	11	2
CELTIS.						
orientalis.	B. Jeebuna.6	India	Indigenous.	T.	○	○
occidentalis.		America.	W. Hamilton, Esq.	T.		
tetrandra, R.		Napaul.	Dr. F. Buchanan, 1802.		3	9
NAMA.						
zeylanica.	H. Kanchura.7	India.	Indigenous.	⊙ Cr.	C. S.	H. S.
HYDROLEA.						
zeylanica.8						
GENTIANA.						
heteroclita.9						
diffusa.10						
verticillata.	T. Nella-gulli.	Coromandel.	Nalla-mandu, 1804.	S. ♃ Cr.	○	○
saponaria.		N. America.	W. Hamilton, Esq. 1804.	♃		
HYDROCOTYLE.						
asiatica.	B. Thall-kuri.11	India.	Indigenous.	♂ or ♃ Cr.	R. & C. S.	C. S.
rotundifolia, R.				☉ or ♂ Cr.	R. S.	C. S.
PANAX.						
conchifolium, R.12		Moluccas.	Dr. A. Berry, 1807.	♄ E.		
palmatum, R.		Chittagong.	Mr. J. R. 1810.	♄ E.	8—10.	10—11
digitatum, R.	B. *Soona.*13	Silhet.	Mr. M. R. Smith, 1810.	T.	C. S.	H. S.
fruticosum.14		Moluccas.	H. C. 1798.	♄	5—6.	C. S.
fragrans, R.	B. *Gooti Soona.*	Silhet.	Mr. M. R. Smith, 1811.	T.	C. S.	H. S.
DAUCUS.						
Carota.	H. & B. Gajur.15	Persia.		⊙ E.	C. S.	H. S.
CONIUM.						
maculatum.16		Britain.	J. Bentley, Esq. 1812.			
LIGUSTICUM.	[Ajouan.17					
Ajouan, R.	H. & B. *Juvanee*, or	Bengal.		⊙ E.	C. S.	H. S.
diffusum, R.	H. *Bun-ajouan.*18		Indigenous.	☉ D.	C. S.	H. S.
CUMINUM.						
Cyminum.	H. & B. Jeera.19	Persia, &c.		○	C. S.	
PHELLANDRIUM.						
stoloniferum.	B. Pan-trasee.	Bengal.	Indigenous.	⊙ E.	C. & H. S.	H. S.
CORIANDRUM.						
sativum.	H. & B. Dhunya.20			⊙ E.	C. S.	H. S.

1 S. *Vastooka.* 2 *Palunka.* 3 S. *Umlana.* H. Lall Gool-mukhmul. 4 H. Suffet Gool-mukhmul. 5 Tam. *Tambachi.* 6 S. *Jeevunee.* 7 S. *Languli. Kunchura.* 8 See Nama zeylanica. 9 See Pladera sessiliflora. 10 See Pladera virgata. 11 S. *Bheka-purni.* 12 H. A. 4. t. 31. 13 or *Kata-soona.* 14 H. A. 4. t. 33. 15 S. *Grinjuna.* A. Gunir. P. Zerduch. 16 Common Hemlock. 17 S. *Bruhmadurbha. Juvanika*, &c. 18 *Moduyuntee.* 19 S. *Jeeruka.* A. *Kimoon.* 20 S. Dhunya. N. Danga.

SESSELI.						
bengalensis, R.		Bengal.	Indigenous.	⊙ E.	D. S.	H. S.
PASTINACA.						
sativa.	E. Parsnep.					
ANETHUM.						
Fœniculum.	E. Fennel.			♃ E.	D. S.	H. S.
graveolens.	E. Dill.	Europe.		☉ E.	D. S.	H. S.
Sowa, R.	H. Sowa.1	India.	Before 1794.	⊙ E.	D. S.	H. S.
Panmorium, R.	H. Panmuhooree.2		1794.	⊙ E.	D. S.	H. S.
PIMPINELLA.						
Anisum.	B. Muhooree.	Egypt.		⊙		
APIUM.						
petroselinum.	E. Parsley.		Before 1794.	♂ E.	C. S.	H. S.
involucratum, R.	B. Radhoonee.3	Bengal.	1794.	☉ E.	D. S.	H. S.
graveolens.	E. Celery.	England.	1794.	♂ E.	D. S.	H. S.
SEMECARPUS.						
Anacardium.	B. H. Bhela.4	India.	1797.	T.	R. S.	C. S.
Cassuvium, R.5		Moluccas.	H. C. 1798.	T.	H. S.	R. S.
cuneifolium, R.		Hindoostan.	J. Pierard, Esq. 1808.	T.		
HOLIGARNA.						
longifolia, R.		Chittagong.	Dr. F. Buchanan, 1797.	T.	C. S.	5–6
racemosa, R.		Silhet.	Mr. M. R. Smith, 1812.	T.	2–3	6–7
RHUS.						
glabrum.		N. America.	W. Hamilton, Esq.	♃ Cr.		
Toxicodendron.			W. Hamilton, Esq.			
vernix.			W. Hamilton, Esq. 1801.	T.		
succedaneum.		China.	Dr. A. Berry, 1800.	T.	12–3	3
copallinum.		N. America.	W. Hamilton, Esq.			
Bucki-amelam, R.	Nep. Bucki-amelo.	Napaul.	King of Napaul, 1798.	S. T.	8–9	C. S.
radicans.		Virginia.	Ld. Viscount Valentia, 1807.	♄		
lucidum.		Cape of G. H.	1807.	♄ E.		
parviflorum, R.		Napaul.	Dr. F. Buchanan, 1801.	T. D.	R. S.	
lævigatum.		Cape of G. H.	1799.	S. T.		
SAMBUCUS.						
nigra.		Europe.	1799.	S. T.		
Ebulus.		Bengal, &c.	Dr. F. Buchanan, 1809.	H. E.	R. S.	C. S.
TAMARIX.						
indica.	B. Jhawoo.6	India.	Indigenous.	T.	R. S.	C. S.
dioeca, R.	B. H. Lall Jhawoo.7	Ganges.	Capt. Frazer, 1800.	T.	R. S.	C. S.
TURNERA.						
angustifolia.		America.	W. Hamilton, Esq. 1805.	♂ E.	○	○
PHARNACEUM.						
mollugo.	H. & B. Gima.8	India.	Indigenous.	⊙ D.	R.&C.S.	C. S.
pentagynum, R.	B. Dhoosura.9			⊙ D.	R.&C.S.	C. S.
ALSINE.						
media.	Chick-weed.	Europe.	Accident, 1804.	⊙ D.	C. S.	C. S.
BASELLA.						
rubra.*	B. Pooi.10	India.	Indigenous.	♃ T.	D. S.	H. S.
alba.*	H. Suffet Pooi.11			♃ T.	D. S.	H. S.
lucida, *cordifolia Willd.*	H. & B. Pooi.12		Before, 1794.	♃ S.	D. S.	H. S.

TETRAGYNIA.

EVOLVULUS.						
alsinoides.		India.	Col. Hardwicke.	♃ D.	C. S.	
pilosus. R.			Sir A. Hesleridge.	♃ D.	○	
ARALIA.						
nudiflora.			W. Hamilton, Esq. 1809.	♄ D.		
racemosa.		N. America.	W. Hamilton, Esq.			
digitata, R.	H. Dain.13	India.	Dr. F. Buchanan.	T. D.	H. S.	R. S.
umbraculifera, R.		Moluccas.	H. C. 1797.	T.		
LINUM.						
usitatissimum.	B. Musina.14	India, &c.	Before 1794.	☉ E.	C. S.	H. S.
trigynum.	H. Gool-ashroopi.15	Hindoostan.	1794.	♄ D.	C. S.	H. S.
ALDROVANDA.						
verticillata, R.	D. Malacca-jhanji.	Bengal.	Indigenous.	⊙ A.		
DROSERA.						
Burmanni.		India.	Mr. M. R. Smith, 1810.	⊙	C. S.	C. S.
GISEKIA.						
pharnacioides.			Indigenous.	⊙ D.	C. S.	C. S.
COMMERSONIA.						
echinata.		Moluccas.	H. C.	T.	○	○

1 Shuta-pooshpa. S. Shaleya. B. Suloopha. A Shibit. 2 S. Mudhoorika. 3 H. Ujmood. 4 S. Urooskura. Bhullatakee. T. Nella jedi. 5 H. A. 1. t. 70. 6 S. Jhavooka. 7 S. Pichoola. 8 Greeshmuja. 9 Dhoosura. 10 S. Pootika. B. Racta Bun-pooi. T. Yerra-batsalla. 11 T. Alla-batsalla. * These can only be considered as varieties of one species, and I have doubts if even lucida and cordifolia be any thing more; alba being the original root, or stock from whence they have all sprung. 12 T. Pedda-batsalla. 13 Compare with Hedera teribinthinacea. 14 S. Ooma. Matusee. H. & B. Tisi. E. Flax. 15 and Gooldadee.

CLASSIS VI.

HEXANDRIA MONOGYNIA.

URANIA.						
speciosa.1		Madagascar.	Captain Tennant, 1802.	T. E.	9---10	5---7
BROMELIA.						
Ananas.	H. & B. Ananus.2	S. America.	Before 1794.	B. ♃	H. S.	R. S.
striped leaved.3			Phillips, Esq. 1800.	♃ E.		
Karattas.		America.	W. Hamilton, Esq.	♃		
PITCAIRNIA.						
bromelifolia.		Jamaica.	W. Hamilton, Esq. 1800.	♃	C. S.	
BURMANNIA.						
disticha.	H. Tala-kattee.	India.	Mr. M. R. Smith, 1812.	⊙ E.	R. S.	R. S.
TRADESCANTIA.						
virginiana.		America.	Kew Garden, 1806.	♃ E.	R. S.	C. S.
discolor.			W. Hamilton, Esq. 1800.	♃ E.	O	O
axillaris.	H. Bagha-nulla.4	India.	Indigenous.	⊙ Cr.	R. S.	C. S.
imbricata, R.		India.		⊙ Cr.	R. S.	C. S.
malabarica.5						
PONTEDERA.						
vaginalis.	B. Nouka.6	India.	Indigenous.	♃	R.&C.S.	C. S.
hastata.	T. Neeroo-tamana.			♃	R.&C.S.	C. S.
dilatata.				♃	R.&C.S.	C. S.
sagittata, R.		Chittagong.	1810.			
plantaginea, R.		Bengal.	Indigenous.	♂ or ♃	D. R. S.	C. S.
ovata.7						
LEUCOJUM.						
æsticum.		Europe.	Dr. W. Carey.	H.	H. S.	
NARCISSUS.						
Pœticus.		Europe.	Mr. M'Mahon, 1810.	H.		
incomparabilis.			Mr. M'Mahon, 1810.	H.		
pseudonarcissus.		Europe.	Mr. M'Mahon, 1810.	H.		
Tazetta, yellow.8		S. of Europe.	H. C. 1806.	H.	1---2	
white.	P. & A. Nurgis.9	Persia.	Gen. Martin, 1796.	H.	1---2	
Jonquilla.10		S. of Europe.	1799.	H.		
PANCRATIUM.						
zeylanicum.11		Ceylon, &c.	Genl. H. M'Dowall, 1802.	♃	R. S.	
longiflorum.		Amboyna.	Mr. C. Smith, 1798.	♃	R. S.	
biflorum, R.		Hindoostan.		♃	R. S.	
triflorum, R.	B. Sada-kanoor.	India.	Before 1794.	♃	R. S.	
fragrans.		West Indies.	Kew Garden, 1800.	♃	C. S.	
maritimum.		Europe.	W. Hamilton, Esq. 1801.	♃	R. S.	
littorale.		Tierra-Bambo.	H. C. 1810.	♃	R. S.	R. S.
amboinense.12						
CRINUM.						
amœnum, R.13		Silhet.	Mr. M. R. Smith, 1812.	♃	H.&R.S.	
asiaticum.	H. Sookh-durshun.14	India.	Indigenous.	♃	R. S.	R. S.
ensifolium, R.		Pegu.	Dr W. Carey, 1812.	♃	H. S.	
longifolium, R.		Bengal.	Dr W. Carey, 1803.	♃	R. S.	R. S.
brevifolium, R.15				♃	H.&R.S.	
lorifolium, R.		Pegu.	Mr. F. Carey, 1810.	♃	R. S.	C. S.
sumatranum, R.		Sumatra.	Dr. C. Campbell, 1801.	♃	R. S.	R. S.
canaliculatum, R.				♃	H.&R.S.	
superbum, R.		Sumatra.	Dr. C. Campbell, 1802.	♃	H.&R.S.	R. S.
Toxicarum, R.	H. & B. Bura-kanoor.16	India.	Col. Kyd.	♃	R. S.	R. S.
nervosum.17		Amboyna.	Mr. C. Smith, 1798.	♃	R. S.	R. S.
augustum, R.18				♃	H.&R.S.	R. S.
latifolium, R.19		Bengal.		♃	H.&R.S.	R. S.
zeylanicum.	H. & B. Kanoor.20	India.	Before 1794.	♃	R. S.	R. S.
moluccanum, R.		Moluccas.	H. C. 1798.	♃	R. S.	
erubescens.21		America.	W. Hamilton, Esq. 1801.	♃	R. S.	R. S.
americanum.			W. Hamilton, Esq. 1801.	♃	H.&R.S.	R. S.
	Swamp lily.	N. S. Wales.	Mrs. Blaxland, 1812.			

1 Ravenalia of Sonnerat. 2 B. Anarush. 3 From Malacca. 4 T. Gollagandi. H. M. 10. t. 13. 5 H. M. 9. t. 63. See Commelina nudiflora. 6 H. Louka. T. Neeroo-cancha. 7 See Phrynium capitatum. 8 Yellow polyanthus narcissus. 9 White polyanthus narcissus. 10 Common Jonquill. 11 H. A. 6. t. 70. f. 2. 12 See Crinum nervosum. 13 S. Soodurshuna. 14 H. M. 11. t. 38. 15 See asiaticum. Curt. mag. 1073. 16 C. Toldo. H. A. 6. t. 69. 17 H. A. 6. t. 70. f. 1. 18 From the Mauritius. 19 H. M. 11. t. 30. 20 H. A. 5. t. 105. 21 Curt. mag. 1034.

AMARYLLIS.						
atamasco.		America.	W. Hamilton, Esq. 1798.			
formosissima.		Cape of G. H.		♃	H. & R. S.	
reginæ.			W. Hamilton, Esq. 1798.	♃	H. S.	
equestris.		West Indies.	W. Hamilton, Esq. 1800.	♃	H. S.	R. S.
beladona.		Cape of G. H.		♃		
vitata.			Dr. W. Carey, 1812.			
aurea.		China.	H. C. 1810.	♃	8---9	
sarniensis.			Dr. W. Carey, 1812.			
radiata.	Ch. Yuck-lan.	China.	Before 1794.	♃	R. S.	
crispa.			Dr. W. Carey, 1812.			
zeylanica.1						
LEUCOMIS.						
punctata.		Cape of G. H.	Dr. W. Carey, 1812.			
LILIUM.						
candidum.		Europe.	1799.			
ALLIUM.						
porrum.	B. Puroo.2		Before 1794.	♂	H. S.	
sativum.	B. Loshoon.3		1794.	♂	H. S.	
scorodoprasum.	E. Rocambole.					
ascalonium.	B. Gundhuna.4	India, &c.		♂	C. S.	
cepa.	E. Onion.5	Asia.		♂		
proliferum, R.		China.	H. C. 1796.	♂	C. S.	
tuberosum, R.		In Gardens in Bengal.		♃	R. S.	
TULIPA.						
brujniana.		Cape of G. H.	1799.			
gesneriana.		Europe.	Mr. M'Mahon, 1810.			
ALBUCA.						
major.		Cape of G. H.	1799.	♃	C. S.	
minor.			1799.	♃		
ASPHODELUS.						
clavatus, R.		Bengal.	Indigenous.	⊙ E.	C. S.	H. S.
ANTHERICUM.						
tuberosum, R.	T. Cushelli.6	India.	1800.	H. E.	R. S.	C. S.
GLORIOSA.						
superba.	B. Eesha-nungula.7	India	Indigenous.	H. Sc.	R. S.	C. S.
CURCULIGO.						
orchioides.	B. Tal-moolee.8	India.	Indigenous.	♃	O	
recurvata, R.		Chittagong.	Dr. F. Buchanan, 1798.	♃	R. S.	C. S.
sumatrana, R.9		Sumatra.	Dr. C. Campbell, 1800.	♃	H. S.	
ORNITHOGALUM.						
umbellatum.		India.		H.		
capinsis.			Mr. J. R. 1802..	♃		
thyrsoides, white and yellow.		Cape of G. H.	1799.	♃		
candatum.		Cape of G. H.	Dr. W. Carey, 1812.			
SCILLA.						
peruviana.		Peru.	W. Hamilton, Esq. 1804.	♃		
indica, R.		India.	Mr. W. R. 1800.	H. E.	H. S.	H. S.
coromandeliana, R.		Coromandel.	Dr. A. Berry, 1802.	H. E.	C. S.	H. S.
ASPARAGUS.						
officinalis.	P. & H. Nakdoun.10	Europe.	Before 1794.	H. E.	H. S.	
acerosus, R.		Bengal.	Mr. Creighton.	H. E.	9---12	C. S.
adscendens, R.		Rohilcund.	Mr. Gott, 1804.	H. E.	16---11	2---3
racemosus.	B. Shuta-moolee.11	India.	Indigenous.	♄ Sc.	C. S.	2---3
curillus.	N. Kurilla	Napaul.	Dr. F. Buchanan, 1801.	H. Sc.	R. S.	
DRACÆNA.						
angustifolia, R.12		Moluccas.	1795.	♄ E.	H. S.	R. S.
ferrea.	B. Bijoolee chutuk.13	India.	Before 1794.	♄ E.	H. S.	
terminalis.14		Moluccas.	Mr. C. Smith, 1798.	♄ E.	H. S.	R. S.
spicata, R.		Chittagong.	Dr. F. Buchanan, 1796.	♄ E.	H. S.	R. S.
maculata, R.		Sumatra.	Dr. C. Campbell, 1805.	♄	H. S.	R. S.
terniflora, R. 2 var.	H. Bhoonamtol.	Silhet.	Mr. M. R. Smith, 1812.	♄ E.	1---2	12
atropurpurea, R.				♄ E.	4---5	1
SANSEVIERA.						
zeylanica.	B. Shoochi mookhee.15	India.	Indigenous.	♃ E.	R. S.	R. S.
DIANELLA.						
nemorosa.16		India.	Dr. C. Campbell, 1797.	♃ E.	O	O
TETA.						
viridiflora, R.	H. Teta.	Chittagong.	Dr. F. Buchanan, 1798.	♃ E.	3---4	R. S.
POLYANTHES.						
tuberosa, single.	B. Rujuni-gundha.17		Before 1794.	♃ E.	R. S.	
double.	B. Ruj uni-gundha.18				R. S.	
HYACINTHUS.						
orientalis.	E. Garden Hyacinth.	Levant.		♃	C. S.	

1 And latifolia. Willdenow, see Crinum Zeylanicum, and latifolium. 2 E. Common Leek.
P. Gundana. A. Koorrass. 3 Lushoona. S. Mohoushoodha. A. Soom. P. Seer. 4 S. Pulandoo.
5 A. Besel or Bussel. B. Peeaj. 6 S. Caitra. 7 and Kookoora-neja. H. Kariari. H. Ulut-chandal.
8 or Tal-noor. T. Nella-tada. H. M. 9. t. 59. 9 H. A. 6. t. 53. 10 A. & B. Hilyoon. 11 S.
Shuta-moolee. 12 H. A. 4. t. 35. 13 H. A. 4. t. 34. f. 2. 14 H. A. 4. t. 34. f. 1. 15 S.
Moorva. H. Murta, Moorga. T. Chamakada. 16 Dracæna ensifolia, Willd. H. A. 5. t. 73.
17 E. Tuberose. H. Gool Shaboo. 18 E. Tuberose. H. Bura Gool Shaboo.

YUCCA.						
superba, R.1		Amorica.	1799.	♃ E.	R. S.	
gloriosa.		S. America.	1799.	♄ E.	R. S.	
filamentosa.		Virginia.	W. Hamilton, Esq. 1799.	♄ E.		
ALOE.	[maree.2					
perfoliata.	H. & B. Ghrita-koo-	India, &c.	Before 1794.	♃ E.	R. S.	
verrucosa.		Africa.	Kew Garden, 1806.	♃		
variegata.		Africa.	Kew Garden, 1806.	♃		
three undetermined species.		Mauritius.				
AGAVE.						
Cantala, R.	S. Kuntala.	India.	Before 1794.	♄ E.	5---6	8---10
lurida.		America.		♄ E.	5---6	8---10
tuberosa.3						
HEMEROCALIS.						
cordata.		China.	Mr. W. Kerr, 1812.	♃	R. S.	
fulva.	H. Gool-nurgis.	North of Bengal.	Dr. W. Carey, 1797.	♃	R. S.	
cœrulea.		China.	Dr. W. Carey, 1811.	♃		
ACORUS.						
Calamus.	B. Shwet Buch.4	India, &c.	1796.	♃	R. S.	
CALAMUS.5						
TACCA.						
pinnatifida.6		Malayana.	H. C. 1798.		R. S.	C. S.
aspera, R.		Chittagong.	Mr. J. R. 1810.	H.	R. S.	C. S.
lævis, R.	B. Moti-munda.	Silhet.	Mr. M. R. Smith, 1811.	H.	H.&R.S.	
CORYPHA.						
umbraculifera.	C. Tala.7	Ceylon.	Genl. H.M'Dowall, 1798.	L. T.		
Utan.8		Moluccas.	H. C. 1798.	T.		
Taliera, R.	S. Talee.9	Bengal.	1800.	T.	3---4	11---12
elata, R.	H. & B. Bajra.10		Before 1794.	L. T.	4	10---11
LICUALA.						
spinosa.		Moluccas.	H. C. 1798.			
peltata, R.		Chittagong.	Mr. W. R. 1801.	S. T.	11---12	5---6
CHAMÆROPS.						
humilis.		Europe.	W. Hamilton, Esq. 1800.	♄ E.	C. S.	H.
mites.			Hon. C. Greville, 1801.			
Hystrix.			W. Hamilton, Esq. 1809.			
NANDINA.						
domestica.		Japan.	Mr. W. Kerr, 1812.	♄		
ACHRAS.						
mammosa.		China.	Mr. W. Kerr, 1807.	T.		
Sapota.		West Indies.		T.	H. S.	H. S.
MIMUSOPS.						
Elengi.	B. Bukool.11	India.	Indigenous.	T.	H. S.	R. S.
Kauki.	B. Ksheerni.12	Malayana.	Dr. A. Berry, 1801.			
hexandra.	T. Palta.13	Coromandel.	1800.	T.	H. S.	R. S.
BERBERIS.						
asiatica, R.		Hindoostan.	Dr. F. Buchanan, 1802.	♄	H. S.	R. S.
pinnata, R.	B. Huding nioong mari.	Munipore.	Mr. M. R. Smith, 1812.	♄ E.	C. S.	
LORANTHUS.						
bicolor, R.	H. & B. Bura munda.14	India.	Indigenous.	S. P.		
globosus, R.	B. Chota-munda.	Bengal.		S. P.	O	O
TRIPHASIA.						
aurantiola.	H. & B. China narangi.15	India.	Before 1794.	♄ E.	O	O
BAMBUSA.						
arundinacea.	B. Bansh.16	India.	Before 1794.	T.	H. S.	
stricta, R.	T. Sadanapa-vedroo.			S. T.	H. S.	
Tulda, R.17		Bengal.	1794.	T.	H. S.	
Balcua, R.18			1794.	T.	H. S.	
baccifera, R.	B. Bish-bansh.19	Chittagong.	F. Pierard, Esq. 1801.	S. T.	R. S.	C. S.
spinosa, R.	B. Behoor-bansh.20	Bengal.	Before 1794.	S. T.	H. S.	
nana, R.		China.	1794.	♄		

DIGYNIA.

ORIZA.						
sativa.21	S. Arunya, also Ashoo vrihi.			⊙ E.	R. S.	C. S.

1 Is Agave tuberosa. Hort. Kew. 2 S. Koomaree, &c. 3 Hort. Kew. is our Yucca superba. 4 S. Vucha. 5 See Dicœcia Hexandria. 6 H. A. 5. t. 112. 7 Tam. Conda pani. 8 H. A. 1. t. 11. 9 H. Taliera. B. Tariyat. 10 or Bajra-batool. 11 S. Kesura Vukoola. H. Mulsari. T. Pagada. 12 S. Ksheerika. H. Kheeri. M. Booaso. 13 Tam. Paltæ. 14 S. Vunda. T. Yellinga. Wadenika. 15 Limonia trifoliata Linn. 16 S. Vungsha. T. Vedroo. H. M. 1. t. 16. 17 Three varieties, viz. Tulda, Bashini, and Peea-bansh. 18 Two varieties, viz. Bhalkoo, or Puloo, and Dhooli-bansh. 19 H. Paya-tullu. H. M. 1. t. 60. 20 H. A. 4. t. 3. 21 B. Dhan the plant and the paddy, and chaul the rice. A. Aruz. T. Oori the plant, Oodloo paddy, or grain in the husk, and bium the rice. The varieties of this most useful grain cultivated over Asia are endless. On the Coast of Coromandel alone I detected and described above forty all well known to the native farmers. The original wild stock, from whence I am inclined to think all have sprung, is still common on the borders of lakes, pools, &c. during the rainy season in many parts of India. Its Sungskrit name is Neevara. The grain of this wild sort, is much esteemed by many of the natives of the northern Circars.

LEERSIA.						
aristata, R.	B. Kolshia.1	Bengal, &c.	Col. Hardwicke.	⊙ A.	R. S.	C. S.
ciliate.			Indigenous.	⊙ A.	10---11	C. S.
RUMEX.						
acutus.	B. Bun-palung.2	Bengal.	Indigenous,	☾ E.	1---3	4---5
vesicarius.	B. Chookа-palung.3	India.	Before 1794.	⊙ E.	R. S.	C. S.
acetosella.		Europe.	Captain Tennent, 1802.	♃	2	
acetosa.			Mr. M'Mahon, 1809.	♃		
FLAGELLARIA.						
indica.	H.&B.Bun-chandur.4	India.	Indigenous.	♄ Sc.	R. S.	C. S.
APONOGETON.						
monostachyon.	H. Ghetchoo.5	India.	Indigenous.	H. A.	R. S.	C. S.
undulatum, R.		Bengal.		H. A.	R. S.	C. S.
microphyllum, R.			Dr. W. Carey, 1799.	H. A.	R. S.	C. S.
HELONIAS.						
asphodeloides.		Virginia.	W. Hamilton, Esq.			

HEXAGYNIA.

DAMASONIUM.						
indicum.	B. Panee-kula.6	India.	Indigenous.	⊙ A.	R. S.	C. S.

1 H. M. 10. t. 12. 2 B. Jol-palung. 3 S. Chookrika. Shutuvedhee. A. Hooman.
4 T. Poindi-pooti. H. Harchara. 5 S. Kakanjee. T. Namma. 6 T. Neer-veneki.

CLASSIS VII.

HEPTANDRIA MONOGYNIA.

PISONIA.						
aculeata.1	B. Bag-achera.	India.		L. Sc. ♄		
mitis.	T. Konki-putri.	Madagascar. Mons. Jannet, 1811.		♄ E.		
JONESIA.						
Asoca.	S. B. Ushoka.2	Bengal.	Before 1794.	T.	3—4	8—9
scandens.		Sumatra.	Dr. C. Campbell, 1804.			
PETIVERIA.						
alliacea.		W. Indies.	W. Hamilton, Esq.	♄		

1 Tam Karu-Indu. Plants reared from seed from the West Indies, do not in any respect differ from those of India. Dr. Kœnig considered it an undescribed genus, when he first found it, and gave it the name Tragularia horrida. 2 H. *Ussu*. H. M. 5. t. 59.

CLASSIS VIII.

OCTANDRIA MONOGYNIA.

TROPÆOLUM.						
majus.	Great Indian Cress.	S. America.	1799.	☉ Sc.	C. S.	
XANTHOPHYLLUM.						
virens, R.	B. Gandhee.	Silhet.	Mr. M. R. Smith, 1812.	L. T.	2---4	6---7
ŒNOTHERA.						
sinuata.		America.	Mr. M'Mahon, 1809.		C.&H.S.	H. S.
purpurea.					C.&H.S.	R. S.
tetraptera.		S. America.	Captain Thomas, 1808.	♃ D.	☾	
longiflora.		America.	Dr. W. Carey, 1811.	♂ D.	3---5	
rosea.		N. America.	Dr. Hornman, 1807.	♃	☽	
grandiflora.		America.	Mr. M'Mahon, 1809.	♂ D.	D. S.	
GAURA.						
biennis.		N. America.	Dr. W. Carey, 1807.	♂		
COMBRETUM.						
purpureum.		Mauritius.	H. Hope, Esq. 1810.	L. Sc. ♄		
decandrum.	T. Arry-koot*a*.	India.	1803.	L. Sc. ♄	C. S.	R. S.
chinensis, R.		China.	Colonel Kyd.	L. Sc. ♄	C. S.	R. S.
extensum, R.		Moluccas.	H. C. 1798.		1---2	R. S.
ovalifolium, R.		Coromandel.	1804.	L. Sc. ♄	3---4	R. S.
pilosum, R.	B. *Jooni*-ugur.	Silhet.	Mr. M. R. Smith, 1811.	L. Sc. ♄	2---3	R. S.
acuminatum, R.	B. Pat-*jooniya*.		Mr. M. R. Smith, 1812.	L. Sc. ♄	3---4	7---8
costatum, R.	B. Tal*i*-*jooniya*.			L. Sc. ♄	3---4	
GRISLEA.						
tomentosa.	B. Dhub.1	India.	Mr. B. Heyne, 1795.	♄ E.	5	2---3
ROXBURGHIA.						
viridiflora.		Chittagong.	Dr. Buchanan, 1798.	♄ T.	H. S.	C. S.
MARLEA.						
begonifolia, R.	B. Marl*iya*.	Silhet.	Mr. M. R. Smith, 1812.	S. T.	3---4	7---8
XYLOCARPUS.						
granatum.	H. & B. *Pusoor*.2	India.	F. Goodlad, Esq. 1798.	T.	R. S.	C. S.
GUAREA.						
binectarifera, R.		Chittagong.	Dr. F. Buchanan, 1798.	T.	5	2---3
paniculata, R.	B. Kalee-koura.	Silhet.	Mr. M. R. Smith, 1812.	L. T.	5---6	4---5
AMYRIS.						
agallocha, R.	S. & B. *Googgool*a.	Garrow hills.	Before 1794.	S. T.	1---2	
gileadensis.		Arabia.	Dr. Berry, 1798.	♄		
acuminata, R.		Moluccas.	H. C. 1798.	T.	H. S.	
pentaphylla.		Hindoostan.	Colonel Hardwicke, 1802.	♄ E.	H. S.	R. S.
heptaphylla, R.	H. Karun-phul.	India.	1796.	♄ E.	H. S.	R. S.
nana, R.		Amboyna.	H. C. 1798.	♄	H. S.	R. S.
fruticosa, R.		Chittagong.	Dr. Buchanan, 1796.	♄ E.	H. S.	R. S.
punctata, R.			Before 1794.	S. T.	H. S.	R. S.
sumatrana, R.		Sumatra.	Dr. C. Campbell, 1807.	T.	5---9	C. S.
ACER.						
negundo.		America.	Monsieur Jannet, 1810.	T.		
XIMENIA.						
americana.	T. Oora-nekra.	India.	Mr. B. Heyne, 1803.	♄ E.		
ægyptiaca.	H. Hingun.3		Dr. F. Buchanan.	S. T.	5---6	
PIERARDIA.						
sapida, R.	H. Lutk*a*.	Chittagong.	Before 1794.	S. T.	2	5
DODONÆA.						
angustifolia.		India.	Mr. Heshusius, 1802.	♄ E.	R. S.	C. S.
dioeca, R.		Hindoostan.	1796.	♄ E.	R. S.	C. S.
LAWSONIA.						
inermis.	H. & B. Mindee.4	India.	Before 1794.	S. T.	☉	
MEMECYLON.						
capitellatum.		India.	Dr. A. Berry, 1810.	♄ E.	2---3	
SYMPHOREMA.						
involucrata, R.	T. Suroodoo.	Coromandel.	Mr. B. Heyne, 1808.	♄ Sc.	2---3	
ORNITROPHE.						
serrata.	B. Rakhal-phul.5	India.	Indigenous.	L. ♄	R. S.	C. S.
glabra, R.		Chittagong.	S. Harris, Esq.	L. ♄	R. S.	C. S.
Aporetica, R.	H. Ghee-kashee.	Silhet.	Mr. M. R. Smith, 1810.	L. ♄	6---7	C. S.
villosa, R.		Chittagong.	Mr. J. R. 1810.	L. ♄		
SCYTALIA.						
Litchi, R.6		China.	Before 1794.	T.	2	4---5

1 S. *Ugni*-jwala. Dhatr*i*-*pooshpika*. H. Dhuw*i*. T. Seringi. 2 S. Puroosh*a*. Tam. Kandalanga. T. Garu. 4 P. Hena. A. Alhinna. T. Gorinta. 5 T. Tantichi. 6 Ch. Leetchee, many varieties.

rimosa, R.	B. Tingurce.	Silhet, &c.	Mr. M. R. Smith, 1811.	L. T.	3---4	8
Longan, R.	B. Ash-phul.	China, Bengal.	Before 1794.	T.	2---3	6
Rambootan, R.	M. Rambootan.	Malayana.	H. C. 1798.	T.		
verticillata, R.		Moluccas.	H. C. 1798.	♄	R. S.	H. S.
Danura, R.	H. Danura.	Chittagong.	Dr. F. Buchanan, 1797.	T.	H. S.	
rubra, R.	H. Koi-poora.	Silhet.	Mr. M. R. Smith, 1812.	T.	2---3	4---5
SCHLEICHERA.						
trija.	C. Kunghas.1	Ipdia.	1797.	T.	H. S.	6---7
pentapetala, R.	B. Poora-koi.	Silhet.	Mr. M. R. Smith, 1811.	T.	H. S.	8---9
MOLINÆA.						
canescens.		Coromandel.	Dr. A. Berry, 1809.	T.	H. S.	
MELICOCCA.						
bijuga.2		Jamaica.	H. C. 1795.	T.	2---3	
SAPINDUS.						
emarginatus.	B. Bura-reetha.3	India.	Sir J. Royds, 1800.	T.	10---11	D. S.
laurifolius.4			Dr. Buchanan, 1800.	T.	12	4
detergens, R.	B. Reetha.5	Bengal,	Indigenous.	T.	H. S.	C. S.
Saponaria.		West Indies.	W. Hamilton, Esq. 1800.	T.	H. S.	C. S.
rubiginosus, R.		India.	Indigenous.	T.	H. S.	C. S.
fruticosus, R.		Moluccas.	H. C. 1797.	♄ E.	H. &R. S.	C. S.
polyphyllus, R.		Pegu.	Mr. F. Carey, 1810.	T.		
BLIGHIA.						
sapida.	Akee.	Africa.	W. Hamilton, Esq. 1797.	T.	♀	♀

TRIGYNIA.

POLYGONUM.						
nutans, R.	[murich.	Bengal.	Indigenous.	☉ E.	R. S.	C. S.
lanatum, R.	B. Shweta-panee-			☉	R. S	C. S.
pilosum, R.	B. Bura-panee-murich.	India.		☉	R. S.	C. S.
tomentosum.	T. Yeati-mallier.	Circars.	Nalla-mundu.	☉	R. S.	C. S.
glabrum.6		India.	Dr. W. Carey.	☉ E.	R.&C.S.	C. S.
tenellum, R.		Bengal.	Indigenous.	☉	R. S.	C. S.
barbatum.	T. Kunda-mallier.	India.	Dr. W. Carey, 1797.	☉	R.&C.S.	C. S.
rivulare, K.7			Indigenous.	☉	R. S.	C. S.
flaccidum, R.	B. Panee-murich.	Bengal.		☉	R. S.	C. S.
perfoliatum.		India.	Dr. Buchanan.	♂ Sc.	♀	♀
horridum.			Mr. J. R. 1810.			
Fagopyrum.		Tibet, &c.	Before 1794.	☉ E.	C. S.	H. S.
aviculari.		Europe.	Accident, 1800.	☉ D.	5---6	
elegans.	H. Kunuk-jhinghi.	India.	Indigenous.	♃ D.	♀	D. S.
COCCOLOBA.						
uvifera.		W. Indies,	W. Hamilton, Esq. 1802.	T.	H. S.	10
PAULINIA.						
carthageniensis.		S. America.	W. Hamilton, Esq. 1800.	♄ Cr.	C. S.	
CARDIOSPERMUM.						
Halicacabum.	B. Shibjhool.8	India.	Indigenous.	☉ Sc.	♀	
Corindum.		Brazil.	W. Hamilton, Esq.	☉ Sc.		

TETRAGYNIA.

ODINA.						
Woodier.	B. Jiyul.9	India.	Indigenous.	L. T.	3--4	9---10

1 Tam. May. T. Yœlim-buriki. 2 Genip-tree of the West Indies. 3 T. Kunkoodoo. 4 H. M. 4. t. 19. 5 S. Aristaphenila. 6 H. M. 12. t. 77. 7 H. M. 12. t. 76. 8 and Noaphutuki. S. Jyotishmutee. T. Nalla-golivinda. 9 S. Jeevula. H. Kashmulla. T. Gumpina.

CLASSIS IX.

ENNEANDRIA MONOGYNIA.

LAURUS.						
I. *Leaves opposite.*						
Cinnamomum.	B. Dar-cheeni.1	Ceylon.	Gen. H. M'Dowall, 1802.	T.	1---2	5---6
broad-leaved.			Before 1794.	T.	1---2	5---6
narrow-leaved.	B. Tej.2	India.	1796.	T.	2---4	6---7
Cassia.			Before 1794.	T.		
Malabathrica.3	Mn. Poheree-lawan.	Moluccas.	Mr. C. Smith, 1800.	T.	3---4	
Culitlaban.4		China.	1796.	T.	4---5	
recurvata, R.	Mn. Coolit manes.5	Sumatra.	Dr. C. Campbell, 1802.	T.	1---2	5---6
nitida, R.	B. Kintun.	Silhet, &c.	H. C. 1810.	T.	1---2	7---8
obtusifolia, R.		China.	1795.	T.	H. S.	R. S.
II. *Leaves alternate.*		Europe.	Before 1794.	b		
dulcis, R.	E. Sweet-bay.	West Indies.	1793.	T.	2---3	7---8
nobilis.	E. Avocado.6	Silhet.	Mr. M. R. Smith, 1811.	L. T.	2---4	5---6
Persea.	B. Orook.		Mr. M. R. Smith, 1811.	T.	2---4	5---6
glaucescens.	B. Sundhigool.					
lanceolaria, R.	S. Kurpoora.7	Sumatra, &c.	Monsieur Cere, 1802.	T.	3---4	
camphorifera.	Mn. Cayoo-gaddees.8	Sumatra.	Dr. C. Campbell, 1803.	T.	C. S.	5
porrecta, R.		America.	W. Hamilton, Esq. 1809.	T.		
Sassafras.		N. America.	W. Hamilton, Esq. 1804.			
chloroxylon.		Virginia.	W. Hamilton, Esq.			
Benzoin.		Tipperah.	Mr. S. Harris, 1797.	T.	2	6
bilocularis, R.						
ANACARDIUM.						
occidentale.	B. Hijilee-budam.	India.	Before 1794.	T.	3---4	R. S.
CASSYTA.						
filiformis.	B. Akash-bulee.9	India.	Indigenous.	P. Cr.		

TRIGYNIA.

RHEUM.						
compactum.		China.	1806.	H.		
palmatum.			1807.			

1 C. or Rasse-coronde. S. Daroositi. This variety of the Cinnamon tree was introduced into Bengal and Tinnevelly while the Dutch were in possession of Ceylon, and is probably Burman's Cinnamomum perpetuo-florens, or one of his inferior varieties mentioned in page 62 and 63 of his Thesaurus Zeylanica. 2 The tree, and Tej-pat, the leaves. S. Twuk-putra. 3 H. M. 5. t. 53. 4 H. A. 2. t. 14. 5 Marsd. Sumat. 125. 6 Or Alligator-pear. 7 A. Kafur. Mn. Kapoor. 8 Marsd. Sumat. p. 129. 9 S. Akasha-vullee. T. Pauncha-tiga.

CLASSIS X.

DECANDRIA MONOGYNIA.

SOPHORA.						
tomentosa.		Ceylon.	1796.	♄ E.	☉	
robusta, R.	B. Peeplur.	Silhet.	Mr. M. R. Smith, 1811.	L. T.	4	7
PODALYRIA.						
bracteata, R.	B. Gopooree.	Silhet.	Mr. M. R. Smith, 1812.	♄ E.	5---6	12---1
capensis.		Cape of G. H.	Capt. Thomas, 1799.	T.		
monosperma.		W. Indies.	A. B. Lambert, Esq. 1806.	T.		
australis.		N. America.	W. Hamilton, Esq. 1798.	♄		
CERCIS.						
canadensis.		N. America.	W. Hamilton, Esq. 1798.	♄		
BAUHINIA.						
variegata.	B. Rukta-kanchun.1	India.	Indigenous.	T.	H. S.	R. S.
candida.	B. Kana-raj.	India.	Before 1794.	T. E.	H. S.	C. S.
purpurea.	B. Deb-kanchun.2		Mr. Heshusius, 1804.	T.	C. S.	H. S.
triandra, R.		Bengal.	Dr. W. Carey.	T.	H. & C. S.	H. S.
parviflora.	S. Vana-raja.3	Coromandel.	1796.	T.	5---6	C. S.
retusa, R.		Bengal.	Before 1794.	T.	R. & C. S.	H. S.
acuminata.	B. Kanchun.	India.	1794.	♄ E.	☾	☾
tomentosa.4	Tam. Ati.	Malabar.	W. Hamilton, Esq.	♄ E.	H. & R. S.	C. S.
racemosa.	H. Mawal.5	India.	Before 1793.	L. Sc. T.	H. S.	C. S.
anguina, R.	B. Nag-poot.	Garrow-hills.	Mr. M. R. Smith, 1811.	L. Sc. ♄	R. S.	C. S.
corymbosa, R.		China.	Mr. W. Kerr, 1796.	♄ Sc.	H. S.	R. S.
semibifida, R.		Sumatra.	Mr. W. R. 1804.	♄ Sc.	10---11	4---5
malabarica, R.		Malabar.	M. Brown, Esq. 1808.	T.	10---11	
scandens.	B. Gundha-gila.	India.	Mr. M. R. Smith, 1812.	♄ Sc.	3---4	10
HYMENÆA.						
Courbaril.		America.	W. Hamilton, Esq. 1805.	T.		
PARKINSONIA.						
aculeata.		America.	W. Hamilton, Esq. 1797.	T.	H. S.	R. S.
CASSIA.						
exigua, R.		Bengal.	Dr. F. Buchanan, 1812.	☉	8---10	C. S.
Absus.		Coromandel.	1797.	♂	☾	☾
bacillaris.		Surinam.	Sir G. Younge, 1801.	L. ♄	C. S.	R. S.
Tora.	B. Chakoonda.6	India.	Indigenous.	☉	☾	☾
toroides, R.	Can. Taga-shay.	Mysore.	Dr. F. Buchanan, 1801.	☉ E.	R. S.	C. S.
bicapsularis.		Uncertain.	Accident, 1799.	L. Sc. ♄	9---10	D. S.
purpurea, R.	B. Kala-kulkashinda.7	India.	Indigenous.	☉ E.	R. & C. S.	D. S.
occidentalis.				☉ E.	☾	☾
obtusa.8		Mysore.	Dr. F. Buchanan, 1801.	♂ D.	☾	☾
Fistula.*	B. Sonali.9	India.	Indigenous.	T.	5---6	2---3
rhombifolia,* R.		Ceylon.	Gen. H. M'Dowall, 1802.	T.	5---6	2---3
arborescens.10		India.	Before 1794.	S. T.	R. & C. S.	D. S.
Senna.	B. Sona-pat.11	Arabia, &c.	Dr. W. Carey, 1800.	♂ E.	R. & C. S.	D. S.
esculenta, R.	B. Kas-murdun.12	Circars, &c.	1801.	☉ E.	C. S.	H. S.
Sophora.13	B. Kulkashinda.	India.	Indigenous.	☉ E.	R. & C. S.	D. S.
speciosa, R				S. T.	☾	D. S.
sumatrana, R.		Sumatra.	Dr. C. Campbell, 1798.	T.	R. & C. S.	D. S.
nodosa.*		Chittagong.	Dr. F. Buchanan, 1798.	T.	4---5	C. S.
auriculata.	T. Tangheroo.	India.	1797.	L. ♄.	10---12	D. S.
bacillus, Gært.*		Malay Islands	Dr. C. Campbell, 1800.	T.	H. S.	C. S.
alata.	B. Dadoo-murdun.14	Uncertain.	Before 1794.	L. ♄	9	C. S.
marginata,* R.		Coromandel.	Dr. A. Berry, 1800.	T.	R. S.	C. S.
glauca, R.			Dr. A. Berry, 1800.	S. T.	10---11	3---4
marilandica.		N. America.	W. Hamilton, Esq. 1798.	♃ E.	R. & C. S.	D. S.
grandis.		Surinam.	W. Hamilton, Esq. 1800.	T.	3---5	2---4

1 S. Kovidara, Yooga-putra, (i. e. double-leaved). 2 H. Sona. 3 Tam. Areka. marum. T. Arroe. 4 H. M. 1. t. 35. 5 T. Ada. 6 S. Chukra-murda. T. Chacunda. 7 T. Kunda-kashinda. 8 Indian senna, and used as such by the natives. 9 S. Soovurnuka. H. Amultus. T. Rela. 10 H. M. 6. t. 9 & 10. 11 A. Suna or Sena. H. Sona-mookhee. 12 T. Nitty-kashinda-kura. 13 H. M. 2. t. 52. 14 S. Dudrooghna. T. Wandukolli.

* These five might, I think, with much propriety be formed into a separate genus under the following, or some such essential character, Calyx pentaphyllous, Petals five, subequal. Filaments very unequal, lower three with a double curve below the middle. Anthers erect, with two slits on the front. Loment cortical, cylindric, many-celled. They are all trees of great beauty, particularly while in flower. The leaves pinnate; no glands on the common petioles, but stipulate at the base; flowers on axillary racemes; calyx of five equal leaflets; corol of five nearly equal petals; stamina very unequal, the three lower filaments much longer than the rest, with a double curve below the middle, and in two of them a large swelling at the middle; loment cortical, cylindric, pendulous, many-celled, with one seed in each cell; embryo straight, furnished with an hard perisperm.

frondosa.		W. Indies.	W. Hamilton, Esq. 1800.			
ligustrina.		N. America.	W. Hamilton, Esq. 1800.	♄		
Chamæcrista.		Cuba.	W. Hamilton, Esq. 1804.	☿ E.	○	○
prostra a, K.	T. Nalla Teeloога.	India.	Indigenous.	☿ D.	R. S.	C. S.
sensitiva, R.		Coromandel.		☿ D.	R. S.	C. S.
dimidiata.		Napaul.	Dr. F. Buchanan, 1801.	☾ E.	R. S.	C. S.
tenella, R.		Bengal.	Dr. W. Carey.	♂ E.	R. S.	C. S.
nictitans.		N. America.	Dr. W. Carey, 1809.	☾ E.	R. S.	C. S.
aurata, R.		Malayana.	1809.	♄ E.	R. S.	C. S.
GYMNOCLADUS.						
canadensis.			Mr. M'Mahon.			
POINCIANA.						
pulcherrima.	H. & B. Krishna-choora.	India, &c.	Before 1794.	♄	○	○
elata.			Dr. Berry, 1799.	T.	○	○
CÆSALPINIA.						
Sappan.	B. Bukum.1	India.	Before 1794.	S. T.	H. S.	C. S.
Bonduc.		Sumatra.	Dr. C. Campbell, 1802.	L. sc. ♄		
Bonducella, R.	B. Nata.2	India.	Indigenous.	L. Sc. ♄	R. S.	C. S.
Olæosperma, R.	B. Umul-koochi.3	[Ganges.		L. Sc. ♄	R. S.	C. S.
cuculata, R.		Delta of the	Dr. W. Carey.	L. Sc. ♄	R. S.	C. S.
Simora.		Mysore.	Dr. Buchanan.	Sc. ♄	C. S.	H. S.
sepiaria, R.	Mysore thorn.		General Martin.	L. Sc. ♄	R. & C. S.	R. S.
resupinata, R.		Moluccas.	Mr. Smith.	Sc. ♄	R. S.	C. S.
enneaphylla, R.		Chittagong.	Dr. Buchanan.	Sc. ♄	R. & C. S.	H. S.
paniculata, 4 R.		India.	1800.	L. Sc. ♄	C. & H. S.	R. S.
chinensis, R.		China.	Mr. W. Kerr, 1807.	L. Sc. ♄		
tortuosa, R.		Sumatra.	Dr. C. Campbell, 1797.	L. Sc. ♄	2---3	R. S.
sumatrana, R.			Dr. Campbell.	L. Sc. ♄	H. & R. S.	C. S.
polyphylla, R.		Pegu.	Mr. F. Carey, 1810.	L. Sc. ♄		
GUILANDIA.						
Bonducella.5						
HYPERANTHERA.						
Morunga.	B. Shojina.6	India.	Indigenous.	T.	H. S.	C. S.
CYNOMETRA.						
cauliflora.	Mu. Nam Nam.	Sumatra.	Dr. C. Campbell, 1800.	T.		
polyandra, R.	B. Peng.	Silhet.	Mr. M. R. Smith, 1811.	L. T.	3---4	7---8
GÆRTNERA.	[luta.7					
racemosa.	H. & B. Madhubee-	India.	Before 1794.	L. Sc. ♄	R. & C. S.	C. S.
obtusifolia, R.		China.	1794.	Sc. ♄	2---4	R. S.
ADENANTHERA.						
pavonina.	S. & H. Koochunduna.8	India.	Before 1794.	T.	H. S.	R. & C. S.
HÆMATOXYLON.						
campechianum.	E. Log-wood.	S. America.	Monsieur Cere, 1810.	T.	12---1	3---4
PROSOPIS.						
spicigera.	S. Shumee.9	India.	Captain Wilford, 1800.	S. T.	C. S.	R. S.
RUTA.						
graveolens.	E. Rue.10	Europe & Asia.	1800.		C. S.	H. S.
MURRAYA.						
exotica.11		India, China.	Before 1794.	S. T.	H. & R. S.	C. S.
sumatrana, R.		Sumatra.	Dr. C. Campbell, 1801.	L. ♄	○	○
BERGERA.						
Kœnigii.	H. Bursunga.12	India.	Mr. B. Heyne, 1800.	S. T.	C. S.	R. S.
integerrima.	B. Bun-koonchа.	Bengal.	Dr. F. Buchanan, 1797.	S. T.	C. S.	R. S.
LIMONIA.						
monophylla.	T. Udivi-nima.	Coromandel.	Dr. Rottler, 1803.	♄ E.	10---11	
bilocularis, R.			Dr. A. Berry, 1807.	S. ♄	5---6	8---9
citrifolia, R.		Chittagong.	Mr. J. R. 1810.	♄	H. S.	
scandens, R.	S. Luvunga-luta.	Silhet.	R. K. Dick, Esq. 1805.	Sc. ♄	H. S.	9---10
pentaphylla.	B. Ash-shoura.13	India.	Indigenous.	♄ E.	○	○
crenulata, R.	T. Torellega.			S. T.	H. S.	R. S.
pentagyna, R.	T. Chitreka.	India.	S. Harris, Esq. 1796.	T.	H. S.	R. S.
COOKIA.						
punctata.	Ch. Whong-pi.14	China.	Before 1794.	T.	H. S.	R. S.
BOSWELLIA.						
thurifera.	H. Salace.15	Coromandel.	Dr. Berry, 1803.	L. T.	2---4	R. S.
BUCHANANIA.						
latifolia, R.	S. & B. Piyala.16	India.	H. Colebrooke, Esq.	T.	2---3	R. S.
angustifolia, R.		Mysore.	Dr. F. Buchanan, 1801.	T.	5---6	3---4
lanceafolia, R.		Chittagong.	Mr. J. R. 1810.	L. T.		
WALSURA.						
piscidia, R.	T. Wallursi.17	Coromandel.	P. Heshusius, Esq. 1800.	S. T.	C. S.	
robusta, R.	B. Uphing.	Silhet.	Mr. M. R. Smith, 1812.	L. T.	3---4	6---7

1 S. Potanga. H. P. & A. Bukum. 2 S. Pootikarunja. H. Katkuleja. C. Komboorel etta. 3 T. Noonі-ghietsa. 4 H. M. 6. t. 19. 5 And paniculata, see Cæsalpinia. 6 S. Sobhanjuna. Tam. Morunga. 7 S. Utimookta, and Madhuveeluta, 8 B. Ructa-chunduna or Ranjuna. Can. Mandateea. T. Manseni-kotta. 9 T. Shumee. 10 S. Brahmi, also Soma-luta. H. Saturi, also Allooda. A. Sudab. 11 Chalcas paniculata. Mant. is the same plant. 12 T. Kari-vepa. Tam. Kari-pilli. 13 S. Ushwa-shakhota. T. Gollunga. 14 or Wampee. 15 S. Shullukee. B. Luban. E. Olibanum tree. 16 Tam. Moreeda. T. Tsaroo-mamadi 17 Tam. Walsura.

HEYNIA. trijuga, R.		Napaul.	Dr. F. Buchanan, 1801.	T.	9	10
EKEBERGIA. indica, R.	T. Poroodonda.1	India.	H. Russell, Esq. 1802.		O	O
SANDORICUM. indicum, R.	Mn. Sandoricum.	Malacca,&c.	Before 1794.	T.	2---3	9---10
MELIA. Azadirachta.	H. & B. Nim.2	India, &c.	Indigenous.	T.	2---3	5---6
sympervirens.	H. Bukain.3	Indies.	Before 1793.	S. T.	O	O
Azederach.		India, China.	1798.	L. T.	3	11---12
superba, R.		Soonda.	Dr. Berry, 1804.	L. T.	2---3	10
robusta, R.		Malabar.	Dr. Berry, 1802.	L. T.	3---4	11---12
SWIETENIA. Mahogony.	E. Mahogony.	West Indies.	H. C. 1795.	L. T.	H. S.	
febrifuga, R.	H. B. Rohun.4	India.	1791.	T.	3---4	7---8
chickrassa, R.	B. Chukrasi.		Dr. J. Fleming.	L. T.	H. S.	R. S.
chloroxylon, R.	T. Billoo.5		Mr. Heyne, 1801.	T.	3---4	C. S.
GARUGA. pinnata, R.	B. & H. Toom.6	India.	S. Harris, Esq. 1798.	L. T.	H. S.	10---1
TRIBULUS. lanuginosus.	B. Gokhoor.7	India.	Indigenous.	♂ D.	O	O
cistoides.		N. America.	W. Hamilton, Esq. 1799.	♂ D.	O	O
JUSSIEUA. repens.	B. Kachura dam.8	India.	Indigenous.	☉ A.	R. S.	C. S.
exaltata, R.	B. Bun-lung.9	India.		♃ E.	R. S.	C. S.
MELASTOMA. aspera.	B. Chota-Phootika.	Bengal.	Dr. W. Carey, 1810.	S. ♄ E.	C. S.	
malabathrica.	B. Bura-Phootika.10	India.	R. W. Cox, Esq. 1795.	♄ E.	O	O
vagans, R.		Chittagong.	Mr. J. R. 1810.	♄ Sc.	5---7	
cernua, R.				♄ Sc.	10---11	2---3
Spec. nov.	B. Juya Phootika.	Silhet.	Mr. M. R. Smith, 1812.	♄ Sc.	R. S.	
GASTONIA. palmata, R.		Chittagong.	Mr. J. R. 1810.	♄ E.		
RHODODENDRON. punicеum, R.	H. Boorans.	Sirinagur.	Col. Hardwicke, 1811.	S. T.	4---5	
CERATOSTEMA. variegata, R.	B. Tala-moot.	Silhet,Garrow.	Mr. M. R. Smith, 1812.	♄ E.	C. S.	6---7
vaccinacea, R.	B. Kesa-proom.	Garrow hills.	Mr. M. R. Smith, 1812.	♄ E.	3---4	6---7
FERONIA. elephantum.	H. & B. Kuth-bel.11	India.	Indigenous.	T.	3	10
STYRAX. Benzoin.		Sumatra.	Dr. Lumsdain, 1812.	T.		
serrulata, R.		Chittagong.	Mr. J. R. 1810.	T.	H. S.	R. S.
INOCARPUS. edulis.12		Moluccas.	H. C. 1798.	T.	H. S.	C. S.
CASEARIA. Vareca, R.	B. Titta cheera.	Silhet.	Mr. M. R. Smith, 1811.	♄ E.	H. S.	9---10
glabra, R.		Moluccas.	H. C. 1798.	L. T.	R. S.	
glomerata, R.	B. Loorjoor.	Silhet.	Mr. M. R. Smith, 1811.	♄ E.	12	3
AQUILARIA. Agallocha, R.	E. Aloe-wood tree.13	Eastern frontier.	R. K. Dick, Esq. 1803.	L. T.	3---4	R. S.
HARDWICKIA. binata, R.	Tam. Acha or atti.14	Coromandel.	Dr. Berry, 1800.	T.		
pinnata, R.			Dr. Berry, 1809.	T.		
GETONIA. nutans, R.		Rajemahl.	Mr. W. R. 1800.	Sc. ♄	C. S.	H. S.
TERMINALIA. Catappa.	H. Budam.	India.	Before 1794.	T.	2---6	R. S.
procera, R.		Andamans.	Major Kyd.	L. T.	2---3	R. S.
Bellerica.	H. & B. Buhura.15	India.	Before 1794.	T.	H. S.	C. S.
Chebula.	B. Huritukee.16	India.	Dr. A. Berry, 1803.	L. T.	H. S.	1---2
moluccana.				T.		
angustifolia.	Can. Mungachi.	Travancore.	Mr. G. Young, 1808.	T.		
Citrina, R.		Silhet, &c.	Mr. M. R. Smith, 1811.	L. T.	H. S.	1---2
gangetica, R.			Kurrum kaun, 1779.	T.	H. S.	C. S.

1 Can. Bettuda-bewoo. 2 S. *Nimba*. T. Vepa. Tam. Vepam. 3 A. Ban. T. Turka vepa. 4 T. Soymida. S. *Rohuna*. Tam. Woomæ marum. 5 E. Satin-wood. Tam. Moodooda. C. Boorooch-gaha. 6 T. Garugo. H. M. 4. t. 33. 7 or Gokhyoor. S. *Gokshooruka*. 8 S. Langulee. H. Canchara. T. Neer-batsola. 9 S. *Bhoo-luvunga*. T. Neeroo-aghiondraparoo. H. M. 2. t. 50. 10 H. M. 4. t. 42. 11 S. *Kupittha*. T. Yellinga. 12 H. A. 1. t. 65. 13 S. *Ugooroo*. H. & B. *Ugoor*. 14 Can. Caratchu. 15 S. *Vibheetukee*. T. Toandi. P. Belele. 16 S. Hureetukee. H. Hur or *Hura*. T. *Kudooke*. P. Halileh. A. Ahlilej.*

* "Under the head *Ahlilej* (the Arabic name answering to the Persian *Halileh*) the author distinguishes the following kinds, as the produce of the same tree (*Terminalia Chebula*) gathered at different degrees of maturity: 1. *Halileh Zira*, gathered when the fruit is just set; being dried, it is about the size of the *Zira* (*Cumin seed*). 2. *H Jawi*, when more advanced it is the size of a barley-corn (*Jaw*). 3. *H. Zengi*, *Hindi* or *Aswed*, when the young fruit is still further advanced; being dried, it is of the size of a raisin, and is black, whence its name (*Aswed black*, or *Zengi negro*). 4. *H. Chini*, gathered when the nut has acquired some degree of hardness; the dried fruit is of a green colour, inclining to yellow. 5. *H. Asfer*, when approaching to maturity: the fruit, when dry, is of a reddish yellow, whence the name (*Asfer yellow*). 6. *H. Cabuli*, when the fruit is come to full maturity." *Fleming in Asiatic Researches.*

PENTAPTERA.						
tomentosa, R.	M. & B. *Usun.*1	India.	Before 1793.	L. T.	H. S.	C. S.
glabra, R.	T. Della-madoo.		1803.	L. T.	H. S.	C. S.
Arjuna, R.	B. *Urjoon.*2		Before 1793.	L T.	H. S.	C. S.
Coriacea, R.	Tam. Anemutti.	Coromandel.	Dr. Berry.	L. T.	H. S.	C. S.
crenulata, R.		India.	Dr. Berry.	L. T.	H. S.	C. S.
bialata, R.			Before 1793.	L. T.	H. S.	C. S.
paniculata, R.	T. Nimmeeri.3	Coromandel.	Dr. Berry, 1803.	L. T.	11---12	5---6
angustifolia, R.		Balla-gaut.	Dr. A. Berry, 1800.	L. T.	H. S.	C. S.
CONOCARPUS.						
latifolia, R.	T. Sirimaun.	Coromandel.	Dr. A. Berry, 1800.	T.	C. S.	H. S.
acuminata, R.	T. Panch-maun.		Dr. A. Berry, 1803.	T.	C. S.	H. S.

DIGYNIA.

HYDRANGIA.						
hortensis.		China.	Mr. W. Kerr, 1812.	♄ E.		
TRIANTHEMA.						
obcordata, R. white.	H. Shwet Sabuni.	India.	Indigenous.	⊙ D.	R.&C.S.	C. S.
red.	H. Lall Sabuni.			⊙ D.	R.&C.S.	C. S.
decandra.	B. Gadha-buni.			⊙ D.	R.&C.S.	C. S.
crystallina.	T. *Kooka-pel-koora.*			♃ E.	R.&C.S.	C. S.
SAPONARIA.						
perfoliata, R.		Bengal.	Accident.	⊙ E.	C. S.	R. S.
DIANTHUS.	[alnan.					
caryophyllus, single.4	P. Gool-*kurunphul*	Europe, Persia.	Dr. W. Carey, 1800.	♃	H. S.	
double.5			Mrs. Birch, 1808.	♃	H. S.	
chinensis.	China-pink.	China.	H. C. 1796.	♂	C. S.	H. S.
barbatus.	Sweet-William.	Europe.	H. C. 1802.	♃ E.		
superbus.				♃ E.	○	

TRIGYNIA.

SILENE.						
indica, R.		Napaul.	Dr. F. Buchanan, 1802.	⊙ E.	3---5	4---6
bellidifolia.		Cape of G. H.	1798.	⊙		
antirrhina.		Virginia.	Mr. Lambert, 1802.	⊙	2	
nutans.		Europe.	Dr. Hornman, 1807.	♃		
chlorantha.			Dr. Hornman, 1807.	♃		
Atocion.			Sir A. Hume, 1812.			
MALPIGIA.						
lucida.		W. Indies.	W. Hamilton, Esq. 1802.	L. ♄	3---5	
HIRAEA.						
nutans, R.		Bengal.	Dr. F. Buchanan.	Sc. ♄	7---8	11---12
ERYTHROXYLON.						
monogynum, R.	T. *Udivi-gorinta.*	Coromandel.	Gen. M'Dowall, 1802.	S. T.	R.&C.S.	C. S.
BANISTERIA.						
auriculata.		S. America.	Dr. Wallick, 1808.	T.	1---7	
STELLARIA.						
dichotoma.		N. America.	Accident, 1803.			
ARENARIA.						
flaccida, R.		Bengal.		⊙ D.		

PENTAGYNIA.

AVERRHOA.						
Carambola, acid.	S. *Kurmu-runga.*6	India.	Before 1794.	S. T.	H.&R.S.	C. S.
sweet.	H. & B. Meetha Ka-		1794.	S. T.	H.&R.S.	C. S.
Bilimbi.7	[*ma-runga.*		1800.	S. T.	H. S.	R. S.
SPONDIAS.						
axillaris, R.	N. Neema.	Napaul.	Dr. F. Buchanan, 1802.	T.	3---4	R. S.
mangifera.	H. & B. Amra.8	India.	Indigenous.	T.	3	12---1
dulcis.	E. Otahiete apple.	Otaheiti.	Captain B. Blake, 1801.	T.	3	12---1
acuminata, R.		Malabar.	Mr. G. Young, 1808.	T.		
lonifolia, R.			Monsieur Cere, 1810.	T.		
CNESTIS.						
monadelpha, R.	B. *Sookur-thooti.*	Silhet.	Mr. M. R. Smith, 1811.	L. ♄	6---7	10
COTYLEDON.						
laciniata.	S. *Hema-sagura.*9	India.	Before 1793.	H. E.	R. S.	C. S.
rhizophylla, R.		Moluccas.	Lady Clive, 1799.	♃ E.	C. S.	H. S.
heterophylla, R.		Mysore.	Dr. F. Buchanan, 1801.	♃ E.	C. S.	H. S.
BERGIA.						
verticillata.	H. & B. Lal-keshura.10	India.	Indigenous.	⊙ Cr.	R. S.	C. S.
ammannioides, R.	H. Dhoolacheri.11			⊙ E.	R. S.	C. S.
OXALIS.						
corniculata.	H. & B. Amrool.12	India.	Indigenous.	⊙ Cr.	○	C. S.
caprina, single & double.		Cape of G. H.	1799.	H.	C. S.	

1 S. *Usuna.* T. Nellamadoo. 2 S. *Urjoona.* *Kukoobha.* H. Kahoowa. 3 Tam. Po-
karakai. 4 or clove-pink. 5 or carnation. 6 H. & B. *Kama-runga.* 7 H. M. S.
t. & 45 46. 8 S. *Amratuka.* T. Awatum. 9 H. & B. Hemsagur. 10 T. Nee-
roo-pavala. 11 Lechea verticillata, Willd. 12 S. *Umla-lonika, chookrika.*

sensitiva.	H. Lucbhanno.1	India.	Indigenous.	☿ E.	O	O
flava.	Single & double.	Cape of G. H.	1799.	H.		
purpurea.			1799.	H.		
hirta.			1799.	H.		
sessilifolia.			1799.	H.		
versicolor.			1799.	H.		
violacea.		America.	A. Roberts, Esq. 1811.		C. S.	
CERASTIUM.						
cordifolium, R.		Bengal.		⊙ D.	C. S.	H. S.
vulgatum.		Europe.	W. Hamilton, Esq. 1800.	⊙ D.	3	
SPERGULA.						
arvensis.		Europe.	W. Hamilton, Esq.	⊙ D.	C. S.	C. S.
AILANTHUS.						
excelsa, R.		Coromandel.	1803.	L. T.	C. S.	

DECAGYNIA.

PHYTOLACCA.						
octandra.		Mexico.	Mr. M'Mahon, 1809.	H. E.	D. S.	H. S.
decandra.		America.	W. Hamilton, Esq. 1801.	H. E.	D. S.	H. S.
acinosa, R.		Napaul.	Dr. F. Buchanan, 1802.	H. E.	C.&H.S.	H. S.

1 B. Bun narunga.

CLASSIS XII.

DODECANDRIA MONOGYNIA.

AGATHOPHYLLUM.						
aromaticum.1		Madagascar.	Captain Tennent, 1802.	T.		
TOMEX.2						
CASSIA.3						
RHIZOPHORA.						
conjugata.		India.	--- Goodlad, Esq. 1796.	T.		
gymnorhiza.	B. Kakra.4		--- Goodlad, Esq.	T.	H. S.	C. S.
cylindrica.		Sunderbunds.	--- Goodlad, Esq.			
Mangle.	B. Bhorar.5	India.	--- Goodlad, Esq.	T.	H. S.	C. S.
decandra, R.	B. Guran.		--- Goodlad, Esq.	T.	H. S.	C. S.
candel.6			Dr. F. Buchanan.	T.		
GARCINIA.7						
CANELLA.						
alba.		West Indies.	W. Hamilton, Esq. 1804.	T.	H. S.	
CRATEVA.8						
TRIUMFETTA.						
Bartramia.	H. & B. Bun-okra.9	India	Indigenous.	⊙ E.	R.&C.S.	C. S.
semi-triloba.		West Indies.	W. Hamilton, Esq.			
trilocularis, R.		India.	Accident, 1803.	♄ E.	10---11	1---2
TRIUMFETTA.						
annua.		India.	Indigenous, 1802.	⊙	1---4	
orbiculata.		Circars.	Naila-mandu, 1803.		9---10	C. S.
PORTULACCA.						
Oleracea, 3 or 4 var.	H. & B. Noonya.10	India.	Before 1794.	⊙ E.	R. S.	C. S.
meridiana.	H. & B. Noonya.11		Indigenous.	⊙ Cr.	☾	○
quadrifida.	T. Pedda Poil-pail-kura.			⊙ Cr.	○	○
pilosa.	H.&B.Bura-noonya.			⊙ E.	○	○
linifolia, R				⊙ D.	○	○
TALINUM.						
cuneifolium.		India.	Dr. Heyne, 1804.	H. E.	R.&C.S.	C. S.
LYTHRUM.12						
KLIENHOVIA.13						
STERCULIA.14						

TRIGYNIA.

RESEDA.						
Luteola.		Europe.	H. C. 1800.	⊙ D.	C. S.	H. S.
odorata.	E. Mignonette.		Before 1794.	⊙ D.	C. S.	H. S.
undata.			Accident, 1808.	⊙ E.	C. S.	H. S.
EUPHORBIA.						
antiquorum.	H. & B. Nara-shij.15	India, &c.	Indigenous.	S. T.	C. S.	H. S.
arborescens, R.			1800.	S. T.		
lactea, R.		Moluccas.	H. C. 1798.	S. T.		
trigona, R.			Mr. Smith.	♄ E.	H. S.	
nereifolia.	B. & H. Shij.	India.	Indigenous.	T.	3---4	
Ligularia, R.	B. & H. Munsa-shij.		Before 1793.	S. T.	C. S.	
Tirucalli.	B. Lunka-shij.16	Coromandel.			R. S.	
sessiliflora, R.		Pegu.	Mr. F. Carey, 1808.	♃	2---3	
hirta.	B. Bura-kheerooee.	India.	Indigenous.	☾ E.	○	○
uniflora.				⊙ D.	○	○
thymifolia.	B. Shwet-kheerooee.			⊙ D.	○	○
Chamaesyce.	H. Choota-kheerooee.			⊙	○	○
parviflora.				⊙	○	○
tithimalbides.		America.	Before 1794.	♄ E.	H. S.	
dracunculoides.	H. Chagul-putputee.	India.	Dr. Buchanan.	⊙ E.	C. S.	H. S.
peltata, R.		Circars.	1794.	⊙ E.	C. S.	H. S.
hyberna.		Siberia, &c.	W. Hamilton, Esq.			
maculata.		N. America.	W. Hamilton, Esq.	⊙		
prunifolia.		Peru.	Dr. Jacquin, 1806.	⊙ E.	○	○
peplis.		Europe.	Dr. W. Carey, 1813.	⊙ E.		
acaulis, R.		Bengal.	B. Smith, Esq. 1810.	H.	3	

1 Ravansara. Sonnerat. 2 See Tetranthera. 3 See Polyandria Monogynia. 4 H. M. 6. t. 30 & 31. 5 T. Upoo-punna. 6 H. M. 6. t. 25. 7 See Polyandria Monogynia. 8 See Feronia & Aegle. 9 H. A. 6. t. 25. f. 2. 10 S. Lonika. A. Tooruk. 11 T. Pail-kura. H. M. 10 t. 31. 12 See Grislea. 13 See Monadelphia. 14 See Monadelphia. 15 S. Sihoonda. 16 Tam. Tirucalli.

CLASSIS XIII.

ICOSANDRIA MONOGYNIA.

CACTUS.						
mamillaris.	Prolifer.	W. Indies.	W. Hamilton, Esq. 1800.	♄	D. S.	
cylindricus.			Kew Garden, 1806.	♄ D.		
tetragonus.			W. Hamilton, Esq. 1801.		R. S.	
hexagonus.		Surinam.	W. Hamilton, Esq. 1799.	♄ E.	R. S.	
peruvianus.		Peru.	W. Hamilton, Esq.	♄ E.		
Royeni.		America.	W. Hamilton, Esq.	♄ E.		
grandiflorus.		W. Indies.	W. Hamilton, Esq. 1801.	♄ Cr.	R. S.	
flagelliformis.		America.	W. Hamilton, Esq. 1801.	♄ Cr.		
triangularis.		W. Indies.	Dr. Berry, 1795.	♄ Cr.	R. S.	
coccinellifer.		S. America.	Dr. Anderson.	♄ E.	R. S.	
chinensis, R.		China.	Before 1793.	♄ E.		
Opuntia.		America.	W. Hamilton, Esq. 1793.	♄ D.		
curassavicus.		Curassao.	Kew Garden, 1803.	♄ E.	R. S.	
Ficus indica.		America.	W. Hamilton, Esq.		R. S.	
Tuna major.			W. Hamilton, Esq.	♄ E.	R. S.	
Tuna elatior.	E. Prickly Pear.	S. America.	Dr. J. Anderson, 1801.	♄ E.	R. S.	R.&C.S
indicus, R.	H.&B.Nag-phunee.1	India.	Indigenous.	♄ E.	☋	☋
Pereskia.		America.	W. Hamilton, Esq.	♄ Sc.		
METROSIDEROS.						
vera.2		Moluccas.	Captain Anderson, 1802.	T.	7---8	9---0
linearis.		N. S. Wales.	Colonel Paterson, 1800.	♄ E.	5---6	C. S.
PSIDIUM.						
pyriferum.	B. Peyara.3	W. Indies.	Before 1794.	T.	☋	☋
pomiferum.	B. Peyara.4	W. Indies.	1794.	T.	☋	☋
guinensis.		Guinea.	Sir G. Young, 1801.	♄	H. S.	R. S.
CARALLIA.						
lanceofolia.		Sumatra.	Dr. C. Campbell, 1803.	T.	H. S.	
EUGENIA.						
malaccensis.	B. Malacca jamrool.	Moluccas.	Before 1794.	T.		
purpurea.			H. C. 1797.	T.		
amplexicaulis.		Chittagong.	Mr S. Harris, 1799.	T.	☋	☋
Jambolana.						
obtusifolia, R.		Amboyna.	1798.	T.	H. S.	
operculata, R.		Moluccas.	Before 1794.	T.	3---4	5---6
caryophyllifolia, R.	B. Chota jamb.	India.	Indigenous.	L. T.	3---4	5---6
fruticosa, R.	B. Bun jamb.	Chittagong.	Dr. F. Buchanan, 1796.	S. T.	3---4	6---7
brachiata, R.		Moluccas.	H. C. 1798.	S. T.	5	7
claviflora, R.	B. Lumba-nuli-jamb.	Chittagong.	Mr. J. R. 1811.	T.	2---3	5---6
myrtifolia, R.		Sumatra.	Dr. C. Campbell, 1803.	S. T.	3---4	5---6
bracteata,	H. Hijulee-mindee.5	India.	Before 1794.	L. ♄	☋	☋
venusta, R.		Tipperah.	S. Harris, Esq.	S. T.		
ternifolia, R. 2 var.	H. Phul-jamb.6	Chittagong.	Mr. J. R. 1810.	T.	4	6---7
acuminata, R.		Amboyna.	H. C. 1798.	T.	3	6---7
cymosa, R.	H. Buti-jamb.	Silhet.	Mr. M. R. Smith, 1811.	L. T.	12---1	
alba, R.	B. Jamrool.	Malayana.	Before 1794.	T.	H.& R. S.	R. S.
aquea, R.	B. Jamb.7	Moluccas.	H. C. 1798.	T.	H.& R. S.	R. S.
oblata, R.	H. Goolum.	Silhet.	Mr. M. R. Smith, 1811.	T.	3	6---7
lanceæfolia, R.	H. Poora-jam.	Silhet.	Mr. M. R. Smith, 1812.	L. T.	11	2
lanceælaria, R.	H. Panee-jamb.		Mr. M. R. Smith, 1811.	S. T.	4---5	
Jamboo.	B. & H. Gulab-jamb.8	India.	Indigenous.	T.	C. S.	R. S.
caryophyllata.	S. Luvunga.9	Moluccas.	H. C. 1797.	T.		
pulchella, R.		Moluccas.	H. C. 1798.	T.	3	R. S.
inophylla, R.			H. C. 1797.	T.	6---7	9---10
glandulifera, R.		Sumatra.	Dr. C. Campbell, 1804.	♄ E.		
macrocarpa, R.		Chittagong.	Mr. J. R. 1811.	T.		
corymbosa.						
MYRTUS.						
communis.	B. Vilati Mindi.		Before 1794.	♄ E.	☋	
tomentosa.		China, &c.	H. C. 1798.	♄ E.		
Pimenta.						
long-leaved.10		W. Indies.	Mr. W. Aiton, 1794.	T.	3---4	7---8
broad-leaved.11		W. Indies.	Hon. Sir J. Banks, 1795.	T.	3---4	7---8

1 Phuni-munusa. Tam. Nagakali. 2 H. A. 3. t. 7. 3 E. White guava. H. Suffri-am.
4 E. Red guava. H. Lal-suffriam. 5 T. Garoog-moodi. 6 White and Lall Phul-jamb
7 The rose coloured, and ayer, the white. 8 Also Jamau. S. Jambu. 9 H. & B. Lung.
A. Kurunful. P. Meykoo. 10 Pimento or Jamaica all-spice. 11 Pimento or Jamaica all-spice.

CALYPTRANTHES.1						
PUNICA.						
granatum.2	B. Dalim.	Persia, &c.	Before 1794.		○	○
double-flowered.	H. Gool-anar.		1794.	♄ E.	R. S.	
nana.		W. Indies.	W. Hamilton, Esq. 1800.	S. ♄ E.		
ALLANGIUM.						
hexapetalum.	B. Bagh-ankra.3	India.	Indigenous.	T.	H. S.	R. S.
SONNERATIA.						
acida.4	B. Our-chaka.	India.		T.	H.&R.S.	C. S.
apetala.	B. & H. Kheora.		Indigenous.	T.	H. S.	
AMYGDALUS.						
persica.	P. Shuft-alloo.5	China, &c.	Before 1794.		1---2	5
Flat peach.	Ch. Pin-to.	China.	1794.		1---2	5
Nectarine.		Persia.	Dr. W. Carey, 1800.		1---2	6---7
communis, sweet.	Almond.6					
bitter.	Almond.					
cordifolia, R.		China.	Before 1793.	T.	C. S.	H. S.
PRUNUS.						
armeniaca.						
of Europe.	Europe Apricot.7		Before 1794.	S. T.	C. S.	H. S.
of China.	China Apricot.8	China.	Before 1794.	S. T.	C. S.	H. S.
Cerasus.	A. Kerosya.9	Asia, &c.				
triflora, R.	Ch. Hong-sum-li.	China, &c.	Mr. W. Kerr, 1811.	L. ♄	2	5---6
CHRYSOBALANUS.						
Icaco.		America.	W. Hamilton, Esq.	T.		
LAGERSTRŒMIA.						
indica.	H. Telinga-china.	China.	Before 1794.	S. T.	R. S.	
reginæ, 2 or 3 var.	B. Jarool.10	India.	1794.	L. T.	H. S.	8---9
grandiflora, R.	S. Indradroog.	Chittagong.	Mr. Moddern, 1801.		D. S.	7---8
parviflora, R.	T. Chinamghie.	India.	Dr. Heyne, 1802.	T.	H. S.	8---9
LUDIA.						
spinosa, R.		Sumatra.	Dr. C. Campbell, 1804.	S. T.	5---6	9---10
fœtida, R.	M-n. Somer-meera.	Amboyna.	H. C. 1798.	T.	R. S.	

DIGYNIA.

CRATÆGUS.						
crenulata, R.		Napaul.	Dr. Buchanan, 1802.	L. ♄	4---5	8
indica.			Mr. W. Kerr, 1812.			

PENTAGYNIA.

MESPILUS.						1
japonica.	Ch. Lo-quat.	China.	Before 1793.	T.	R.S.&12-1	9
bengalensis, R.		Chittagong.	Mr. J. R. 1810.	L. T.	12---1	6---7
PYRUS.						
communis.	P. Nash-patee.11		1799.			
chinensis.	Ch. Sha-li.12	China.	Before 1794.	T.		
Cydonea.	H. Behee or Abee.13		1794.			
Malus.	S. Seva.14		1794.		○	
japonica.	Ch.Tut-chi-hoi-tong.	Japan.	Mr. W. Kerr, 1811.	T.	H. S.	
spectabilis.	Ch. Sai-fou-hoi-tong.	China.	Mr. W. Kerr, 1812.			
indica, R.		Garrow hills.	Mr. M. R. Smith, 1809.	S. T.		
SPIRÆA.						
corymbosa.	C. La-poy.	China.	Before 1794.	♄ E.	○	

POLYGYNIA.

ROSA.						
triphylla, R.	Ch. Cha-te-qui-fa.	China.	Before 1794.	♄ Sc.		
indica.		Rohilcund.	J. Brooke, Esq. 1811.	♄ E.		
inermis.	Ch. Pah-mo-li.15		Mr. W. Kerr, 1812.	♄		
recurva, R.		Napaul.	Dr. F. Buchanan, 1802.	♄		
centifolia.	H. & B. Goolab.16	Asia.	Before 1794.	♄	○	
chinensis.	H.&B.Kanta Goolab.	China.	1794.	♄ E.	C. S.	
semperflorens.			J. Harington, Esq. 1803.	S. ♄	○	
glandulifera, R.	H. & B. Shewuti.17	India.	Before 1794.	Sc. ♄	○	
pimpinellifolia.		Europe.	Mr. M'Mahon, 1810.			
involucrata, R.		Bootan.	Dr. W. Carey, 1797.	♄	C. S.	H. S.
microphylla, R.	Ch. Hai-tong-hang.	China.	Mr. W. Kerr, 1812.	♄		
procumbens, R.			Mr. W. Kerr, 1810.	♄ D.		
RUBUS.						
hirtus, R.		Mysore.	Dr. Berry, 1801.	♄ Sc.	C. S.	

1 See Eugenia. 2 A decoction of the bark of the root is a specific for the tape worm. S. Darimba. P. & H. Anar. A. Rumma. 3 S. *Unkotha*. *Nikochuka*. H. Akarkanta. 4 H. M. 3. t. 40. 5 A. Tuffa-Parsee. Ch. Tow. Conic peach. 6 A. H. & B. Badam. 7 H. & P. Khoobanee. A. Tuffee *Urmenee*. 8 Three varieties. 9 or Jerosya. Ch. Ying-thow. 10 S. *Urjoona*. Burm. Peema. 11 A. *Umrood*. 12 China pear. 13 A. Sufir-jal. 14 P. Seeb. A. Tuffah. 15 The double white, and Wong-mo-ue-he-vong, the double yellow. 16 A. Ward. P. Gool. 17 S. Sevuti.

rosæfolius.		Mauritius.	Capt. Tennent, 1802.	♃ E.	1---2	2---3
paniculatus, R.		Uncertain.	Uncertain, 1799.	♄ E.	○	○
Gowry-phul.	H. Gouree-phul.	Sewalick mts.	Col. Hardwicke, 1802.	♄ Sc.	2---3	
albescens, R.		Malabar.	Capt. Dickenson, 1801.	♄ Sc.	C. S.	H. S.
gracilis, R.		Napaul.	Dr. Buchanan, 1802.	♄ Cr.	C. S.	H. S.
hexagynus, R.		Silhet.	Mr. M. R. Smith, 1810.	♄ Sc.	C. S.	H. S.
FRAGARIA.						
indica, R.		Asam, &c.	Dr. F. Buchanan.	♂ Sar.	C. S.	C. S.
vesca.	Pine Strawberry.	Europe.	Dr. A. Berry, 1808.		12---2	1---3
	Wood Strawberry.		Lady A. Hume, 1808.		1---3	2---4
	Hautboy Strawberry.		Lady A. Hume, 1808.		1---3	2---4
chiloensis.	Chili Strawberry.	S. America.	Capt. Richardson, 1808.		1---5	2---5
maylayana, R.		Pullo-Pinang.	Mr. W. R. 1802.	♂ Sar.	○	○
POTENTILLA.						
reptans.		Europe.	Dr. W. Carey, 1811.	♃ Cr.	C. S.	
argentea.			A. B. Lambert, Esq. 1802.		2	
GEUM.						
urbanum.		England.	H. C. 1806.			
COMARUM.						
flavum.		Napaul.	Dr. F. Buchanan, 1802.		2	
CALYCANTHUS.						
floridus.		Carolina.	W. Hamilton, Esq. 1804.			
præcox.		China.	Mr. Kerr, 1811.	♄ E.	C. S.	

CLASSIS XIV.

POLYANDRIA MONOGYNIA.

TERNSTROMIA.						
serrata, R.	B. Walooa.	Silhet.	Mr. M. R. Smith, 1811.	S. T.	4---5	7---8
BASSIA.						
latifolia, R.	S. Mudhooka.1	India.	Before 1794.	T.	1---2	R. S.
longifolia.	Tam. Illipee.	Coromandel.	Rev. Dr. Rottler, 1800.	T.	H. S.	R S.
SYMPLOCOS.						
spicata, R.	B. Booree.	Silhet.	Mr. M. R. Smith, 1812.	T.	R. S.	C. S.
racemosa, R.	B. Lodh.2	Bengal.	T. Marriot, Esq. 1808.	L. T.	C. S.	H. S.
ferruginea, R.	B. Dookatiya.	Silhet.	Mr. M. R. Smith, 1812.	S. T.	9---10	
DIOSPYROS.						
Kaki.3		China, &c.	Before 1794.	S. T.	2---3	R. S.
Ebenum.		Ceylon, &c.	Dr. A. Berry.	T.	3---5	R. S.
melanoxylon.	B. Kiew.4	Coromandel.	Dr. A. Berry, 1804.	T.	4---5	8---10
tomentosa, R.	B. Tumala.5	Bengal.	Before 1793.	T.	4---5	
glutinosa.	B. Gab.6	India.	Indigenous.	T.	3---4	R. S.
Sapota.7		Mauritius.	Dr. A. Berry, 1804.	T.		
Mabola, R.		Philippines.	Monsieur Jannet, 1811.	T.		
sylvatica, R.	T. Tella-gada.	Circars.	Nalla-mandu, 1803.	S. T.		
racemosa, R.	B. Goolal.	Tipperah.	Before 1794.	T.	3---4	
montana, R.	T. Yerra-gada.	Circars.	Nalla-mandu, 1803.	S. T.	3---4	
stricta, R.		Silhet, &c.	S. Harris, Esq. 1797.	T.	3---4	
chloroxylon, R.	T. Nella-wooly-mera.	India.	Mr. Snodgrass.	S. T.	3---4	
cordifolia, R.	H. Bun-gab.8		Dr. W. Carey.	S. T.	3---4	R. S.
ramiflora, R.	B. *Oori*-gab.	Tipperah.	Before 1794.	T.	3---4	R. S.
MIMOSA.						
I. *Unarmed. Spikes globular.*						
Xylocarpa, R.	P. Pingado.9	India.	Dr. Buchanan.	L. T.	H. S.	8
lucida, R.		N. of Bengal.	Before 1793.	L. T.	H. S.	C. S.
Kœringa.						
Sirissa, R.	S.H.&B.Shireesh.10	India.	Indigenous.	T.	H. S.	H. S.
speciosa.			M. Brown, Esq. 1808.	T.		
heterophylla, R.	B. Kawa-hurunee.	Silhet.	Mr. M. R. Smith, 1811.	T.	2---4	5---6
odoratissima.	T. Shinduga.11	Coromandel.	Dr. Berry, 1800.	T.	5---6	C. S.
elata, R.		Bengal.	Before 1793.	L. T.	5---6	C. S.
Kalkora, R.		Goalpara.	Mr. M. R. Smith, 1810.	L. T.	5---6	C. S.
pulchella, R.		Malabar.	Dr. Berry, 1804.	L. T.		
amara, R.	T. Nella-renga.12	Coromandel.	Dr. Berry, 1797.	T.	H. S.	C. S.
Smithiana, R.	B. Suris.	Silhet.	Mr. M. R. Smith, 1811.	L. T.	5	12
semicordata, R.		Malabar.	M. Brown, Esq. 1808.	T.		
fruticosa, R.	Ch.Shan-yang-thow.	China.	Mr. W. Kerr, 1812.	♄ E.		
stipulata, R.	B. Amlukee.	Bengal.	Dr. W. Carey, 1797.	L. T.	H. S.	C. S.
pedunculata, R.		Malay Islands.	Dr. Berry, 1799.	L. T.		
triquetra.		Coromandel.	Dr. Rottler, 1810.	☿ D.	☾	☾
microphylla, R.	B. Tetooleeya.	Silhet.	Mr. M. R. Smith, 1812.	S. T.	5---6	3---4
biglobosa, R.		Silhet.	Mr. M. R. Smith, 1812.	L. T.	12	4
II. *Unarmed. Spikes cylindric.*						
natans.	H. Pan lajuk.13	India.	Indigenous.	A. Cr.	R.&C.S.	C. S.
adenanthera, R.		Uncertain.	Accident, 1803.	☿ E.	☾	☾
scandens.	T. Gila-tiga.	Indies.	Captain Denton, 1797.	L. Sc. ♄	H. S.	C. S.
III. *Thorny. Spikes globular.*						
dulcis, R.		Manilla.	1792.	L. T.	C. S.	H. S.
corcondiana, R.		Coromandel.	1802.	S. T.	H. S.	C. S.
farnesiana.	B. Gooya-babula.14	India.	Indigenous.	L. ♄	12---1	R. S.
arabica.	H. & B. Babula.15	India.		T.	5---10	3---4
leucophlea, R.	T. Tella-tooma.16	Coromandel.	Dr. Berry, 1799.	T.	R. S.	C. S.
tomentosa, R.	B. Shalshaeei babula.17	India.	Indigenous.	S. T.	C.&H.S.	R. S.
unguis-cati.		West Indies	Hon. C. Greville, 1807.	T.	C. S.	H. S.
IV. *Thorny. Spikes cylindric.*						
dumosa, R.		Hindoostan.		S. T.		
latronum.	T. Peeki-tooma.	Coromandel.	Dr. Berry, 1801.	S. T.	R. S.	C. S.
farnesiana.18		America & Asia.	Before 1793.	T.	C. S.	R. S.
cinerea.	Tam. Wara-taræ.19	Coromandel.	Dr. Berry.	S. T.	H. S.	8

1 T. Ipee. H. & B. Moula. 2 S. Lodhra. 3 C. Chi, several varieties, all edible. 4 T. Tumida. Tam. Tumbali. S. Kakindoo. 5 Tumala. 6 S. Tinduka. T. Tumika. 7 Sapota-nigro, Sonnerat. 8 T. Kawoolimera. 9 T. Konda-tangheroo. 10 Sireessa. T. Dirshena. 11 Tam. Sœlœ-marum. 12 Tam. Shikrœn. 13 T. Neeroo-tavalapoo. B. Lajoolshya. 14 S. Urimeda. T. Kustoori, or Peetooma. 15 S. Barbura. T. Nella-tooma. 16 Tam. Veel-velum. 17 Tam. Kodi-velœ. 18 Gleditsia Linn. 19 S. Yeltoor.

V. Prickly. Spikes cylindric.						
ferruginea, R.	T. Woni.	Coromandel.	Mr. Heshusius, 1801.	T.	H. S.	C. S.
Sundra, R.	T. Sundra.	Coromandel.	Mr. Snodgrass.	T.	R. S.	C. S.
Suma, R.	B. Shaee.1	Bengal.	Indigenous.	T.	R. S.	C. S.
Catechu.	B. & H. Khuera.2	Bengal.	Mr. R. W. Cox, 1800.	T.	H.&R.S.	C. S.
VI. *Prickly. Spikes globular.*						
pudica.	B. Lajuk.3	India.	Indigenous.	♃	R. S.	C. S.
Cavalum.		Bengal.	Dr. Buchanan, 1812.	T.		C. S.
mutabilis, R.		Bengal.	Mr. Turnbull, 1803.	♄	R. S.	C. S.
octandra, R.	B. Koochee-kanta.4	India.	Indigenous.	♄	R. S.	C. S.
Intsia.5	T. Korinta.	Coromandel.	H. C. 1801.	Sc. ♄	R. S.	C. S.
concina.	B. Reetha.6	India.	Indigenous.	Sc. T.	R. S.	C. S.
cæsia.	T. Konda-korinta.	Coromandel.	Dr. Berry, 1802.	Sc. ♄	R. S.	C. S.
pennata.	H. Pahari-aiel.	India.	Col. Hardwicke, 1801.	Sc. ♄	R. S.	C. S.
torta.		Coromandel.	H. C. 1803.	Sc. ♄	R. S.	H. S.
tenuifolia.	Tam. Kodi-vallen.	Coromandel.	H. C. 1803.	Sc. ♄	R. S.	C. S.
VII. *From America & Africa.*						
virgata.		W. Indies.	W. Hamilton, Esq. 1797.	Sc. ♄ ♂	R. S.	C. S.
glauca.		America.	W. Hamilton, Esq. 1797.	S. T.	R. S.	C. S.
herbacea.		America.	W Hamilton, Esq. 1799.	H.	R. S.	C. S.
asperata.		Vera Cruz.	W. Hamilton, Esq. 1798.	♄	R. S.	C. S.
illinoensis.		America.	W. Hamilton, Esq.		R. S.	C. S.
capensis.		Cape of G. H.	1799.	S. T.		
miamensis.		America.	W. Hamilton, Esq.	H.	R. S.	
grandiflora.			W. Hamilton, Esq. 1805.	♄ E.	R.&C.S.	
brachyloba.		N. America.	W. Hamilton, Esq. 1804.	♄ E.	R. S.	C. S.
tortuosa.			Hon. C. Greville, 1807.			
altissima.		Trinidad.	Mr. W. Lockhead, 1808.	Sc. ♄	9---10	
sensitiva.		Brazil.	Mr. W. Lockhead, 1808.	S. ♄		
lævigata,7 R.		Mauritius.	Col. Hardwicke, 1811.	T.		
CAPPARIS.						
spinosa.		Europe.	Hon. C. Greville, 1808.	♄		
acuminata.	H. Kalu-kera.8	India.	Indigenous.	Sc. ♄	C. S.	R. S.
zeylanica.	T. Adonda.	India.		Sc. ♄	R. S.	
sepiaria.	B. Kanta-goor-kamace	9. India.		♄	H. S.	
aphylla, R.	H. Keril.	Hindoostan.	Col. Hardwicke, 1807.	S. ♄		
frondosa.		America.	W. Hamilton, Esq.			
heteroclita, R.	T. Putta-tiga.	Coromandel.	Nalla-mandu, 1803.	Sc. ♄	12---2	
trifoliata, R.	B. Buroon.10	India.	Indigenous.	T.	1---2	R. S.
linifolia, R.		Coromandel.	Dr. B. Heyne, 1808.	L. T.		
CHELIDONIUM.						
majus.		Europe.	J. Bentley, Esq. 1812.			
PAPAVER.						
somniferum, single.	B. & H. Post.11		Before 1794.	☉		
red single.12				☉		
dubium.						
Rhoeas.						
ARGEMONE.						
mexicana.	H.&B.Burashial kanta.13	W. Indies.	Before 1794.	☉ E.	D. S.	D. S.
BIXA.						
orellana.	H. Lut-kun.14	W. Indies.	Before 1793.	T.	R. S.	C. S.
ANNESLE, A.						
spinosa, R.	B. Mukhana.	Chittagong,	S. Harris, Esq. 1802.	☉ A.	R. S.	C. S.
NYMPHÆA.						
rubra, R.	B. Ructa kumbula. 15	India.	Indigenous.		H.&R.S.	R. S.
Lotus.	B. Shalook.16	India.			R.&C.S.	R. S.
versicolor, R.		Bengal.	Kurrum Khan, 1803.		H.&R.S.	R. S.
cyanea, R.	B. Soondhi.17	India.	Indigenous.		H.&R.S.	R. S.
esculenta, R.	B. Chota soondhi.18	India.			H.&R.S.	R. S.
stellata.	B. Chota shalook.19	India.			H.&R.S.	R. S.
ÆGLE.						
marmelos.	H. & B. Bel.20	India.	Indigenous.	T.	3	10
CALLOPHYLLUM.						
inophyllum.	H. Sultan champa.21	Asia.	Indigenous.	T.	O	O
angustifolia, R.	M. Poon.	Pullo-Pinang.	Mr. Smith.	T.		
lanceolarium, R.		Mauritius.	Monsieur Jannet, 1810.	T.		
Bintagor, R.22		Moluccas.	Monsieur Jannet, 1810.	T.		
MAMMEA.						
americana.		W. Indies.	Sir G. Young, 1801.	T.	H. S.	
MESUA.						
ferrea.	B. Nagkeshur.23	India.	Before 1794.	S. T.	H. S.	C. S.

1 S. Shumee. H. Saee-kanta. 2 S. Khadira. 3 S. Lujjaloo. 4 T. Walla-doora, or Poota-korinta. 5 H. M. 6. t. 4. 6 T. Chikai. 7 Acacia Willd. 8 T. Palliky. 9 T. Nella-upi. 10 S. Vuroona, also Tikta-shaka. T. Tella-woolymera. 11 White or opium. P. S. Chosa. P. Kooknar. A. Khus-kash. 12 also, Double white, red, &c. many varieties. 13 or Bherband. 14 Can. Kuppa-manhala. H. A. 2. t. 19. 15 S. Hulusa, Ructa Sindhuka. T. Yerra-kalwa. 16 S. Koomooda. T. Tella-kalwa. 17 S. Kuhlara, also Neelumboojunma and Indeevura. T. Neety-kalwa. 18 S. Koomooda, and Kiruva. T. Kokta. 19 H. M. 11. t. 27. 20 S. Vilva. T. Maredoo. Shreephula, the S. H. & B. name of a small variety. 21 Tam. & T. Poona. 22 H. A. 2. t. 71. 23 S. Naga-keshura.

VATERIA.						
indica.	Can. Pene.1	Malabar.	Samuel Dyer, Esq. 1809.	L. T.	H. S.	8
lanceolaria, R.		Chittagong.	Mr. J. R. 1810.	T.	4—5	7—8
HOPEA.						
odorata, R.		Chittagong.	G. Dowdeswell, Esq. 1796.	L. T.	4—5	R. S.
eglandulosa, R.		Tipperah.	S. Harris, Esq. 1797.	L. T.		
DIPTEROCARPUS.						
alatus, R.		India, Pegu.	Dr. W. Carey, 1809.	L. T.		
turbinatus.	B. Shweta-gurjun.	Chittagong.	Before 1794.	L. T.	3	5
costatus.	B. Telia-gurjun.2	Chittagong.	Mr. J. R. 1810.	L. T.	C. S.	4—5
incanus, R.	B. Gurjun.	Chtttagong.	Mr. J. R. 1810.	L. T.	11—12	4
SHOREA.						
robusta, R.	S. Sala.3	Morung.	1796.	L. T.	4—5	R. S.
Tumbuggaia, R.	Tam. Tumbugai.	Coromandel.	Dr. A. Berry.	T.		
GARCINIA.						
Mangostana.						
Cornea.						
Cowa, R.						
Kydiana, R.						
purpurea, R.						
pictoria, R.						
paniculata, R.						
pedunculata, R.						
Cambogia.						
Gutta.						
lanceæfolia, R.	H. Kurendra.	Silhet.	Mr. M. R. Smith, 1811.	S. T.	1—2	6—7
Boobicowa, R.	B. Bhoomi-kuwa.	Silhet.	Mr. M. R. Smith, 1811.	S. T.		7—8
XANTHOCHYMUS.						
pictorius, R.	B. Dephul.4	India.	Before 1794.	T.	R. S.	R. S.
dulcis, R.		Moluccas.	H.C. 1798.	T.	H. S.	R. S.
ovalifolius, R.	Cin. Ambul-Goorka.	Ceylon.	Gen. H. Macdowall, 1804.	T.	H. S.	R.
GREWIA.						
orientalis.	T. Peyar.5	Coromandel.	Mr. Heshusius, 1802.	L. ♄	H. S.	C. S.
oppositifolia.		Napaul.	Dr. F. Buchanan, 1802.	S. T.	H.&R.S.	C. S.
asiatica.	B. & H. Phulsha.	India.	Before 1794.	T.	C. S.	H. S.
carpinifolia.	T. Nullie.	Coromandel.	Mr. Heshusius, 1809.	♄	○	C. S.
pilosa, R.		Bengal.	Dr. W. Carey, 1800.	♄ E.	R. S.	C. S.
polygama, R.		Bengal.	Dr. W. Carey, 1800.	♄ E.	R. S.	C. S.
sepiaria, R.	B.&H.Panee-Syora.6	Bengal.	Indigenous.	♄ E.	R. S.	C. S.
sapida, R.		Bengal.	Dr. W. Carey, 1797.	♄ E.	2—3	5—6
sclerophylla, R.		Hindoostan.	Col. Hardwicke, 1804.	♄ E.	4—5	10—11
umbellata, R.		Sumatra.	Mr. W. R. 1803.	♄	4—5	
aspera, R.		Coromandel.	Mr. P. Heshusius.	♄ Sc.	H.&R.S.	C. S.
ulmifolia, R.	Ch. Ki-yong.	China.	Mr. W. Kerr, 1805.	T.	H.&R.S.	C. S.
ELEOCARPUS.						
aristatus, R.	B. Ramduliya.	Silhet.	Mr. M. R. Smith, 1812.	L. T.	4	8
serratus.	H. & B. Julpai.	Chittagong.	Before 1794.	T.	H. S.	R. S.
serrulatus, R.		Tinnevelly.	G. Young, Esq. 1808.	S. T.		
Ganitrus, R.	S. Roodraksha.7	India.	Hon. C. A. Bruce, 1809.	T.	2—3	11
rugosus, R.	B. Bun Julpai.	Chittagong.	Mr. J. R. 1810.	L. T.	2—3	6—7
robustus, R.	B. Dulpai.	Silhet.	Mr. M. R. Smith, 1811.	T.	6—7	9—10
lucidus, R.		Chittagong.	Mr. J. R. 1810.	T.		
lanceæfolius, R.	B. Shweta-pai.	Silhet.	Mr. M. R. Smith, 1812.	T.	6—7	9—10
CORCHORUS.						
olitorius, green var.*	B. Pat.	India.	Indigenous.	⊙ E.	R. S.	9—10
reddish var.	H. & B. Bun-Pat.			⊙ E.	R. S.	9—10
capsularis, reddish var.*	H. Ghinalita-Pat.	India.		⊙	R. S.	9—10
green var.	B. Koshta.8			⊙ E.	R. S.	C. S.
fuscus, R.		India.		⊙	R.&C.S.	C. S.
fascicularis.	H. Junglee-pat.	India.		⊙ E.	C. S.	R. S.
decemangularis.		Bengal.	Dr. F. Buchanan, 1812.	⊙ E.	8—9	10—11
trilocularis.		Bengal.		⊙ E.	8—9	10—11
BERRIA.						
Ammonilla.	Cin. Ammonilla.	Ceylon.	Dr. Berry, 1802.		5	8
THEA.						
Bohea.		China.		♄ E.	C. S.	
viridis.		China.		♄ E.		
OCHNA.	[champaka.9					
squarrosa.	S.&H.B.Kunuk-	India.	Ram-dhun Choudri, 1806.	S. T.	2—3	5—6
HUMEA, R.						
elata.	H. Mous-jat.	Chittagong.	Mr. J. R.	L. T.	4—5	9—10
ROYDSIA, R.						
suaveolens.	H. Madhubi-lata.	Silhet.	Mr. M. R. Smith, 1810.	L. Sc. ♄	3	8

1 Its resin East India Copal. 2 or wood oil tree. 3 also Ushwukurnuka. H. & B. Shal.
4 S. Tumala. 5 H. M. 5. t. 46. 6 or Panshiooli. 7 B. Roodrakhya. 8 Tittha-Pat.
9 T. Yerra-juvi.

* These two are extensively cultivated in Bengal during the rainy season for the fibres of their bark, there called Jute, of which coarse sackcloth for rice bags, cordage, twine, &c. are manufactured.

DIGYNIA.

PÆONIA.						
Moutan.	Ch. Mou-tan.1	China.	Mr. W. Kerr, 1812.			
herbacea.	Ch. Teheuak-yeuak.	China.	Mr. W. Kerr, 1812.			

TRIGYNIA.

DELPHINIUM.						
Ajacis.	Larkspur.2	Helvetia.				
NIGELLA.						
indica.	B. Kala-jeera.3	Hindoostan.	Before 1794.	⊙ E.	C. S.	C. S.
damascena.		Europe.	Colonel Garstin.	⊙ E.	C. S.	C. S.

POLYGYNIA.

NELUMBIUM.						
speciosum.	S. B. H. Pudma.4	India.	Indigenous.	A ☽	H.&R.S.	C. S.
white.	B. Shweta-pudma.5	India.	Indigenous.			
rosy.	B. Rukta-pudma.6					
crimson.	Ch. Hung-lin.	China.	Mr. W. Kerr, 1809.			
DILLENIA.						
augusta, R.		Garrow-hills.	Mr. M. R. Smith, 1810.	L. T.		
speciosa.	B. Chalta.7	India.	Indigenous.		7	2
pentagyna, R.	T. Rawadam.		Mr. J. R.	L. T.	4---5	12
pilosa, R.		Goalpara.	Mr. R. Kyd, 1810.	L. T.		
scabrella, R.		Chittagong.	Mr. J. R. 1810.	L. T.		
LIRIODENDRON.						
grandiflora, R.	B. Doodiya-champa.	Silhet.	Mr. M. R. Smith, 1810.	T.	H. S.	C. S.
Liliifera.	B. Auoa-champa.	Silhet.		T.		
MAGNOLIA.						
grandiflora.		N. America.	W. Hamilton, Esq. 1808.	T.		
glauca.		America.	W. Hamilton, Esq. 1809.			
obovata.	Ch. Hong-yu-lan.	China.	Before 1794.	♄ S.		
pumila.	Ch. Ye-hup.	China.	1794.	♄ E.	☊	
fuscata.	Ch. Hum-sen.	China.	1794.	♄ E.	3---4	
Yulan.	Ch. Yu-lan.	China.	Mr. W. Kerr, 1810.)	♄ E.		
MICHELIA.						
Champaca	S. Champaka. H.&B.	India.	Before 1794.	T.	H.&R.S.	C. S.
ANNONA.						
squamosa.	B. & H. Ata.8		Before 1794.	S. T.	H. S.	7---10
reticulata.	H. & B. Nona.9	India.	Before 1794.	T.	H. S.	7---11
muricata.	Sour-Sop.	W. Indies.	A. Roberts, Esq. 1811.	T.		
Surriffa, R.		Silhet.	Mr. M. R. Smith, 1794.	T.		2---4
UVARIA.						
undulata, R.		Chittagong.	Mr. J. R. 1811.	♄ E.		11
cerasoides, R.	T. Chetta-duduga.	Coromandel.	1803.	T.	H. S.	
suberosa, R.	H.& B. Burai-chauli.10	India.	Indigenous.	S. T.	☊	C. S.
odorata.	Mn. Kananga.	Sumatra.	Dr. C. Campbell, 1798.	T.	☊	C. S.
longifolia.	B. Debdaroo.11	India.	Before 1793.	T.	2---3	R. S.
villosa, R.		Bengal.	1796.	S. T.	3---4	6---7
ventricosa, R.		Tipperah.	S. Harris, Esq. 1797.	T.	3	7
tripetala, R.		Moluccas.	H. C. 1797.	S. T.	4---9	
grandiflora, R.		Sumatra.	Mr. W. R. 1804.	♄	☊	C. S.
elongata, R.		Chittagong.	1806.	L. Sc. ♄	R. S.	
odoratissima, R.	C. Ang-chou-tan.	China.	Before 1793.	L. Sc. ♄	☊	☊
macrophylla, R.	B. Bagh-runga.	Bengal.	Kurrum, 1806.	L. ♄	R. S.	
bracteata, R.	B. Jupa-bun-kula.	Silhet.	Mr. M. R. Smith, 1811.	♄ Sc.	4---5	8---9
Badajamba, R.	B. Bada-jamba.	Chittagong.	Mr. J. R. 1810.	T.		6---7
heteroclita, R.	B. Tubee-kura.	Silhet.	Mr. M. R. Smith, 1812.	♄ Sc.	6---7	10---11
UNONA.						
longiflora, R.	H. Kula-kura.	Silhet.	Mr. M. R. Smith, 1810.	S. T.	4---5	
dumosa, R.	B. Tubee-kura.	Silhet.	Mr. M. R. Smith, 1812.	Sc. ♄	4---5	9---10
ATRAGENE.						
zeylanica.	H. Chagool-batee.	India.	Indigenous.	♄ Sc.	R. S.	C. S.
CLEMATIS.						
gouriana, R.		Bengal.	H. Creighton, Esq. 1800.	L. Sc. ♄	9	1---2
THALICTRUM.						
bractiata, R.		Bengal.	Dr. W. Carey, 1797.	H. Sc.	C. S.	
RANUNCULUS.						
indicus, R.		Bengal.	Indigenous.	☊ E.	C. S.	C. S.
bulbosus.		England.	J. Bently, Esq. 1811.	H. E.	3---4	3---5

1 Three varieties. 2 Several varieties. 3 S. Musavi. 4 P. Neelufer. 5 Pudma. Ch. Young-gaw. T. Tella Tamara. 6 T. Yerra Tamara. 7 S. Ruvya. T. Uva. 8 Custard Apple. S. Ganda-gatra. 9 Luvunee. 10 T. Chelta-duduga. 11 S. Deva-daroo.

CLASSIS XIV.

DIDYNAMIA GYMNOSPERMIA.

AJUGA.						
repens, R.		Chittagong,	Mr. J. R. 1810.		1---2	2---3
disticha, R.	B. & H. Gobura.1	India.	Indigenous.	L. E.	R.&C.S.	C. S.
fruticosa, R.	Tam. Rette peme-[retti.2	Coromandel.	Rev. Mr. Gericke, 1799.	L. ♃ E.	R.&C.S.	C. S.
TEUCRIUM.						
canadense.		Canada.	W. Hamilton, Esq.			
stoloniferum, R.		Bengal.	Mr. Abraham, 1801.	H. E.	R. S.	C. S.
SATUREJA.						
hortensis.	Summer Savory.	Europe.		☉ E.	D. S.	H. S.
HYSSOPUS.						
officinalis.	Hysop.	Europe.	B. Boswell, Esq. 1796.			
nepetoides.		N. America.	W. Hamilton, Esq. 1799.			
LAVANDULA.						
Spica.		Europe.	Mr. M'Mahon, 1810.			
LAVANDULA.3						
MENTHA.						
verticillata, R.	H. & B. Panee-kula.	Bengal.	Indigenous.	☉ E.	R. S.	C. S.
secunda, R.	H. Beel-luta.			E.	C. S.	H. S.
sativa.	H. Podeena.		1795.	♃ Cr.		
imbricata.		Napaul.	Dr. F. Buchanan, 1802.		9	
quadrifolia, R.	T. Konda-jajari.	India.	Dr. F. Buchanan.	♃ E.	R.&C.S.	C. S.
perilloides.		Napaul.	Dr. F. Buchanan, 1802.	☉ E.	C. S.	C. S.
PERILLA.4						
GLECHOMA.						
erecta, R.	H. Najbo.	Skirts of Na-[paul.	Dr. F. Buchanan, 1802.	☉ E.	C. S.	C. S.
LAMIUM.						
amplexicaule.		Europe.	W. Hamilton, Esq. 1800.	☉		
purpureum.		Europe.	1807.			
STACHYS.						
alpina.		Europe.	H. C. 1806.			
BALLOTA.5						
MARRUBIUM.						
supinum.		Europe.	Accident, 1801.			
vulgare.		Europe.	Dr. Hornman, 1807.			
LEONURUS.						
tataricus.		India.	Before 1794.	☉ E.	♀	♀
PHLOMIS.						
nepetifolia.	H. Khujoor-churi.	India.	Indigenous.	☉ E.	C. S.	H. S.
zeylanica.	B. & H. Hulkusa.6			☉ E.	C. S.	H. S.
cephalota, K.	H. Bura hulkusa.			☉ E.	R.&C.S.	C. S.
biflora.	H. Shwet busunta.	Bengal.		♃ Cr.	R.&C.S.	C. S.
Leonurus.		Cape of G. H.	W. Hamilton, Esq. 1809.	♄ E.	♀	♀
esculenta, R.	B. Chota hulkusa.	India.	Indigenous.	☉ E.	R.&C.S.	C. S.
pilosa, R.		Bengal.		♃ E.	R. S.	C. S.
CLINOPODIUM.						
incanun.		America.	W. Hamilton, Esq.			
repens.		Napaul.	Dr. F. Buchanan, 1802.	♃ Cr.	♀	♀
angustifolium.			W. Hamilton, Esq.			
ORIGANUM.						
Majorana.	Sweet marjoram.	Asia.	B. Boswell, Esq. 1796.	♂ E.	C. S.	H. S.
THYMUS.						
vulgaris.	Garden Thyme.	Europe.	Before 1794.	♄ D.		
Acinos.		Europe.	W. Hamilton, Esq. 1800.	☉		
MELISSA.						
officinalis.		Europe.	1799.	♂ D.	C. S.	
DRACOCEPHALUM.						
Moldavica blue flowered.		Europe.	1796.	☉ E.	C. S.	C. S.
pink.		Persia.	Captain Malcolm.	☉ E.	C. S.	C. S.
canariensis.		Canary Islands.	Lady Ann Barnard, 1799.			
denticulata.		Carolina.				
OCYMUM.						
villosum, R.	H. & B. Toolusee.7	India.	Indigenous.	♃ E.	♀	♀
sanctum.	H. & B. Kala Toolsee.8			♃ E.	♀	♀
album.		Uncertain.	W. Hamilton, Esq.	☉ E.	R.&C.S.	C. S.
polystachion.	T. Neeroo-toolusee.	Circars.	Nalla-munda.	☉ E.	R. S.	C. S.

1 T. Mabera. Ballota disticha. 2 Nepeta malabarica. 3 See Plectranthus.
4 See Mentha. 5 See Ajuga. 6 T. Poo-alla-toomi. 8 S. Urjaka. H. M. 10. t. 87.
9 S. Purnasa. T. Krishna-toolusee.

caryophyllatum, R.	B.&H.*Goolal-toolshee*.	India.	Dr. Rottler, 1807.	♃ E.	R.&C.S.	C.S.
pilosum, R.	H. *Babooi-toolshee*.1		Indigenous.	♃ E.	○	○
gratissimum.	H.&B.Ram-toolshee.	India.	Gardeners.	♄ E.	○	○
thyrsiflorum.			Dr. Rottler, 1807.	♂ E.	☾	○
cristatum, K.		Coromandel.	H. Russell, Esq. 1804.	☉ E.	R. S.	C. S.
inodorum.		Circars.	Nalla-munda, 1804.	♃ or ♂ E.	R.&C.S.	C.S.
Basilicum.	P. Deban-shab.2	Persia.	General Malcolm, 1808.	☉ E.	R.&C.S.	C.S.
PLECTRANTHUS.						
secundus, R.		Mysore.	Mr. B. Heyne, 1800.	☉ E.	10---11	1---2
scutellaroides.3		Moluccas.	H. C. 1798.	♃ E.	3---4	4---5
monodelphia. R.		Mysore.	Mr. B. Heyne.	☉ D.	C. S.	H. S.
aromaticus, R.	B. Pathur-choor.4	India.	Before 1794.	♓ E.	R. S.	
strobiliferus, R.5			Dr. Buchanan.	☉ E.	C. S.	R. S.
SCUTELLARIA.						
indica.6		Moluccas.	Accident.	☉ E.	C. S.	C. S.
perigrina.		Napaul.	Dr. F. Buchanan, 1801.	☉ E.	C. S.	H. S.
COLEBROOKIA.						
ternifolia.		Mysore.	Dr. F Buchanan, 1801.	L. E. ♄	2---3	H. S.
oppositifolia.		Napaul.	Dr. F. Buchanan.	L. E. ♄	2---3	H. S.
PRASIUM.						
melissifolium.	B. Hursmun.	Silhet.	Mr. M. R. Smith, 1810.	♄ Sc.	R. S.	C. S.

ANGIOSPERMIA.

BARTSIA.						
coccinea.		N. America.	Mr. M'Mahon, 1809.	♃ D.		
ANTIRRHINUM.						
triphyllum.			From the Cape.	☉ E.	C. S.	
junceum.		Europe.		☉ E.	C. S.	
delawariana.		America.	W. Hamilton, Esq.	D.		
Cymbalaria.			W. Hamilton, Esq.	☉ D.		
Linaria.		Europe.	W. Hamilton, Esq.	♃		
rosmarinifolia.7						
Orontium.		Europe.	Dr. P. Russell.	☉		
purpureum.			Sir A Hume, 1811.	☉	C. S.	C. S.
triste.						
speciosum.			Sir A. Hume, 1812.			
MARTYNIA.						
diandra.		Mexico.	W. Hamilton, Esq. 1802.	☉ E.	R. S.	C. S.
TORENA.						
diffusa, R.		Coromandel.	Accident.	☉ D.	R. S.	R. S.
multiflora, R.		Bengal.	Indigenous.	☉ E.	R. S.	R. S.
varians, R.				☉ E.	R. S.	R. S.
hians, R.			Accident.	☉ D.	R. S.	R. S.
CELSIA.						
coromandeliana.	B. Kookshima.	India.	Indigenous.	☉ E.	C. S.	H. S.
cretica.			Sir A. Hume, 1812.			
DIGITALIS.						
stricta, R.		Bengal.	Indigenous.	☉ E.	R. S.	C. S.
CRESCENTIA.						
Cujete.	Calabash tree.	S. America.	W. Hamilton, Esq. 1795.	S. T.		
BRUNFELSIA.						
undulata.		W. Indies.	W. Hamilton, Esq. 1800.	♄ E.	6	
BROWALLIA.						
elata.		Peru.	W. Hamilton, Esq. 1800.	☉ E.	○	○
OROBANCHE.						
indica.8		Bengal.	Indigenous.	H. E.	R. S.	
acaulis, R.9				H. E.	R. S.	
pedunculata, R. 10				H. E.	R. S.	
THUNBERGIA.						
fragrans, R.	T. Yeanadari-tiga.	Coromandel.	1794.	♄ T.	○	○
grandiflora, R.		Bengal.	Indigenous.	♄ T.	○	○
ACANTHUS.						
ilicifolius.	H. Harkuch-kanta.11	India.	Indigenous.	♄ E.	☾	C. S.
madraspatensis.		Coromandel.	Nalla-mandu, 1803.	H. D.	C. S.	H. S.
BARLERIA.						
Prionitis	B. Kanta jatee.12	India.	Indigenous.	♄ E.	C. S.	C. S.
spicata, R.13		Bengal.	Jandi Mali, 1804.	♄ E.	C. S.	
buxifolia.14		Mysore.	Dr. F. Buchanan, 1801.	♄ D.	C. S.	
cristata, purple.	S.B. Jhintee.	India.	Before 1794.	♄ E.	10---11	2---3
rosy.	H. Pia-bausa.	Cawnpore.	Colonel Hardwicke.	♄ D.	10---11	2---3
ciliata, R.	B. *Oojootee*.15	Bengal.	Dr. W. Carey, 1799.	♄ D.	C. S.	H. S.
dichotoma, R.	B. Sada jatee.		Kurrum-khan.	♄ E.	C. S.	H. S.
cœrulia, R.	S. Dasee.16	India.	Before 1794.	♄ E.	C. S.	H. S.

1 S. *Barbura*. P. Tokma-rihan. H. Rehan, also Tookmareea. 2 And Deban Macawassi. 3 S. Pashana-bhedee. 4 H. A. 5. t. 101. 5 H. M. 10 t. 90. 6 H. A. 5. t. 170. 7 Two varieties, white and rosy. 8 On the root of tobacco. 9 On Sugar-cane root. 10 On the root of Andropogon muricatum. 11 S. Harikussa. 12 S. Kuruntuka. 13 A handsome variety of prionitis. 14 H. M. 2. t. 47. 15 S. Kuruvuka. 16 Also Vana.

RUELLIA.						
infundibliformis, R.1		India.	Dr. F. Buchanan.	♄ E.	☋	C. S.
zeylanica.2			Dr. F. Buchanan.	♄ Sc.	☋	C. S.
comosa, R.		Moluccas.	H. C. 1798.	♄ D.	R. S.	C. S.
repens.			Klein and Rottler, 1780.			
cernua, R.3		Mysore.	Dr. F. Buchanan, 1800.	♄ E.	R. S.	C. S.
hirta.		Circars.	Nalla-mandu, 1803.	H. Cr.	C. S.	H S.
latebrosa.			Nalla-mandu, 1803.	H. Cr.	C. S.	H S.
imbricata.		India.	Indigenous.	☿ D.	R.&C.S.	C. S.
dependens, R.		Mysore.	Dr. Buchanan, 1801.	☿ E.	R. S.	C. S.
salicifolia, R.		Circars.	Nalla-mandu, 1803.	☉ E.	R. S.	C. S.
longifolia, R.	B. Kanta-koolika.4	India.	Indigenous.	H. E.	R. S.	C. S.
obovata, R.		Chittagong.	Mr. J. R. 1810.	☉ D.	9---11	2---3
ringens.5		India.	Nalla-mandu.	☿ Cr.	☋	C. S.
fasciculata.		Circars.	Nalla-mandu, 1803.	☿ D.	D. S.	H. S.
trifiora.		Bengal.	Indigenous.	☉ D.		
LEPIDAGATHIS.						
cristata.		Coromandel.	H. Russell, Esq. 1803.	☿ D.		
ROSCOEA.						
pentandra, R.	B. Kungeea.	Silhet.	Mr. M. R. Smith, 1811.	♄ Sc.	3---4	R. S.
CLERODENDRUM.						
inerme.	B. Bun-joma.	India.	1799.	S. S.		
phlomoides.6	T. Teloki.		Mr. Gibbons, 1798.	♄ E.	R. S.	
VOLKAMERIA.						
infortunata, R	B. Bhant.7	India.	Indigenous.	♄ E.	2---3	4---5
Kæmpferi.		China.	Before 1793.	♄ E.		
urticifolia, R.		Pegu.	Mr. F. Carey, 1809.	♄ E.	8---11	
dentata, R.		Silhet.	Mr. M. R. Smith, 1810.	♄ E.	5---8	
Buchananii, R.		Chittagong.	Dr. F. Buchanan, 1797.	♄ E.	R. S.	
serrata.		Mysore.	Dr. F. Buchanan.	♄ E.	5---6	9---11
herbacea, R.		Napaul.	Dr. F. Buchanan, 1802.	H. E.	5---6	10---11
nereifolia, R.		Chittagong.	Mr. J. R. 1810.	♄ E.	2	
odorata.	B. Bhantee.	Napaul.	Dr. F. Buchanan, 1802.	L. ♄	1---2	
Madœera.	H. Madœer.	Cawnpore.	Col. Hardwicke, 1800.			
farinosa, R.		Silhet.	Mr. M. R. Smith, 1811.	L. ♂	R. S.	
HASTINGIA.						
coccinea.	B. Ungootee.8	Bengal.	Before 1793.	L. ♄	C. S.	H. S.
SIPHONANTHUS.						
indica.	B. Bamunhatee.9	Bengal.	Before 1793.		4-5 & 8-12	12--1
hastata,R.	H. Hatti-kana.	Silhet.	Mr. M. R. Smith, 1811.	♄ E.	4---5	6---7
VITEX.						
Haynga.	H. Haynga.	Chittagong.		T.		
trifoliata.10		Pegu.	Mr. F. Carey, 1810.	S. T.	5---6	
Negundo.	H. Nisinda.11	India,	Indigenous.	S. T.	☋	C. S
altissima.		Goalpara.	S. Harris, Esq. 1797.	L. T.	4---5	8---9
alata, R.	B. Usoola.	Chittagong	Dr. F. Buchanan, 1798.	T.	4---5	8---9
incisa.		China.	Mr. W. Kerr, 1803.	L. ♄	☋	C. S.
arborea, R.	T. Boosee.	Circars.	1799.		4	8
Leucoxylon.		Ceylon.	Dr. Berry, 1800.	T.	4	8
saligna, R.		Coromandel.	Dr. B. Heyne, 1808.	T.	4	
heterophylla, R		Tipperah.	S. Harris. Esq. 1797.	T.		
PREMNA.						
esculenta, R.		Chittagong.	Mr. J. R. 1810.	♄ E.	☋	☋
latifolia, R.	T. Padda-nella kura.	Coromandel.	1800	S. T.	4	8
serratifolia,	B. Bhoot bhiruvee.		Gardeners.	S. T.	R. S.	C. S.
spinosa, R.	B. Guniaree.12	Bengal.	Indigenous.	S. T.	R. S.	C. S.
integrifolia.13		India.		♄	H. S.	R. S.
longifolia, R.		Bengal.	Indigenous.	S. T.	H.&R.S.	C. S.
herbacea, R.	B. Bhooi-Jamba.14		Dr. W. Carey, 1801.		2---3	
GMELINA.						
asiatica.	T. Goomoodoo.	Coromandel.	Dr. Berry, 1799.	L. ♄	☋	C. S.
villosa, R.		P. of W. Isl.	Dr. W. Hunter.	S. T.	☋	C. S.
arborea, R.	H. & B. Gumhar.15	India.	1795.	L. T.	H. S.	C. S.
parvifolia, R.	Tam. Goomadi.16	Coromandel.		♄	C. S.	
AVICENIA.						
tomentosa.	B. & H. Bina.	India.	Mr. Goodlad.	S. T.	H. S.	
LANTANA.						
indica, R.		Mysore.	Dr. B. Heyne.	♄	☋	
aculeata.		America.	W. Hamilton, Esq. 1809.	L. ♄	☋	☋
trifolia.			Dr. W. Carey, 1807.		☋	☋
STREPTIUM.						
asperum, R.	T. Ubeera.	Coromandel.	Nalla-mandu, 1803.	☿ E.	C. S.	H. S.

1 Justicia Linn. H. M. 9. t. 62. 2 Two varieties, purple and white. H. M. 9. t. 45. 3 H. A. 6. t. 23. f. 1. 4 S. Gokantuka. T. Neer-goobi. 5 H. M. 9. t. 64. 6 Two varieties. 7 Clerodendrum, L. S. Bhantee. 8 Also Doodmari. 9 S. Brahmunayustika. 10 H. A. 4. t. 18. H. M. 2. t. 10. 11 S. Sindhooka. T. Wayalakoo. 12 S. Gunikarika. H. Ustabunda. 13 H. A. 3. t. 193. 14 S. Bhoomi-jumbooka. 15 S. Gumbharee. H. M. 1. t. 41. 16 Can. Koomatha.

LIMOSELLA.						
diandra.		India.	Indigenous.	A.		
ERINUS.						
bilabiatus, R.		Moluccas,&c.	H. C. 1798.	⊙ E.	C. S.	C. S.
CAPRARIA.						
gratissima, R.	H. Kala-kurpoor.	Coromandel.	1797.	⊙ E.	C. S.	H. S.
diffusa, R.		Pegu.	Mr. F. Carey, 1808.	♂ D.	C. S.	H. S.
monnieria.1						
STEMODIA.						
ruderalis.	B. Huldi-busunta.	Bengal.	Indigenous.	⊙ D.	R. S.	C. S.
viscosa, R.	T. Boda-sarum.	Coromandel.	Mr. Heshusius, 1802.	⊙ E.	C. S.	C. S.
SESAMUM.						
orientale.	B. & H. Til.2	India.	Gardeners.	⊙ E.	H. S.	H. S.
indicum.	H. Krishna til.		Gardeners.	⊙ E.	R. S	R. S.
a downy variety.		Napaul.	Dr. F. Buchanan, 1802.	⊙ E.		
COLUMNEA.						
balsamica, R.	B. Kurpoor.3	India.	Gardeners.	⊙ E.	R.&C.S.	C. S.
heterophylla, R.		Bengal.	Indigenous.	⊙	C. S.	H. S.
tomentosa, R.	B. Chota bhooi kamadi.			⊙ D.	C. S.	H. S.
minuta, R.		India.		⊙ D.	R.&C.S.	C. S.
INCARVILLEA.						
parasitica, R.		Silhet.	Mr. M. R. Smith, 1811.	♄ D.	R. S.	9--10
BIGNONIA.						
stipulata, R.		Pegu.	Mr. F. Carey, 1810.	T.	H. S.	
undulata, R.		Cawnpore.	Capt. Hardwicke, 1802.	S. T.	2---3	7---8
catalpa.		America.	W. Hamilton, Esq.	S. T.		
spathacea.	F. Woodi.	Coromandel.	Dr. A. Berry.	S. T.	H. S.	C. S.
suaveolens, R.	B. Parool.4	Hurdwar.	Col. A. Kyd, 1799.	T.	H. S.	C. S.
crispa.		Mysore.	Dr. Buchanan.	T.	6	12
Chelonoides.	T. Tagada.5	India.	Mr. Snodgrass, 1800.	L. T.	6	12
xylocarpa, R.		Malabar.	Dr. Berry, 1804.	T.	3---4	2---3
indica.	H. Shona.6	India.	Indigenous.	T.	6	C. S.
suberosa, R.		Tanjore.	Dr. J. Anderson, 1794.	S. T.	C. S.	R. S.
grandiflora.	C. Tung-wang-fa.	China.	Before 1794.	♄ Cr.	6---8	C. S.
radicans.		America.	H. C. 1807.	♄ Cr.		
crucigera.			W. Hamilton, Esq.	♄ Cr.		
PEDALIUM.						
murex.	Tam. Amei-nerunshil.	Madras.	Dr. A. Berry, 1811.	⊙ D.		O

1 See Gratiola. 2 S. Tila. P. Roghen. A. Duhn. 3 H. M. 9. t. 85, & 12. t. 36.
4 S. Patuli. H. Purula. 5 Tam. Poupadyræ. 6 S. Shyonaka. Mundooka-purna. T. Pam pena. Tam. Vaag-marum.

CLASSIS XV.

TETRADYNAMIA SILICULOSA.

MYAGRUM.						
hispanicum.		Spain.	M. De Cossigny, 1802.		3	
sativum.		Europe.	Dr. Hornmann, 1807.	⊙ E.	2	
perfoliatum.		France.	A. B. Lambert, Esq. 1803.		3---4	
BUNIAS.						
Erucago.		Europe.	A. B. Lambert, Esq. 1802.		1---2	
LEPIDIUM.						
Thlaspi.						
sativum.	B. Halim.1	Asia.	Before 1794.	⊙ E.	C. S.	C. S.
bonariense.			Accident.			
THLASPI.						
Bursa pastoris.	Shepherd's purse.	Europe.	Accident.	⊙	C. S.	H. S.
campestre.		Europe.	Accident, 1812.	⊙ E.	C. S.	H. S.
COCHLEARIA.						
flava.		Bengal.	Dr. F. Buchanan, 1812.			
Armoracia.	Horse-raddish.	Europe.	1798.	H.		
Coronopus.		Europe.	Accident, 1812.	⊙ D.	C. S.	H. S.
ALYSIUM.						
deltoidum.		Europe.				

SILIQUOSA.

SISYMBRIUM.						
nasturtium.	Water-cress.	Europe.	Captain Simson, 1795.			
ERYSIMUM.						
Barbarea.	Winter-cress.2	Europe.	1809.			
CHEIRANTHUS.						
Cheiri.	H. Lahoree-shuboo.3	Britain.	General Martin.			
mutabilis.		Madeira.	Mr. J. R.			
incanus.4		Europe.				
maritimus.						
BRASSICA.						
Rapa.	B. Shalgum.5				D. S.	
oleracea.	H. & B. Kopee. 6				D. S.	
erucoides, R.	B. Shwet *Surisha*.7	Bengal.	Accident.		D. S.	
Eruca.		Europe.	Mr. Forsyth, 1803---4.		D. S.	
SINAPIS.						
nigra.	Black mustard.	Britain.	Colonel Garstin.	⊙ E.	D. S.	D. S.
alba.	White mustard.		Mr. P. Bowil, 1802.	⊙ E.	D. S.	D. S.
arvensis.		Europe.	Dr. Hornmann, 1807.	⊙ E.	D. S.	D. S.
dichotoma, R.	B. *Surisha*.8	India.	Indigenous.	⊙ E.	D. S.	D. S.
glauca, R.	B. Shwet raee.			⊙ E.	D. S.	D. S.
ramosa, R.	B. *Jooni* raee.9	Bengal.		⊙ E.	D. S.	D. S.
brassicata.	Ch. yea-tsai.10	China.	Mr. J. Miller.	⊙ E.	D. S.	D. S.
trilocularis, R.		Napaul.	Dr. Buchanan, 1802.	⊙ E.	D. S.	D. S.
cuneifolia, R.		Tibet.	Dr. Buchanan, 1802.	⊙ E.	D. S.	D. S.
rugosa, R.			Dr. Buchanan, 1802.	⊙ E.	D. S.	D. S.
erysimoides, R.		Malabar.	Captain Dickenson.	⊙ E.	D. S.	D. S.
divaricata, R.	B. Bun-raee.	Bengal.	Indigenous.	⊙ E.	D. S.	D. S.
patens, R.	H. Beel raee.			⊙ E.	D. S.	D. S.
pusilla, R.		Coromandel.	Dr. Berry.	⊙ E.	D. S.	D. S.
prostrata, R.		China.	Mr. W. Kerr, 1812.	⊙ D.	C. S.	C. S.
RAPHANUS.						
sativus.	H. & B. *Moola*.11	India.				
CLEOME.						
monophylla.		Coromandel.	1807.	⊙	R. & C. S.	C. S.
pentaphylla.	B. & H. Karela.12	India.	Indigenous.	⊙	○	○
Chelidonii.	T. Kar-wominta.		H. T. Colebrooke, Esq.	⊙	R. S.	R. S.
viscosa.	H. & B. *Hoor-hoorya*.		Indigenous.	⊙	○	○
heptaphylla.13		W. Indies.	W. Hamilton, Esq.	⊙	○	○

1 Garden cress two varieties. P. Tereezách. 2 Also French or American cress. 3 Wall-flower. 4 Purple or violet stock-gilly flower. 5 P. Shellam. 6 P. Ghellum. A. Kelum. 7 S. *Siddhartha*. 8 S. *Surshupa*. B. Shanchi *sursha*. H. Surson. 9 S *Rajika*. Several varieties. 10 White China Cabbage. 11 S. Moolaka. P. *Toorb*. A. *Fujl*. 12 S. *Kara*-vela. G. Wominta. 13 Does not seem to differ from the next.

CLASSIS XVI.

MONADELPHIA TRIANDRIA.

TAMARINDUS.1						
FERRARIA.						
undulata.		Cape of G. H. 1798.			3	

PENTANDRIA.

WALTHERIA.						
indica.		India.	Indigenous.	♃ E.		
elliptica.		Bengal.	Dr. F. Buchanan, 1812.			
HERMANNIA.						
alnifolia.		Cape of G. H.	1799.	P.		
scordifolia.			1809.	♄	○	○
MELOCHIA.						
corchorifolia.	B. Tiki-opra.2	India.	Indigenous.	♃	R.&C.S.	C. S.
OCHROMA.						
Lagopus.		W. Indies.	Sir G. Young, 1800.	T.	4	C. S.
PASSIFLORA.						
laurifolia.	Water lemon.	W. Indies.	H. C. 1795.	♄ Sc.	R. S.	C. S.
cœrulea.		Brazil.	1799.	♄ Sc.	R. S.	
minima.		Curassao.	W. Hamilton, Esq. 1800.	♄ Sc.	○	○
suberosa.		W. Indies.	H. C. 1805.	♄ Sc.	R. S.	
lutea.		Jamaica.	Loddiges & Son, 1807.	♄ Sc.		
MODECCA.						
trilobata, R.	H. Akanda-phul.	Chittagong.	Mr. J. R. 1810.	♄ Sc.	R. S.	9---10
tuberosa, R.3		Malabar.	Mr. M. Brown, 1808.	♄ Sc.	R. S.	10
dubia, R.		Sumatra.	Uncertain, 1804.	♄ Sc.	R.	
ERODIUM.						
malacoides.		Europe.	A. B. Lambert, Esq. 1802.	☉		
Gruinum.		Africa, &c.	A. B. Lambert, Esq. 1802.	☉	C. S.	
cicutarium.		Europe.	Dr. Hornmann, 1807.	☉ D.	2	
ÆGICERAS.						
majus.4	B. Hulsi.	India.	— Goodlad, Esq. 1797.	S. T.	H. S.	
PISTIA.						
Stratiotes.	H. Toka-pana.5	Bengal.	Indigenous.	☉ A.	H.&R.S.	R.&C.S.

HEXANDRIA.

CANARIUM.						
bengalense, R.		Silhet.	Mr. M. R. Smith, 1811.	L. T.	5--6	1--2
commune.		Moluccas.	H. C. 1798.	L. T.		
sylvestre.			H. C. 1798.	T.		
nigrum, R.			H. C. 1798.	T.	5	
strictum, R.		Tinnevelly.	Mr. G. Young, 1808.	T.		

HEPTANDRIA.

PELARGONIUM.	[ranium.					
inquinans.	Scarlet flowered Ge-	Cape of G. H.		♄	C. S.	H. S.
zonale.	[um.	Cape of G. H.	Sir A. Hume, 1812.	♄		
capitatum.	Rose-scented Gerani-	Cape of G. H.	T. Brooke, Esq. 1802.	♄ D.	C. S.	H. S.
vitifolium.	Balm-scented Gerani-	Cape of G. H.	Sir J. Royds, 1809.	♄	C. S.	H. S.
tomentosum.	[um.	Cape of G. H.	Mr. M'Mahon, 1812.	♄		

DECANDRIA.

CONNARUS.						
paniculatus, R.		Chittagong.	Mr. J. R. 1810.	L. T.	6	
nitidus, R.	H. Toog.	Silhet.	Mr. M. R. Smith, 1810.	T.	H. S.	9--10
pentandrus, R.						
HELICTERES.						
Isora.6	Can. Ky-walla.	India.	1795.	S. T.	R.&C.S.	D. S.
KLEINHOVIA.						
hospita.7		Moluccas.	H. C. 1798.	T.	○	C. S.

1 See Diadelphia. 2 H. Beel-paat. H. M. 9. t. 73. 3 Modecca. H. M. 8. t. 20.
4 H. M. 6. t. 36. 5 S. Kumbhika and Vari-purnee. B. Pana. 6 H. M. 6. t. 30. 7 H. A. 3. t. 119.

STERCULIA.						
Balanghas.1	China Chesnut.	China.	Before 1794.	T.	H. S.	R. S.
urens, R.	T. Kavali.2	India.	D. Turnbull, Esq. 1808.	T.	C. S.	H. S.
colorata, R.	T. Karaka.		R. Pierard, Esq. 1797.	T.	C. S.	H. S.
villosa, R.		Bengal.	S. Harris, Esq. 1797.	T.	C. S.	H. S.
alata, R.	B. *Buddha*-Narikella.3	Chittagong.	Before 1794.	L. T.	2---3	C. S.
guttata, R.4		Malabar.	Capt. Dickinson, 1802.	T.	C. S.	6
lanceæfolia, R.	B. *Ushlea*.	Silhet.	Mr. M. R. Smith, 1811.	T.	C. S.	H. S.
coccinea, R.			Mr. M. R. Smith, 1811.	T.	C. S.	H. S.
parviflora, R.		Tipperah.	S. Harris, Esq. 1797.	T.	2---4	
angustifolia, R.		Napaul.	Dr. F. Buchanan, 1802.	T.	3	
fetida.5		India.	Dr. A. Berry, 1802.	T.	H. S.	
populnifolia, R.		Coromandel.	Dr. B. Heyne, 1800.	T.		
HERITIERA.						
littoralis.6		Mauritius.	Monsieur Cere, 1810.	T.		
minor, R.	B. S. Soonduree.7	Bengal.	--- Goodlad, Esq. 1796.	T.	3---4	9---10
THEOBROMA.						
Cacao.	Chocolata nut tree.	S. America.	H. C. 1798.	S. T.	R. S.	H. S.
BUBROMA.						
Guazuma.	Bastard cedar.	Jamaica.	Dr. J. Anderson, 1790.	T.	H.&R.S.	C. S.
ABROMA.						
augusta.8	B. *Oolut*-kumbul.	India.	Before 1794.	♄ E.	H.&R.S.	C. S.
DOMBEYA.						
angulata.		Mauritius.	Monsieur Jannet, 1811.	♄ E.		
tomentosa.		Madagascar.	Monsieur Jannet, 1811.	♄ E.		
tileœfolia.		Mauritius.		♄ E.	10---11	
PENTAPETES.						
phœnicea.	B. Doopahariya.9	India.	Indigenous.	☉ E.	R. S.	C. S.
PTEROSPERMUM.						
suberifolia.	S. *Moochukoonda*.10	India.	Before 1794.	T.	C.&H.S.	C. S.
acerifolia.	B. *Kunuk*-champa.11		1793.	T.	3---5	C. S.
semisagittata.		Chittagong.	Dr. F. Buchanan, 1796.	T.	3---5	12---1
lanceofolium, R.		Silhet.	Mr. M. R. Smith, 1811.	T.	5---6	C. S.
canescens, R.		Ceylon.	Gen. H. M'Dowall, 1803.	T.	3---5	11---12
ADANSONIA.						
digitata.		Africa.	Before 1794.	T.	H. S.	C. S.
BOMBAX.						
pentandrum.	B. Shwet *Shimool*.12	India.	Indigenous.	T.	2---3	4---5
heptaphyllum.	B. *Rukta Shimool*.13	India.	Indigenous.	T.	2---3	4---5
Ceiba.		W. Indies.	Mr. W. Lockhead, 1808.	T.		
gossypinum.		India.		T.		
SIDA.						
I. *Capsules from* 5 *to* 10.						
angustifolia.		Bourbon.	Monsieur Cere, 1810.	♃ E.	○	○
microphylla.		Bengal.	Indigenous.	♃	R.&C.S.	C. S.
cuneifolia, R.	T. Mutto-apullacum.	Circars.	Nalla-mandu, 1802.	♃	R.&C.S.	C. S.
acuta.14	B. *Kureta*.	India.	Indigenous.	♃	R.&C.S.	C. S.
geniculata.		America.	W. Hamilton, Esq.	♃ E.		
humilis.	B. *Junka*.15	India.	Indigenous.	♃ D.	○	○
nudiflora.		S. America.	Dr. Wallich, 1809.	♃ E.	R.&C.S.	C. S.
periplocifolia.		Sumatra.	Dr. C. Campbell, 1802.	♄ E.	R.&C.S.	C. S.
Jamaicensis.		Jamaica.	Hon. C. Greville, 1807.		R.&C.S.	D. S.
alba.		Coromandel.	Nalla-mandu, 1802.	♃ E.	R.&C.S.	D. S.
polyandra, R.			Dr. A. Berry, 1805.	H. E.	R.&C.S.	C. S.
Napaea.		Uncertain.	Accident.	♄ E.	R.&C.S.	C. S.
lanceolata.		India.	Indigenous.	♃ E.	R.&C.S.	C. S.
retusa.		Bengal.		♃ D.	R.&C.S.	C. S.
truncata.		W. Indies.	W. Hamilton, Esq. 1802.	☉	R.&C.S.	C. S.
carpinifolia.		Madura.	W. Hamilton, Esq. 1800.	♃ E.	R.&C.S.	C. S.
rhomboidea, R.	B. Shwet-baryala.	India.	Indigenous.	♃ E.	R.&C.S.	C. S.
rhombifolia.	B. *Rukta* Baryala.			♃ E.	R.&C.S.	C. S.
cordifolia.	B. Baryala.16	Coromandel.	Dr. Rottler, 1803.	H. E.	R.&C.S.	C. S.
II. *With more than* 10 *Capsules.*						
crispa.		Coromandel.	Dr. Rottler.	♃ D.	R. S.	C. S.
mauritiana.		Mauritius.	Dr. Klein, 1803.	♃		
Abutilon.	Ch. King-ma.	Moluccas &	Accident.	☉ E.	R. S.	C. S.
hastata.		[China.	W. Hamilton, Esq. 1802.	☉	R. S.	C. S.
Dilleniana.		Mexico.	Governor of Manilla, 1810.	♃ D.	C. S.	C. S.
tomentosa, R.		India.	1796.	☉ E.	R.&C.S.	C. S.
asiatica.	B. Jhampi Petaree.17	India.	Indigenous.	H. E.	R.&C.S.	C. S.
populifolia.	T. Nugoo-benda.			H. E.	R.&H.S.	C. S.
indica.				H. E.	R.&C.S.	C S.
graveolens, R.	H. *Bura*-kungee.	Duab.	Col. Hardwicke, 1800.	H. E.	C. S.	H. S.
cristata.		Mexico.	W. Hamilton, Esq.	♃ D.	C. S.	H. S.

1 H. M. 1. t. 49. 2 H. *Kooli*. 3 In Silhet *Toola*. 4 H. M. 4. t. 61. 5 H. A. 4. t. 107. 6 Balanopteris Tothila. Gœrt. 7 B. Minor. Gœrt. 8 Its bark abounds in strong very white fibres, which make good rope, &c. 9 S. *Ruktuka*, or *Bundooka*. 10 Tam. Taddæ marum. 11 S. *Kurnikara*. H. *Kuniyar*. 12 T. Kadami. 13 S. Shalmuli. The pod containing the cotton is called *Pakura*. T. Boorgha. 14 H. M. 10. t. 53. 15 T. Nellabenda. 16 S. *Batyalaka*. 17 G. Tootra-benda.

MALACHRA.						
capitata.		W. Indies.	Monsieur Jannet, 1810.	☉ E.	O	O
ALTHÆA.						
officinalis.	Marsh mallow.	Europe.	W. Hamilton, Esq. 1801.	♃ E.		
rosea.	H. Gool-khiree.	China.	Before 1794.	☉ E.	C. S.	H. S.
MALVA.						
americana.		N. America.		☉		
rotundifolia.		Persia.	Major Malcolm.	☉ D.	C. S.	H. S.
verticillata.		China.	A. B. Lambert, Esq. 1802.	☉	2---3	
capensis.		Cape of G. H.	1799.			
hispanica.		Europe.	A. B. Lambert, Esq.	☉ E.		
crispa.						
caroliniana.		Carolina.	Dr. Jacquin, 1806.		D. S.	
LAVATERA.						
arborea.		Europe.				
KITAIBELIA.						
vitifolia.		Europe.	Mr. M'Mahon, 1810.			
URENA.						
lobata.	B. & H. Bun-okra.	India.	Indigenous.	♃ E.	R.&C.S.	C. S.
sinuata.	B. & H. Kungooya.1			♃ E.	R.&C.S.	C. S.
repanda.		Coromandel.	H. C. 1804.	♄ E.	9---11	12---2
GOSSYPIUM.						
obtusifolium, R.		Ceylon.	General M'Dowell.	♄ E.	O	O
arboreum.	Red flowered.	India.	Before 1794.	♄ E.	O	O
herbaceum.*	S. Karpasee.2	India.	Before 1794.	♃ E.	O	O
religiosum.	Nankeen, or Tawny	China.	Hon. C. Grivell, 1809.	♃ D.	O	O
vitifolium.	[Cotton.			L. ♄ E.	O	O
acuminatum, R.		Hindoostan.	--- Robertson, Esq. 1800.	L. ♄ E.	O	O
barbadense.	E. Bourbon Cotton.	America.	Dr. J. Anderson, 1790.	♃ E.	O	O
hirsutum.3			W. Hamilton, Esq.	♄ D.	O	O
mexicanum.		Mexico.	W. Hamilton, Esq. 1804.			
KYDIA.						
calycina, R.	H. Chouputeeya.4	India.	Before 1794.	T.	R.&C.S.	C. S.
HIBISCUS.						
I. *Leaves entire, or but slightly lobed, or angular.*						
populnea.	B. & H. Purush.5	India.	Before 1794.	T.	R. S.	C. S.
populneoides, R.	B.&H. Purush-pipool.6	India.	1797.	T.	R.&C.S.	C. S.
tiliaceus.7		Moluccas.	H. C. 1798.	T.	O	O
tortuosus, R.	S. & B. Bala.8	India.	Indigenous.	T.	H.&R.S.	C. S.
macrophyllus, R.	H. Kasya-pal.9	Chittagong.	Mr. J. R. 1810.	T.		
Rosa sinensis.	B. Juba.10	India&China.	Before 1794.	♄	O	
phœniceus.	S. Soora-yamuni.11	India.		T.	R.&C.S.	C. S.
syriacus.	B. Juba.12	Syria.	Before 1794.	♄ E.	O	
carolinus.		W. Indies.	W. Hamilton, Esq.			
rigidus.	T. Salla-burta.	Coromandel.	1808.	♄ E.	R. S.	R. S.
tubulosus.		India.	Dr. W. Carey, 1800.	☿ & ♂ E.	9---10	1---2
pruriens, R.			Mr. B. Heyne, 1802.	♂ E.	H.&R.S.	R. S.
Solandra.		Mysore.	Mr. B. Heyne.	☉ E.	R. S.	R. S.
Lampas.		Bengal.	Mr. W. R. 1801.	♄ E.	R.&C.S.	C. S.
collinus, R.	T. Konda-gong.	Coromandel.	1804.	S. T.	R.&C. S.	C. S.
vitifolius.	B. Bun-kupasi.13	India.	Indigenous.	☉	R.&C.S.	C. S.
mutabilis.	B. Thol-pudma.14	India&China.	Before 1794.	S. T.	R.&C.S.	C. S.
Abelmoschus.	H. Kala-Kustooree.	India.	Indigenous.	☉ E.	R.&C.S.	C. S.
palustris.		N. America.	W. Hamilton, Esq.	♃ E.		
cancellatus, R.		Napaul.	Dr. F. Buchanan, 1801.	☉ E.	R.&C.S.	C. S.
truncatus, R.		Ceylon.	Gen. M'Dowell, 1802.	☉ E.	R.&C.S.	
scandens, R.		Chittagong.	Mr. J. R. 1811.	♄ Sc.	4---5	
chinensis, R.		China.	Mr. W. Kerr, 1812.	☉ E.	R.&C.S.	C. S.
II. *Leaves deeply divided.*						
tricuspis.		Society Islds.	Lord Minto, 1809.	T.	C. S.	H. S.
militaris.		America.	Dr. W. Carey, 1807.			
hirtus.		Mysore.	Dr. F. Buchanan.	☉ E.	R.&C.S.	C. S.
furcatus, R.		Bengal.	Dr. W. Carey.	♃ E.	R. S.	C. S.
bifurcatus.		S. America.	Mr. W. Lockhead, 1808.	♃ E.	R. S.	C. S.
surattensis.		India.	Dr. Klien, 1803.	♃ Sc.	R.&C.S.	D. S.

1 T. Palier-mankena. 2 B. Kapasa. H. Rooi. A. Kootn. P. Poombeh or Pembeh. 3 Two or three varieties. 4 T. Pandiki. 5 T. Gangari. 6 S. Gurdubhanda. T. Mooni Gangari. 7 H. A. 2. t. 73. 8 H. Bola. H. M. 1. t. 30. 9 Ropes are made of the bark. 10 S. Juva. Of this we have three varieties, viz. single red, double red, and double pale yellow called Patkili. 11 Red B. & H. Lall Surjya-muni. T. Yerra-Salla-burta. White B. & H. Sufet Soorjya-muni. T. Tella Salla-burta. 12 Single white. H. Sada-juba. Single purple. H. Neel-juba. Double white, and double purple. 13 S. Bharudwajee. 14 Single and double, rosy and white. S. Pudma-charinee. Sthula-pudma.

* Of this there are several varieties, very generally cultivated over various parts of India, &c. one about Dacca, of which their fine muslins are made; another in Berar, and from thence the cotton is carried into the northern Circars, where the fine Madras long cloths are made of it. A third has been introduced into this garden from China.

strictes, R.1	H. Buri-harul.	Rajmahl.	Accident, 1793.	☉ E.	R. S.	C. S.
prostratus.		Coromandel.	1803.	☉ D.	R.&C.S.	C. S.
diversifolius.2		Africa.	1799.	♃	C. S.	H. S.
Sabdariffa.		W. Indies.	Dr. Rottler.	☉ E.	R. S.	C. S.
cannabinus.	H. & B. Mesta-pat.3	India.	Dr. W. Carey, 1801.	☉ E.	R. S.	C. S.
radiatus.		Uncertain.	Before 1794.	♂ D.	C. S.	C. S.
longifolius.	H. Ram-turi.4	India.	1794.	♂ E.	R. S.	C. S.
esculentus.5		W. Indies.	W. Hamilton, Esq. 1802.	☉ E.	R. S.	C. S.
speciosus.		N. America.	1802.	☉ E.	R. S.	C. S.
tetraphyllus, R.		Bengal.	Indigenous.	☉	R. S.	C. S.
pentaphyllus, R.		China.	Mr. Kerr.		R.&C.S	C. S.
pungens, R.		Napaul.	Dr. F. Buchanan, 1802.	♃ E.	R.&C.S.	C. S.
trionum.		Cape of G. H.	1798.	☉ E.	R.&C.S.	C. S.
ternatus.			1790.			
PAVONIA.						
odorata.	T. Chittle-bunda.	Coromandel.	Dr. Rottler, 1802.	☉ E.	H. S.	C. S.
ACHANIA.						
pilosa.		W. Indies.	W. Hamilton, Esq. 1810.		C. S.	H. S.
GORDONIA.						
integrifolia, R.	B. Bun-nukh.6	Silhet.	Mr. R. Kyd, 1810.	L. T.	3—4	12
CAMELLIA.						
japonica.7	C. Fa-cha-fa.	China.	Sir C. Blunt, 1796.	♄ E.	C. S.	
MESUA.8						
BARRINGTONIA.						
speciosa.		Malay Islds.	Dr. John, 1808.	T.		
acutangula, R.	B. Hijjul.9	India.	Indigenous.	T.	H. S.	R. S.
racemosa, R.			H. C. 1798.	T.	H. S.	R. S.
CAREYA.						
herbacea, R.	B. Bhooi-dalim.10	Bengal.	Dr. W. Carey, 1799.	H.	H. S.	R. S.
arborea, R.	T. Kumbi.11	India.	Col. Hardwicke, 1802.	T.	H. S.	R. S.
spherica, R.		Chittagong.	S. Harris, Esq. 1798.	T.	H. S.	R. S.

1 The bark of this more than any other of the order known to me, abound in strong white, fibres, fit for twine, cordage, &c. 2 Armatus. B. H. 3 Also Pulooa. T. Gong-kura. 4 B. Dhenroos, T. Binda. 5 Two varieties. 6 H. Makurjal. 7 Many varieties. 8 See Polyandria. 9 S. Ijjul. T. Kadami. 10 S. Bhoomi-darimba. 11 Tam. Poottannimarum.

CLASSIS XVII.

DIADELPHIA TRIANDRIA.

TAMARINDUS.						
indica.	S. *Umlika*.1	India.	Indigenous.	L. T.	5—6	1—2
FUMARIA.						
officinalis.		Europe.	1799.			
parviflora.	H. *Bun-shulpha*.2	Bengal.	Indigenous.	☉ D.	D. S.	D. S.
vescicaria.		Cape of G. H.				

OCTANDRIA.

SECURIDACA.						
paniculata, R.	B. *Furrinjur*.	Chittagong.	Mr. J. R. 1810.	Sc. ♄	R. S.	3—4
POLYGALA.						
arvensis.	B. Merradoo.	India.	Indigenous.	H. D.	R. S.	C. S.
telephi.				H. D.	R.&C.S.	C. S.

DECANDRIA.

DALBERGIA.						
ougeinensis, R.	S. *Tinisha*.3	Hindoostan.	Dr. W. Hunter.	L. T.	H. S.	9—10
latifolia, R.	B. Sweta shala.4	India.	J. Glass, Esq. 1808.	L. T.	H. S.	9—10
Sissoo, R.	H. & B. Sissoo.5	Bengal.	Before 1794.	L. T.	H. S.	C. S.
emarginata, R.	Andaman Sissoo.	Andaman.	Before 1793.	L. T.	R. S.	C. S.
frondosa, R.		Coromandel.	Genl. H. M'Dowell, 1801.	L. T.	5—6	8—9—10
paniculata, R.	T. Patseroo.6	Circars.	1802.	L. T.	H. S.	C. S.
zeylanica, R.	C. Belloo-labba.	Ceylon.	Genl. H.M'Dowell, 1801.	L. T.	5—6	8—9—10
robusta, R.	H. Krowee.	Silhet.	Mr. R. Kyd, 1810.	L. T.	H. S.	C. S.
marginata, R.	B. Makura-*gila*.	Silhet.	Mr. M. R. Smith, 1812.	♄ Sc.	R. S.	H. S.
scandens.	B. Noa-luta.7	India.	Indigenous.	♄ Tw.	R. S.	C. S.
tamarindifolia, R.	B. Tento*olee*.	Silhet.	Mr. M. R. Smith, 1811.	♄ Sc.	3—4	C. S.
stipulacea, R.		Silhet.	Mr. M. R. Smith, 1811.	♄ E.	3—4	11
rimosa, R.	B. *Koogroom*.	Silhet.	Mr. M. R. Smith, 1811.	♄ E.	3—4	11—12
reniformis, R.	B. *Kurisa*.	Silhet.	Mr. M. R. Smith, 1811.	T.	3—4	11—12
	[*duna*.8					
PTEROCARPUS.						
Santalinus.	H. & B. *Rukta* Chun-	Coromandel.	Dr. A. Berry.	T.	H. S.	
marsupium, R.	H. Peet Shal*a*.9	India.	Before 1794.	L. T.	R. S.	C. S.
dalbergioides, R.	Andaman red-wood.	Andamans.	Col. A. Kyd, 1795.	L. T.	6—7	11—12
indicus.		Malayana & China.	Sir J. Roydes, 1808.		4	10—11
Draco.		America.	Captain Young, 1812.	T.		
GALEDUPA.						
arborea, R.	B. & H. *Kurunja*.10	India.	Indigenous.	T.	H. S.	C. S.
uliginosa, R.	B. Panee-luta.	Bengal.		♄ Tw.	D. S.	D. S.
elliptica, R.		Malayana.	H. C.	♄ Tw.	3	8
macrocarpa, R.		P. of W. Isl.	Dr. W. Hunter.			
piscidia, R.	B. Chunkee.	Silhet.	Mr. M. R. Smith, 1811.	S. T.	6—7	C. S.
marginata, R.	H. *Ook*nee.	Silhet.	Mr. M. R. Smith, 1811.	♄ Sc.	4—5	9—10
bijuga, R.						
BUTEA.						
frondosa.	B. & H. Pulas.11	India.	Indigenous.	T.	2—3	5—6
superba, R.	T. Tiga-maduga.	Coromandel.	Captain A. Denton, 1797.	L. Sc. ♄	2—3	
parviflora, R.		Coromandel.	Captain A. Denton, 1797.	L. Sc. ♄	11—12	4—5
	[*dar*.12					
ERYTHRINA.						
indica.	H. & B. Pal*ita*-man-	India.	Indigenous.	T.	H. S.	H. S.
stricta, R.	Can. En-mouricou.	Malabar.	M. Brown, Esq. 1808.	T.	3	5
ovalifolia, R.	H. Hur*i*-kek*r*a.13	India.	Indigenous.	T.	C. S.	H. S.
suberosa, R.	T. Mooni maduga.	India.	Indigenous.	S. T.		
arborescens, R.		Napaul.	Dr. F. Buchanan.	♄ E.	9—10	
herbacea.		Carolina.	W. Hamilton, Esq. 1804.	H.	C. S.	H. S.
resupinata, R.		Cawnpore.	Col. Hardwicke, 1802.	H.	C. S.	H. S.
cafra.		S. Africa.	Col. Hardwicke, 1811.	T.		

1 *Tintiree*. H. & B. *Ambli*. B. Amlee, Tent*ool*. Of this, we have three varieties, viz. the common sort, which differs from that of the West Indies in the fruit being much longer, with from six to twelve seeds; a second with sweet pulp; and a third with pink coloured pulp. 2 B. Sholoopha. 3 Also Syandu*na*. 4 T. Veru-gudu-chawa. 5 S. *Shingshupa*. 6 Can. Hasur-gunni. 7 T. Scherata*lli*-badoo. 8 S. Koochundu*na*. 9 T. Yeangasha. 10 Kurun-*juka*. S. 11 S. Pulasha. *Kingshooka*, &c. T. Maduga. 12 Or Furrud. S. Mundara. T. Badata. 13 S. Par*ibhudra*.

ABRUS.						
precatorius, seed red.	B. Koonch.1	India.	Indigenous.	♄ Tw.	C. S.	H. S.
seed white.	B. Sweta koonch.2		1803.	♄ Tw.	C. S.	H. S.
seed black.3		Mysore.	Mr. B. Heyne, 1800.	♄ Tw.	C. S.	H. S.
SPARTIUM.						
scoparium.	Common Broom.	Europe.	B. Boswell, Esq. 1799.	♄ E.		
junceum.	Spanish Broom.		M. De Cossigny, 1802.	♄ E.		
ULEX.						
europaeus.	Common Furze.4	Britain.		♄		
AMORPHA.						
fruticosa.		America.	A. B. Lambert, Esq. 1803.	♄ E.		
cœrulea.			A. B. Lambert, Esq. 1803.	♄ E.	3	
CROTALARIA.						
I. *Leaves simple.*						
juncea.*	H. & B. Sun.5	India.	Before 1794.	☉ E.	R.&C.S.	C. S.
linifolia.		India.	Col. Hardwicke.	☿ E.	C. S.	C. S.
tenuifolia, R.		Coromandel.	Dr. Klein, 1809.	☿ E.	C. S.	H. S.
sagittalis.		W. Indies.	W. Hamilton, Esq.	☉		
tetragona, R.		Napaul.	Dr. F. Buchanan.	☿ E.	R.&C.S.	C. S.
stipulacea, R.		Mysore.	Mr. B. Heyne, 1801.	☿ E.		
parviflora.		N. America.	Mr. M'Mahon, 1809.	☉	2---3	H. S.
montana, R.		Circars.	1796.	☿ E.	R.&C.S.	C. S.
stricta, R.	T. Moonga.	India.	Dr. W. Carey, 1804.	☉ E.	C. S.	C. S.
fulva, R.		Mysore.	Dr. F. Buchanan, 1801.	☿ E.	C. S.	C. S.
pulcherrima, R.			Dr. F. Buchanan.	☿ E.	C. S.	H. S.
ramosissima, R.		Bengal.	Dr. W. Carey, 1804.	☿ E.	R.&C.S.	C. S.
chinensis.		Coromandel.	Dr. Rottler.	♄ E.	10---11	1---3
prostrata, R.	B. Bhooi-Sun.6	India.	Nalla-mandu.	☿ D.	C. S.	C. S.
hirsuta.			P. Hessusius, Esq. 1803.			
biflora.			Mr. B. Heyne, 1803.	H. D.	C. S.	C. S.
retusa.	H. Beel-jhunjhun.7	India.	Indigenous.	☉ or ♂	C. S.	H. S.
sericea.	B. Peeyooli.8	Bengal.		☉ E.	R.&C.S.	C. S.
verrucosa.	H. & B. Bun-Sun.9	India.	Indigenous.	☉ E.	C. S.	H. S.
semperflorens.		Sumatra.	W. Ewer, Esq.	☿ E.	11---12	1---3
bialata, R.		Silhet.	Mr. M. R. Smith, 1810.	☿ D.	R.&C.S.	C. S.
angulosa.10		Mauritius.	Col. Hardwicke, 1811.	☉ E.	○	C. S.
paniculata.		Coromandel.	Dr. Klein, 1804.	♄ E.	C. S.	H. S.
II. *Leaves compound.*						
laburnifolia.	B. Muna.11	India.	Dr. John, 1795.	♄ E.	○	○
elliptica, R.		China.	Mr. W. Kerr, 1805.	♄ D.	8---9	C. S.
cytisoides.		Napaul.	Dr. F. Buchanan, 1801.	♄ E.	C. S.	H. S.
trifoliastrum.	T. Naga-galli-gheetsa.	Circars.	Nalla-mandu, 1804.	♄ E.	C. S.	H. S.
argentea.		Cape of G. H.	Mr. J. R. 1804.		C. S.	
capensis.		Cape of G. H.		♄ E.	C. S.	
lanceolata, R.		W. Indies.	Mr. W. Lockhead, 1808.	♄ E.	C. S.	C. S.
bracteata, R.		Chittagong.	Mr. Moddern.	♄ E.	9---10	1---2
quinquefolia.	T. Neeru-galli-gheetsa.	India.	Dr. A. Berry, 1802.	☉ E.	C. S.	C.&H.S.
ARACHIS.						
hypogœa.	C. Fa-sang.12	China, &c.	Before 1794.	H. D.	R.&C.S.	D. S.
fruticosa.	T. Sakilli-kampa.13	Ceylon, &c.	Dr. John, 1800.	☿ E.	○	○
LUPINUS.						
varius.		Cape of G. H.				
CARPOPOGON.						
monospermum, R.	T. Enuga.	India.	Captain Dickinson.	♄ Tw.	R.&C.S.	C. S.
pruriens, R.	B. Alkooshi.14	India.	Indigenous.	☉ Tw.	C. S.	H. S.
atropurpureum, R.	T. Tilla-kodi.	Travancore.	Dr. A. Berry, 1806.	♄ Tw.	1---2	R. S.
niveum, R.	H. Khamach.15	Bengal.	Mrs. Honeycomb, 1796.	☉ Tw.	R. S.	C. S.
giganteum, R.16		Chittagong.	Mr. J. R. 1810.	♄ Tw.	R.&C.S.	D. S.
imbricatum, R.		Silhet.	Mr. M. R. Smith, 1810.	♄ Tw.	R. S.	C. S.
bracteatum, R.		Chittagong.	Mr. J. R. 1811.	♄ Tw.	C. S.	
PHASEOLUS.17						
I. *Twining.*						
vulgaris.			Before 1794.	☉ Tw.	C. S.	C.&H.S.
lunatus.	E. Lima bean.18	Uncertain.	1794.	♂ Tw.	C. S.	H. S.
alatus.	B. Bun-burbutee.19	Bengal.	Indigenous.	☿ Tw	C. S.	H. S.
sublobatus, R.	B. Gora-moog.	Hindoostan.	Dr. W. Hunter.	☉ Tw.	D. S.	D. S.
calcaratus, R.	B. Ranga-moog.	India.	Mr. B. Heyne, 1800.	♂ Tw.	C. S.	H. S.
dolichoides, R.		Chittagong.	Mr. Moddern, 1803.	☿ Tw.	R. S.	C. S.
minimus, R.	Ch. Cham-loe-to.	China.	Mr. Kerr, 1809.	☉ Tw.	D. S.	D. S.

1 S. Krishnula Goonja, and Ructika, the seed. 2 Ranga Gooncha, and Rutti, the seed.
Ch. Lin-tin. 3 And white flowers. 4 Or Whin. 5 S. Shuna. T. Shanamoo.
C. Henna. 6 T. Seri-galli-gheetsa. 7 T. Potu-galli-gheetsa. 8 S. Peeyooksha.
9 T. Alla-galli-gheetsa. 10 Does it differ from verrucosa? 11 T. Pedda-galli-gheetsa.
12 H. Moong-phullee. 13 Stylosanthes of Schreber. 14 Also H. & B. Kiwach. S. Atmagoopta Murkutee, &c. T. Doola-ghonda. 15 An excellent bean. 16 H. M. 8. t. 36.
17 Of these Phaseoli, not less than eleven species, viz. 1, 2, 5, 10, 11, 12, 13, 14, 15, 16, and 18, and their numerous varieties are very generally cultivated for their pulse. 18 Several varieties. 19 T. Kar-alsana.

* This may be reckoned the best Indian substitute for hemp.

angustifolius, R.			Mr. W. Kerr, 1812.	⊙ Tw.	R. S.	C. S.
semierectus.		America.	Mr W. Hamilton.	⊙	D. S.	D. S.
II. *Not twining.*						
nanus.		India.		⊙ D.	C. S.	C S.
radiatus.	B. Mash-kulai.1	India.		⊙ D.	C. S.	C S.
Max.	B. Kalla moog.2	India.		⊙ D.	C. S.	C. S.
Mungo.	B. Hali moog.3	India.		⊙ D.	C. S.	C. S.
glaber, R.		Mauritius.	Mons. Cossigny.	⊙ E.	R. &C. S.	D. S.
aureus, R.	B. Sona-moog.	Bengal.		⊙ E.		
torosus, R.	N. Sweta-mas.	Napaul.	Dr. F. Buchanan, 1801.	⊙ E.	C. S.	C. S.
trilobus.	B. Mooganee.4	India.	Indigenous.	H. D.	C. S.	C. S.
aconitifolius.	H. Moot.	Interior of Bengal.	Dr. W. Hunter.	⊙ D.	C. S.	C. S.
DOLICHOS.						
gladiatus.	B. Mukhun-shim.5	India.	Before 1794.	♃ Tw.	R. S.	C. S.
1 Red.	B. Ranga-Makhon-shim.	India.	Before 1794.	♃ Tw.	R. S.	C. S.
2 Large white.	B. Burra-Makhon-shim	India.	Mr. M. R. Smith, 1810.	♃ Tw.	R. S.	C. S.
3 Small white.	B. Makhun-shim.	India.	Before 1794.	♃ Tw.	R. S.	C. S.
4 Grey.	Not cultivated.	Pegu.	Mr. F. Carey, 1809.	♃ Tw.	R. S.	C. S.
virosus, R.6	H. & B. Kala-shim.	India.	Indigenous.	♂ Tw.	R. S.	C. S.
rotundifolius.7			1804.	♂ Tw.	R. S.	C. S.
obcordatus, R.		China.	Mr. W. Kerr, 1808.	♃ Tw.	R. S.	C. S.
sinensis.	H. Lobia.8	India.	1796.	⊙ Tw.	R. S.	C. S.
Catjang.	H. Burbutee.9		1796.	⊙ Tw.	D. S.	D. S.
tetragonolobus.10			1799.	⊙ or H. Tw	C. S.	C. S.
Lablab.11	S. Shimbee.			☾ or ♂ Tw.	D. S.	H. S.
1 var.	B. Bun-Shim.12					
2 var.	No name.13					
3 var.	T. Annapa.					
4 var.	T. Annapa-chicureay.14					
5 var.	B. Jeea-Shim.15					
6 var.	B. Goordal-shim.					
7 var.	D. purpureus.16					
hexandrus, R.	B. Kussa-gila.	Silhet.	Mr. M. R. Smith, 1811.	♄ Tw.	R. S.	C. S.
lignosus.17		India.	Before 1794.	☉ or ♂ Tw.	D. S.	H. S.
1 var.	B. Panch-shim.18					
2 var.	B. Bagh-nukhooa-shim.					
3 var.	B. Sada-jamai-pooli shim.19					
4 var.	B. Pitulee-jamai-pool-shim.					
5 var.	B. Doodha-pitulee-shim.					
6 var.	B. Kantchi-shim.					
bulbosus.20		Sumatra, &c.	Dr. C. Campbell, 1800.	H. Tw.	R.&C.S.	D. S.
gangeticus, R.	H. Bun burbutee.	Bengal.	Indigenous.	♃ Tw.	○	○
falcatus.	T. Itoo-monghy-tiga.	Coromandel.	Nalla-mandu, 1803.	H. Tw.	D. S.	D. S.
glutinosus, R.	H. Shimi-batrajee.21	India.	Indigenous.	♃ Tw.	C. S.	C. S.
pilosus.	B. Masha-kulai.			♃ Tw.	R.&C.S.	C. S.
biflorus.	H. Koolthee.22		Before 1794.		C. S.	C. S.
Soja.23		China.	Dr. C. Campbell.	⊙ E.	C. S.	C. S.
medicagineus.	T. Gadi-chikurea.	India.	Indigenous.	♂ Tw.	C. S.	C. S.
scarobœoides.	B. Banur kulai.24			♂ Tw.	C. S.	C. S.
phaseoloides, R.		China.	Mr. W. Kerr, 1804.	♃ Tw.	11---12	1---2
fabæformis.	T. Goor-chikurkai.	Coromandel.	Dr. John, 1797.	⊙ E.	D. S.	
unguiculatus.		W. Indies.	Captain Hewitt, 1809.	⊙	○	D. S.
azureus, R.		Silhet.	Mr. M. R. Smith, 1810.	♃ Cr.	10---11	C. S.
GLYCINE.						
debilis.	H. & B. Mashanee.25	India.	Indigenous.	♃ Tw.	R. S.	C. S.
coccinea.		New Holland.	Sir A. Hume, 1812.	♄ Tw.		
labialis.	H. & B. Bun-kulai.			♃ Tw.	R. S.	C. S.
frutescens.		America.	W. Hamilton, Esq. 1804.	♃		
bimaculata.		New S. Wales.	Mr. Bowie, 1801.			
CYLISTA.						
tomentosa, R.		Mysore.	Dr. Buchanan.	♄ Tw.	R.&C.S.	C. S.
scariosa.		Circars.	Mr. Heshusius.	♄ Tw.	R.&C.S.	C. S.
CLITORIA.						
ternata blue.	B. Upurajita.26	India.	Indigenous.	♄ Tw.	○	○
white.	B. Shet Upurajita.27			♄ Tw.	○	○
virginiana.		America.	Hon. C. Greville, 1807.	♃ Tw.	○	○
brasiliana,		Trinidad.	W. Lockhead, Esq. 1808.	♃ Tw.	○	○

1 S. Masha. B. T'hikiree. 2 B. Krishna-moog. T. Nella-pessara. 3 S. Moodga.
4 S. Moodga-purnee. T. Pilli-pessara. 5 S. Shimbee. Four varieties, three of which are cultivated.
6 H. M. 8. t. 45. 7 H. M. 8. t. 48. 8 Two varieties cultivated, viz. white and brown. 9 T. Bobra. Three varieties cultivated, viz. with white, brown, and black seeds. 10 Cultivated for our table. 11 Of this we have seven varieties. 12 Or spicatus. K. Mss. H. A. t. 141. f. 1.
13 These two are wild, and never made any use of. 14 D. Bengalensis. Willd. H. A. 5. t. 137.
15 H. A. t. 136. 16 Willd. The last five are cultivated during the dry season, for their pulse.
17 Of this we have six varieties known to me, all cultivated during the dry season for their pulse. 18 D. Unguiculatus. Loruier. 19 D. Cultratus. Willd. 20 H. A. 4. t. 102. f. 2. 21 T. Kar-kandi.
22 S. Kooluttha. T. Woolawooloo. Two varieties, one with brown, the other black seeds, are cultivated for their pulse. 23 Two varieties, one with white seed, and yellow flowers, the other black seed, and purple flowers
24 H. Belai-chooi. 25 S. Masha-purnee. 26 S. Asphota, also Upurajita. T. Nella dintona.
27 T. Tella dintona.

heterophylla.		Mauritius.	M. Cere, 1810.	S. ♄ Tw.	♀	♀
erecta.		America.		☿ E.	K.&C.S.	C. S.
PISUM.						
sativum, majus.	Bura-mutur. H.1			☉ Sc.	C. S.	C. S.
virida,	H.&B.Chota-mutur.2			☉ Sc.	C. S.	C. S.
	Small white field.3			☉ Sc.	C. S.	C. S.
LATHYRUS.						
Aphaca.	B. Chuna.4	Bengal.	Accident.	☉ D.	D. S.	D. S.
albus.		Europe.	M. De Cossigny.		D. S.	D. S.
sativus.	B. Khesaree.	Bengal.		☉ D.	D. S.	D. S.
tingitanus.		Barbary.	1799.	☉	C. S.	H. S.
odoratus.		Sicily.	1799.	☉	C. S.	
palustris.		Europe.	1804.		C. S.	
VICIA.						
sativa.	H. Ankuri.5	Bengal.	Indigenous.	☉ D.	D. S.	D. S.
Faba, minor.	Horse-bean.	Napaul.	B. Blake, Esq. 1811.	☉ E.	D. S.	D. S.
ERVUM.						
hirsutum.	B. Mussoor.6	Bengal.	Indigenous.	☉ D.	D. S.	D. S.
CICER.						
Lens.	B. Mussoor.	Bengal.	Indigenous.	☉ D.	D. S.	C. S.
arietinum.	B. Chhala.7	India.		☉ D.	C. S.	C. S.
CYTISUS.						
Cajan.	B. & H. Urur.8	India.	Before 1794.	☿ E.	C. S.	C. S.
hirsutus.		Europe.	W. Hamilton, Esq.	☿ E.		
purpureus.		Europe.	Sir A. Hume, 1812.	♄		
nigricans.		Europe.				
Laburnum.		Europe.				
ROBINIA.						
candida, R.	B. Batas-mundulee.	Bengal.	Dr. W. Carey, 1797.	♄ E.	♀	C. S.
suberosa, R.		Bengal.	Dr. W. Carey, 1799.	♄ E.	♀	C. S.
fruticosa, R.			C. Boddam, Esq. 1803.	♄ Sc.	R. S.	C. S.
racemosa, R.		Coromandel.	H. C. 1803.	♄ Sc.	R. S.	9—10
macrophylla, R.		Bengal.		♄ Sc.	6—7	4
COLUTEA.						
arborescens.		Europe.	W. Hamilton, Esq.	♄ E.		
frutescens, R.		Cape.	1799.	♄ E.	5—7	9—10
CORONILLA.9						
sanguinea.			Captain Learmonth, 1812.			
ÆSCHYNOMENE.						
grandiflora, white & red.	B. Buka.10	India.	Indigenous.	S. T.	♀	C. S.
Sesban.	S. Juyuntee.11	India.	Indigenous.	S. T.	R.&C.S.	C. S.
spinulosa, R.	T. Yerra-jeelooga.12	Coromandel.	Nalla-mandu, 1808.	☉ E.	12—1	2—4
paludosa.	B. Shola.13	Bengal.	Indigenous.	☉ E.	R. S.	C. S.
cannabina.14	B. Dhuncha.		Before 1794.	☉ E.	8—9	11—12
procumbens, R.	T. Manchi-jeelooga.	India.	Indigenous.	☉ D.	10—11	
uliginosa, R.		Bengal.		☉ D.	R. S.	C. S.
Pseudo-acacia.		N. America.	Dr. W. Carey, 1808.			
hispida.		America.	Dr. W. Carey, 1807.			
FLEMINGIA.						
procumbens, R.		Sewalick mts.	Accident.	☿ D.	C. S.	H. S.
nana, R.	H. Lipta-busunt.	Hindoostan.	Captain Hardwicke.	☿ E.	H. S.	
congesta, R.	B. Bura-Salpan.	Deltaof the Ganges.	Mr. Meyne, 1796.	♄ E.	R. S.	C. S.
stricta, R.	T. Guidda.	Circars.	1794.	♄ E.	♀	♀
semialata, R.		Napaul.	Dr. F. Buchanan, 1801.	♄ E.	R.&C.S.	C. S.
lineata, R.15		Coromandel.	Dr. Klein, 1800.	♄ E.	R. S.	C. S.
prostrata, R.		China.	Mr. Kerr, 1805.	☿ D.	D. S.	D. S.
SMITHIA.						
sensitiva.	B. Nala-kashina.16	Bengal.	Indigenous.	♂ D.	R. S.	C. S.
aspera, R.	B. Bhat-Shola.	India.		☉ E.	R. S.	C. S.
HEDYSARUM.						
I. *Leaves simple.*						
Alhagi.	H. Juwasa.17	Hindoostan.	Colonel Hardwicke, 1810.	H. E.		
moniliferum.	T. Amera.	India.	Indigenous.	H. E.	R.&C.S.	C. S.
vaginale.				☿ C.	R.&C.S.	C. S.
procumbens, R.		Coromandel.	1800.	☿ D.	R.&C.S.	C. S.
gramineum.	B. Pan-luta.18	India.	Gardeners, 1797.	♂ E.	R.&C.S.	C. S.
bupleurifolium.	H. Allooria.	Bengal.	Indigenous.	☿ D.	R.&C.S.	C. S.
styracifolium.		Coromandel.	Nalla-mandu, 1803.	☿ E.	R.&C.S.	C. S.
glumaceum.			Nalla-mandu, 1802.	♂ D.	R.&C.S.	C. S.
triquetrum.		India.	Before 1794.	H. D.	R.&C.S.	C. S.
alatum, R.	H. Kusrunt.	Cawnpore.	Colonel Hardwicke, 1801.	♄ E.	R.&C.S.	C. S.
gangeticum.	H. & B. Shalpani.19	India.	Indigenous.	♄ E.	R.&C.S.	C. S.

1 Or Marrow-fat. 2 Or field pea. 3 Or Patna pea. 4 Or Bura chima.
5 H. Jhunjhuni-ankuri. 6 S. Musoora. 7 Also Bhoot and Bhoot-kalai. S. Chhunuka.
T. Sunaga. 8 S. Arhukee. T. Kandy. 9 See Æschynomene. 10 Or Bukpoosh'pa. S. Vuka. H. Bassana. T. Avaki. Tam. Agati. 11 Yellow and purple flowered.
H. & B. Juyuntee, or Jainti. T. Siminta. 12 Coronilla aculeata, Willd. 13 B. Kath Shola.
14 Coronilla, Willd. Cultivated during the rains for the fibres of its bark, of which coarse cordage is made.
15 Hedysarum lineatum Linn. 16 H. Oda-breemi. 17 S. Yasa, or Yavusa. 18 H. Bunkajooli. 19 S. Shala-purnee.

colinum, R.		Circars.	Nalla-mandu, 1802.	♄ E.	R.&C.S.	C. S.
latifolium, R.		Hindoostan.	Colonel Hardwicke, 1801.	♄ E.	R.&C.S.	C. S.
strobiliferum.		India.	Before 1794.	♄ E.	R.&C.S.	C. S.
bracteatum, R.		Bengal.	Dr. W. Carey, 1801.	♄ E.	R.&C.S.	C. S.
gyrans.	H. Gora-chand.	Bengal.		♃ E.	R.&C.S.	C. S.
vespertilionis.	B. Chilia-tangi.	Bengal.	Dr. F. Buchanan.	☉ or ♂ E.	○	○
II. *Leaves conjugate.*						
diphyllum.		India.	Indigenous.	H. C.	R.&C.S.	C. S.
III. *Leaves ternate.*						
triflorum.	H. Kodăliya.	India.	Indigenous.	♃ C.	○	○
reptans, R.	[koolya.1	Bengal.	Dr. W. Carey.	♃ C.	R.&C.S.	C. S.
lagopodioides.	B. Gorukhya cha-	India.	Indigenous.	♃ Cr.	R.&C.S.	C. S.
hamosum, R.		Bengal.	Dr. W. Carey.	♃ D.	R.&C.S.	C. S.
alopecuroides, R.		Cawnpore.	Colonel Hardwicke, 1802.	♃ D.	R.&C.S.	C. S.
auriculatum, R.		Bengal.	Indigenous.	☾ Cr.	R.&C.S.	C. S.
orbiculatum, R.	B. Moonjiwra.	Silhet.	Mr. M. R. Smith, 1811.	♄ Sc.	9---10	C. S.
viscidum.		Coromandel.	Nalla-mandu, 1803.	♃ Cr.	R.&C.S.	C. S.
diffusum, R.		Bengal.	Uncertain.	♄ D.	R.&C.S.	C. S.
recurvatum, R.		Cawnpore.	Captain Hardwicke, 1802.	♄ E.	R. S.	C. S.
divergens, R.		Silhet.	Mr. M. R. Smith, 1812.	♄	C. S.	C. S.
purpureum, R.		Bootan.	Dr. W. Carey.	♃ D.	R. S.	C. S.
cephalotes, R.		Bengal.	Dr. W. Carey.	S. T.	R.&C.S.	C. S.
pulchellum.	B. Juta-Shalpanee.	India.	Before 1794.	♄ E.	R.&C.S.	C. S.
canescens.		N. America.	H. C. 1806.	♃ E.	R.&C.S.	C. S.
paniculatum.		America.	Mr. W. Lockhead, 1808.	♄ E.	R.&C.S.	C. S.
patens, R.		Moluccas.	Accident, 1798.	♃ D.	R.&C.S.	C. S.
tuberosum, R.	B. Shimeeya-batrajee.	N. Circars.	Captain Denton, 1795.	♄ T.	R. S.	C. S.
sp. nova.		America.	W. Hamilton, Esq.			
junceum.		Napaul.	Dr. F. Buchanan, 1801.	H. E.	R.&C.S.	C. S.
gyroides, R.	B. Biniyar.	Silhet.	Mr. M. R. Smith, 1811.	♄ E.	10---11	C. S.
rhombifolium, R.		Hindoostan.	Colonel Hardwicke, 1811.	♄ E.	10---11	C. S.
IV. *Leaves pinnate.*						
pictum.	B. Shunkur juta.2	India.	Indigenous.	♃ E.	R.&C.S.	C. S.
crinitum.		Chittagong.	Mr. Maddern, 1801.	♃ D.	R.&C.S.	C. S.
coronarium.		Europe.	Sir A. Hume, 1812.	♃ D.		
sennoides.		Coromandel.	B. Heyne, Esq. 1808.	♄ E.	○	○
lagenaria, R.	H. & B. Phool-shola.3	India.	Indigenous.	♃ A.	R. S.	C. S.
Neli-tali, R.	T. Tiga jeelooga.4			H. D.	R. S.	C. S.
INDIGOFERA.						
linifolia.	B. Bhanga.	India.	Indigenous.	♃ C.	○	○
trita.			Dr. F. Buchanan.	☾ or ♂ E.	○	○
glandulosa.	T. Bura-gadam.	Coromandel.	Nalla-mandu.	♃ D.	R.&C.S.	C. S.
argentea.		India.	Dr. F. Buchanan.	♄ E.	R.&C.S.	C. S.
uniflora.		Mysore.	Dr. F. Buchanan, 1800.	♃ D.	○	○
fragrans.		Coromandel.		♃ E.	R.&C.S.	C. S.
enneaphylla.	T. Cherra-gaddam.	India.	H. Russell, Esq.	♃ C.	○	○
hirsuta.	T. Penni-vempali.	Coromandel.	Dr. Klein.	☾ or ♂ E.	R.&C.S.	C. S.
viscosa.5		India.	Dr. Rottler.	♃ E.	R.&C.S.	C. S.
cœrulea, R.	T. Kar-neelee.	Circars.	1794.	♄ E.	R.&C.S.	C. S.
Anil.		Uncertain.	1810.	♄ E.		
tinctoria.	H. & B. Neel.6	India.	Indigenous.	♄ E.	R.&C.S.	C. S.
elliptica, R.		Bengal.	C. Boddam, Esq. 1803.	♄ E.	2---3	
atropurpurea.		Napaul.	Dr. F. Buchanan.	♄ E.	R.&C.S.	C. S.
pulchella, R.		Coromandel.	1800.	S. T.	R. S.	C. S.
arborea, R.		Sirinagur.	Colonel Hardwicke.	S. T.	2---3	
virgata, R.		Napaul.	Dr. F. Buchanan, 1802.	♄ E.	H. S.	
purpurescens, R.		Coromandel.	Mr. Peter Julian, 1809.	S. T.	H. S.	H. S.
violacea, R.		Hurdwar.	Mr. G. Roxburgh, 1808.	♄ E.	2---3	
Sp. undetermined.		Cuba.	W. Hamilton, Esq. 1800.	♄ E.	R.&C.S.	C. S.
uncinata, R.		Pegu.	Mr. F. Carey, 1809.	♄ E.	R. S.	C. S.
GALEGA.						
pentaphylla, R.		Mysore.	B. Heyne, Esq.	♄ E.	C. S.	C. S.
Heyniana, R.			B. Heyne, Esq.	♄ E.	C. S.	C. S.
incana.			B. Heyne, Esq. 1801.	♄ D.	C. S.	C. S.
villosa.		India.	Indigenous.	♃ D.	○	○
purpurea.	B. Surpunkha.7		1799.	♃ D.	C. S.	C. S.
officinalis.	Goat's rue.	Europe.	Dr. P. Russell, 1801.	♃		
lanceæfolia, R.		Mysore.	B. Heyne, Esq. 1802.	♄ D.	C. S.	C. S.
tinctoria.			Dr. F. Buchanan.	♃	R.&C.S.	C. S.
diffusa, R.		Circars.	Nalla-mandu, 1804.	♃ D.	C. S.	C. S.
ochroleuca.			Dr. Jacquin, 1806.	♃ E.	R. S.	
ASTRAGALUS.						
hamosus.		Persia, &c.	General Malcolm, 1801.		C. S.	C. S.
carolinianus.		N. America.	W. Hamilton, Esq.			
Sp. uncertain.			W. Hamilton, Esq. 1802.			

1 S. Prishnipurnee. H. Goluk-chakulee. 2 H. Dabra. 3 Æschynomene, L. 4 H. M. 9. t. 18. Æschynomene, L. 5 La Marck. 6 S. Neelee. T. Neelee. 7 Shura-punkha.

PSORALEA.						
corylifolia.	H. & B. Hakooch.1	India.	Indigenous.	⊙ E.	R.&C.S.	C.S.
pinnata.		Cape of G. H.	1799.	♄ E.		
aphylla.		Cape of G. H.	1799.	♄ E.		
TRIFOLIUM.						
indicum.	B. Bun-methhee.2	India.	Indigenous.	⊙ E.	D. S.	D. S.
officinale.	H. Ban-piring.	Bengal, &c.		⊙ E.	D. S.	D. S.
spumosum.		S. of Europe.	A. B. Lambert, Esq. 1802.	⊙	D. S.	D. S.
repens.	*White or Dutch clover.*	Britain.	1800.	♃ D.		
pratense.	*Honey suckle clover.*		B. Boswell, Esq. 1796.	♃ D.		
agrarium.	*Hop Trefoil.*	Europe.	B. Boswell, Esq. 1796.	⊙		
stellatum.	*Star-headed trefoil.*	England.	Accident or Col. Garstin.	⊙	D. S.	D. S.
angustifolium.	Narrow-leaved Tre-	Europe.	Accident.	⊙		
alexandrinum.	[foil.	Egypt.	T. Graham, Esq. 1803.	⊙ E.	D. S.	D. S.
cœrulium.		Germany.	A. B. Lambert, Esq. 1802.	⊙ E.	D. S.	D. S.
LOTUS.		[lands.				
jacobæus.		Cape verd Is-	A. Farquhar, Esq. 1798.	♄ E.	R.&C.S.	
tetragonolobus.		Europe.	1804.	⊙		
TRIGONELLA.						
Fœnum-græcum.	B. Methee-shak.3	India.	Indigenous.	⊙ E.	C. S.	C. S.
corniculata.	B. Piring.4	India.	Before 1794.	⊙ D.	C. S.	C. S.
MEDICAGO.						
sativa.	E. Lucern.	England.	B. Boswell, Esq. 1796.	♃ D.	O	O
lupulina.		Europe.				
polymorpha.5		Duab, &c.	Colonel Hardwicke.	⊙ D.	D. S.	D. S.
lappacea.		Bengal.	Dr. F. Buchanan, 1812.			

1 T. Kour-gestum. 2 H. M. 9. t. 40. S. Koostha-nashinee. 3 Methika. 4 S. Lunkaika. Prikka. 5 Two varieties.

CLASSIS XVIII.

POLYADELPHIA DODECANDRIA.

THEOBROMA.1
BUBROMA.2
ABROMA.3

ICOSANDRIA.

CITRUS.						
acida, R.	B. Neboo, &c.4					
1 var.	H.&B.Patee-Neboo.5	Bengal.	Before 1794.	S. T.	H. S.	7---9
2 var.	H.&B.Kagujee-Neboo.6		Before 1794.	S. T.	H. S.	7---8
3 var.	H.&.B.Gora-Neboo.7	Bengal.	Before 1794.	S. T.	H. S.	9---10
4 var.	China-gora-Neboo.8	China.	Before 1794.	S. T.	H. S.	8---9
5 var.	H.&B.Kamaral-Neboo.9	Bengal.	Before 1794.	S. T.	H. S.	11
6 var.	Rungpore.10	Rungpore.	1799.	S. T.	H. S.	8
7 var.	H.&B.Taba-Neboo.11	Bengal.	Before 1794.	S. T.	H. S.	11---12
8 var.	Arabian Lime.12	Arabia.	1800.	S. T.		
9 var.	H. &B. Meetha-Neboo.13		Before 1794.	S. T.	H. S.	R. S.
medica.	H.&B.Korna-Neboo.14		1796.	S. T.		7---8
1 var.	H.&B.Beg-poora.15		S. Hutteman, Esq. 1799.	S. T.	O	O
2 var.	Digitated Citron.16	China.	Mr. W. Kerr, 1811.			
3 var.	Wild Citrons.17	Garrow-hills.	Mr. M. R. Smith, 1810.	S. T.		
Aurantium.	S. Nagrunga.18	Silhet, &c.	Before 1794.	S. T.	D. S.	C. S.
decumanus.	H.&B.Batavi-Neboo.19	Uncertain.	1794.	S. T.	D. S.	9---11
myrtifolia.			W. Hamilton, Esq. 1800.	♄ E.		
inermis, R.	Ch. Kum-guat.	China.	Mr. W. Kerr, 1810.	♄ E.	4---5	8---9

POLYANDRIA.

MELALEUCA.						
Leucadendron.	Mn. Caju-Puti.20	Moluccas, &c.	Lieut. M'Kenzie, 1810.	T.		
Cajuputi.	Mn. Caju-Puti.21		Dr. J. Fleming, 1798.	S. T.	O	O
DURIO.						
zibethinus.	Mn. Durio.22	Moluccas, &c.	W. Ewer, Esq. 1809.	T.		
SYMPLOCUS.23						
HYPERICUM.	[taing.					
monogynum.	Ch. Keem-see-jau-	China.	Before 1794.	♄ E.	C. S.	
cernuum, R.		Sirinagur.	Col. Hardwicke.	♄ E.	C. S.	
canadensis.		America.	W. Hamilton, Esq.			

1 See Monadelphia. 2 See Monadelphia. 3 See Monadelphia. Its bark abounds in strong very white fibres, which make good ropes, &c. 4 Common Lime. S. Jumbeera, Nimbooka. H. Lemoo, Lembo. 5 Common round Lime. 6 Long small Lime. 7 Thick skinned small oval Lime. 8 Or China yellow-juiced Lime. 9 Large oval, smooth-skinned Lime. 10 Round, smooth-skinned Lime. 11 A large, globose, spongy-skinned Lime. 12 A large thick skinned sort from Muscat. 13 Round sweet Lime. 14 Common Lemon. A. Lemoo. 15 Common Citron. A. Utrej. S. Beeja-poora. P. Turenj. 16 Ch. Hevong yeum. 17 Three varieties from the Garrow-hills. 18 Silhet orange. B. Komla-neboo, or Kumla. H. Narungee. 19 Shaddock. 20 H. A. 2. t. 16. 21 Or Caju-kilan. H. A. 2. t. 17. f. 1. 22 Cin. Ratta-goorka. 23 See Polyandria Monogynia.

CLASSIS XIX.

SYNGENISIA ÆQUALIS.

ARNOPOGON.						
picroides.		Europe.	Dr. Jacquin, 1806.	⊙	3	
SCORZONERA.						
hispanica.		Europe.	1799.	♃		
tingitana.		Barbary.	Dr. W. Carey, 1808.	⊙		
SONCHUS.						
orixensis, R.	B. Bun-palung.1	India.	Indigenous.	⊙ E.	D. S.	H. S.
oleraceus, lævis.	T. Etrinta.			☉ E.	D. S.	H. S.
asper.		Accident.				
LACTUCA.						
sativa.	P. Kahoo.2	Persia.	Before 1794.	⊙ E.	D. S.	H. S.
procumbens, R.		India.	Indigenous.	⊙	3—4	4—6
PRENANTHES.						
acaulis, R.		Bengal.	Indigenous.	H.	C. S.	H. S.
linifolia, R.		Bengal.	D. W. Carey, 1812.	♃	C. S.	C. S.
racemosa, R.		Circars.		☉ E.	C. S.	H. S.
sonchifolia.		Circars.		⊙ E.	C. S.	H. S.
procumbens, R.		Bengal.	Indigenous.	H. D.	C. S.	H. S.
asplenifolia.	H. Teak-chama.	India.		H. D.	C. S.	H. S.
lyrata.						
LEONTODON.						
Taraxacum.	E. Dandelion.	Europe.	1807—8.	H.		
APARGIA.						
hamones.		Napaul.	Dr. F. Buchanan, 1802.	⊙ E.	C. S.	H. S.
CREPIS.						
fœtida.		Europe.	Dr. P. Russell, 1803.	⊙	C. S.	
HYOSERIS.						
cretica.		Candia.	A. B. Lambert, Esq. 1802.		C. S.	
KRIGIA.						
virginica.		N. America.	Accident.	⊙		
CICHORIUM.						
Endivia, broad-leaved.	P. Koshee.	India.		⊙ E.	C. S.	H. S.
curled.				⊙ E.	C. S.	H. S.
Intybus.		Europe.		H. E.	C. S.	H. S.
SCOLYMUS.						
maculatus.		Europe.	Accident, 1802.	⊙	C. S.	
SERRATULA.						
anthelmentica, R.3		India.	Indigenous.	⊙ E.	D. S.	D. S.
cinerea, R.	H. Choota kookshima.4			⊙ E.	D. S.	D. S.
carthamoides, R.		Napaul.	Dr. F. Buchanan, 1802.	⊙ E.	D. S.	D. S.
CARDUUS.						
lanatus, R.5		Bengal.	Indigenous.	H. E.	D. S.	H. S.
marianus.		Europe.	1799.	⊙ E.	D. S.	H. S.
radicans, R.		Mysore.	Dr. F. Buchanan.	⊙ D.	D. S.	H. S.
lanceolatus.6		Europe.	Accident, 1802.	⊙ E.	D. S.	
tenuiflorus.		England.	A. B. Lambert, Esq. 1802.	⊙ E.	D. S.	
syriacus.7		Syrica, &c.	Mr. Salisbury, 1807.	⊙ E.	D. S.	
ONOPORDUM.						
Acanthium.		Europe.			C. S.	
CIRSIUM.						
horridulum.			Mr. M'Mahon, 1810.	♂ E.	C. S.	H. S.
CYNARA.						
Scolymus.	P. Hershuf.8	Uncertain.		♃ E.	D. S.	H. S.
CARTHAMUS.						
tinctorius.	S.H.B.&T.Koosum- [bha.9	India, &c.	Before 1794.	⊙	C. S.	H. S.
SPILANTHUS.						
acmella.		India.	Indigenous.	H. D.	R.&C.S.	C. S.
oleracea.	B. Roshuniya.		W. Hamilton, Esq.	⊙ E.	○	○
BIDENS.						
bipinnata.		Rajemahl.	Mr. W. R. 1801.	⊙ E.	R.&C.S.	H. S.
tripartita.		Napaul.	Dr. F. Buchanan, 1802.	⊙ E.	R.&C.S.	H. S.
CACALIA.						
villosa, R.		China.	Mr. W. Kerr, 1812.	⊙ E.	C. S.	C. S.
sonchifolia.	B. Shudi-mudi.10	India.	Indigenous.	⊙ E.	C. S.	C. S.

1 T. Nalla-tapata. 2 A. Khussa. 3 Vernonia, Willd. 4 Conyza, L.
5 Cnicus. Willd. 6 Cnicus. Willd. 7 Cnicus Willd. 8 H. Kungir. 9 H. Kosumbh, or Kooshoom.
10 H. M. 10. t. 68.

bicolor.		Moluccas.	Mr. C. Smith, 1798.	♃ D.	C. S.	C. S.
hieracioides.		Mauritius.	H. Hope, Esq. 1810.	♃ E.	☉	
ETHULIA.						
conyzoides.		India.	Dr. F. Buchanan.	☉ E.	C. S.	C. S.
ramosa, R.		Bengal.	Dr. W. Carey.	☉ E.	D. S.	H S.
EUPATORIUM.						
Aya-pana.		Brazil.	Capt. B. Blake, 1801.	♄ D.	C. S.	
divergens, R.		Mysore.	Dr. Buchanan.	♄ E.	D. S.	H. S.
sp. undermined.		Mauritius.	Mons. Cere, 1810.	♄ E.	C. S.	
asperum.		Bengal.	Dr. F. Buchanan, 1811.	♄ E.	C. S.	H. S.
AGERATUM.						
cordifolium, R.	B. *Ookunchi.*	Bengal.	Indigenous.	☉ E.	C. S.	H. S.
aquaticum, R.	B. *Bura-keshuti.*	India.	Indigenous.	☉ E.	D. S.	H. S.
ATHANASIA.						
crithmifolia.		Cape of G. H.	1798.	♄		
indica, R.	T. Bugati-bunda.	Coromandel.	Nalla-mandu.	☉ E.	B. S.	H. S.

SUPERFLUA.

TANACETUM.						
vulgare.		Britain.	Dr. Hornman, 1807.	♃ E.		
ARTEMISIA.						
vulgaris.	S. H. & B. Naga-dana.	Uncertain.	Before 1794.	H. E.	D. S.	H. S.
indica.	S. Duna, dana.1	India.	Before 1794.	H. E.	D. S.	H. S.
elegans, R.	H. Abbuttna.	Hindoostan.	Genl. Martin.	☉ E.	D. S.	H. S.
paniculata, R.		Persia.	Mrs. Honeycomb.	♃ E.	C. S.	H. S.
parviflora, R.		Napaul.	Dr. F. Buchanan.	H. D.	D. S.	
caruifolia, R.		Napaul.	Dr. F. Buchanan.	☉ E.	D. S.	H. S.
maderaspatana.2		India.	Indigenous.	☉ D		
hemisphærica, R.		Bengal.		☉ E.	D. S.	H. S.
sternutatoria, R.	B. Hachitee.3			☉ pr.	D. S.	H. S.
GNAPHALIUM.						
strictum, R.		India.	Indigenous.	☉ E.	D. S.	H. S.
depressum, R.		India.	Indigenous.	☉ pr.	D. S.	H. S.
alboluteum, R.		India.	Indigenous.	☉ E.	D. S.	H. S.
indicum.		India.	Indigenous.	☉ E.	D. S.	
undulatum.		Africa.	W. Hamilton, Esq.	☉		
CONYZA.		[Ganges.				
corymbosa, R.		Delta of the	Mr. Meyne, 1796.	♄ E.	C. S.	
balsamifera.		India.	Dr. F. Buchanan.	♄ E.	D. S.	H. S.
laciniata, R.		Bengal.	Indigenous.	☉ E.	C. S.	H. S.
lacera.	B. Burra-kooksima.4	India.	Indigenous.	☉ E.	C. S.	H. S.
diffusa, R.		Bengal.	Indigenous.	☉ E.	C. S.	H. S.
aurita.		Coromandel.		☉ E.	C. S.	H. S.
angustifolia, R.		Andamans.	Dr. F. Buchanan.	☉ E.	C. S.	H. S.
pinnatifida.		Napaul.	Dr. F. Buchanan.	♄ E.	H. S.	H. S.
alata, R.		Napaul.	Dr. F. Buchanan.	♄ E.	H. S.	
repanda, R.		Pegu.	Mr. F. Carey, 1809.	☉ E.	C. S.	H. S.
bifoliata.	H. Jurud-banghra.	Coromandel.		H. E.	C. S.	H. S.
lanceolaria, R.	B. Tejmool.	Silhet.	Mr. M. R. Smith, 1811.	♃ E.	3	5---6
stricta.		Bengal.	Accident, 1812.	☉ E.	C. S.	D. S.
ERIGERON.						
heterophyllum.		N. America.	W. Hamilton, Esq.			
purpureum.			W. Hamilton, Esq.			
asteroides, R.		India.	Indigenous.	☉	C. S.	
canadense, R.		America.	Accident.	☉		
carolinianum.		N. America.	1802.			
SENECIO.						
Jacobæa.		Europe.				
Sarracenicus.		Europe.	Accident, 1801.	♃		
vulgaris.		Europe.	Accident, 1803.	☉	D. S.	
ASTER.						
chinensis.	N. Gaynatuka.	Napaul.	Dr. F. Buchanan, 1802.	☉	C. S.	
dumosus.		America.	Accident, 1802.		5---8	
annuus.					8	
trinervius, R.		Napaul.	Dr. F. Buchanan, 1802.	♄ Sc.	C. S.	
DORONICUM.						
calcaratum, R.	T. Paleroopoo.	India.	Indigenous.	☉ E.	C. S.	H. S.
TAGETES.		[ree.5				
patula.	H. & B. Gundha-jaf-	Mexico.	Before 1794.	☉ E.	C. S.	H. S.
erecta.	H. & B. Gendha.	Mexico.	Before 1794.	☉ E.	C. S.	H. S.
ZINNIA.						
multiflora.		America.	A. Hogue, Esq. 1812.	☉ E.	D. S.	H. S.
CHRYSANTHEMUM.						
indicum.	B. Chundru-mullika.6	India.	One yellow var. before 1794.	♄ D.	C. S.	
segetum.		Europe.		☉	C. S.	

1 H. Duna, or Gundhumar. 2 H. M. 10. t. 49. Cotula. Willd. 3 Also Nukchiknee.
4 And Kookkura Shoonga. 5 P. Gool-jafree. 6 H. Gool-daoodi. T. Chamunti. Four varieties, viz. two yellow, one purple, and one white.

Leucanthemum. tricolor.		Europe.	Accident, 1807. Sir A. Hume, 1812.	⊙ E.	H. S.	
PYRETHRUM. indicum, R.		Bengal.	Dr. F. Buchanan, 1809.	⊙ E.	D. S.	H. S.
MATRICARIA. suaveolens.		Europe.	Genl. Martin, 1797.	⊙ E.	C. S.	H. S.
COTULA. sinapifolia, R.		Napaul.	Dr. F. Buchanan, 1802.	⊙ E.	H. S.	H. S.
ANTHEMIS. Cota.		Europe.		⊙	H. S.	
ECLIPTA. prostrata.	B. & H. Keshooriya.1	India.	Indigenous.	⊙ D.	O	O
SIEGESBECKIA. brachiata, R.		India.	Indigenous.	⊙ E.	R.&C.S.	C. S.
VERBESINA. Lavenia.2 calendulacea. biflora. scandens, R. sativa, R.	B. Keshooriya.	Circars. India.	Nalla-mandu. Indigenous.	⊙ E. ♃ Cr. ☺ E. ♃ Sc.	C. S. C. S. R.&C.S.	H. S. H. S. D. S.
	B. & H. Bun-raj. B. Ram-til.3	Bengal. India.	Mr. B. Heyne.	⊙ E.	C. S.	C. S.

FRUSTRANEA.

HELIANTHUS. annuus. indicus. multiflorus. decapetalus. tuberosus.	B. Soorjya-mookhee.4	Mexico.	Before 1794.	⊙ E. ⊙ E. ⊙ E	O	O
		America.	W. Hamilton, Esq. 1800.		R.&C.S.	C. S.
	Jerusalem artichoke.	Brazil.	Colonel Garstin,	H. E.	R.&C. S.	
RUDBECKIA. hirta. purpurea. triloba.		N. America.	W. Hamilton, Esq. W. Hamilton, Esq. J. Bentley, Esq. 1812.	♂ ♃ E.	C. S. C. S.	
COREOPSIS. tripteris.		N. America.	W. Hamilton, Esq.			
DAHLIA. sambucifolia.		S. America.	A. B. Lambert, Esq. 1807.	H. E.	C. S.	H. S.
CENTAUREA. benedicta. moschata. lanata, R.	H. Shah-pusund.	Europe. Persia. Elmora.	1799. Col. Dyer, 1800. F. Pierard, Esq. 1809.	⊙ E. ⊙ E. ⊙ E.	C. S. 3---4---5	H. S. 6---7

NECESSARIA.

CALENDULA. officinalis.		Europe.		⊙		
OTHONA. bulbosa. crispa.		Cape of G. H. Cape of G. H.				
PARTHENIUM. Hysterophorus.		W. Indies.	Accident, 1810.	☺	R. S.	R. S.

SEGREGATA.

ELEPHANTOPUS. scaber.	B. Sham-dulun.5	India.	Dr. W. Carey.	H. E.	C. S.	H. S.
SPHÆRANTHUS. mollis. indicus.	B. Moondee. B. Chhagul-nadi.	India. India.	Indigenous. Dr. W. Carey.	⊙ E.	C. S.	H. S.
ECHINOPS. echinatus, R.		Mysore.	Dr. F. Buchanan. W. Hamilton, Esq. 1809.	⊙ D.	C. S.	H. S.
BROTERA. ovata.						
CÆSULIA. axillaris, R.		India.	Dr. W. Carey.	⊙ D.	R.&C.S.	C. S.
HINGCHA. repens, R.	H. & B. Heluncha.6	Bengal.	Indigenous.	☺ Cr.	H.&R.S.	R. S.

1 Or Keyrati. H. M. 10. t. 41. 2 Lavenia erecta, Willd. 3 Can. Kutrellon.
4 Sun-flower. P. Guli aftaab. 5 S. Gojihwa. H. Sam-dullum. 6 S. Hilumochika.

CLASSIS XX.

GYNANDRIA MONANDRA.

ORCHIS.						
uniflora, R.	B. Kathiya-ram-ram.	Garrow-hills.	Mr. M. R. Smith, 1812.	E.	8---10	
commelinifolia, R.		Bengal.	B. Blake, Esq. 1810.	H. E.	7---8	
PTERYGODIUM.						
sulcatum, R.	B.&H.Shwet-hoolee.1	Bengal.	Indigenous.	H. E.	C. S.	C. S.
EPIPACTIS.						
graminifolia, R.		Garrow-hills.	Mr. M. R. Smith, 1812.	H. E.		
Juliana, R.		Bengal.	Indigenous.	H. E.	H. S.	
plicata, R.	H. & B. Lall-ghurree.	Bengal.	Indigenous.	H. E.	H. S.	H. S.
carinata, R.		Bengal.	Indigenous.	H. E.	H. S.	
babianifolia, R.	H. Chatiya shookar.	Garrow-hills.	Mr. M. R. Smith, 1812.	H. E.	R. S.	
MALAXIS.						
plicata, R.		Sumatra.	Mr. W. R. 1803.	♄ P.	R. S.	
CYMBIDIUM.						
nitidum, R.		Silhet.	Mr. M. R. Smith, 1812.	♃ P.	H. S.	
bambusifolia, R.		Chittagong.	Mr. J. R. 1810.	♃ E.	H. S.	
triste.		Delta of the Ganges.	W. Egerton, Esq. 1809.	♃ P.	H. S.	
alatum, R.		Chittagong.	Mr. J. R. 1810.	♃ P.	H. S.	
verecundum.		W. Indies.	H. C. 1806.	♃ H.	D. S.	
tessalloides, R.	S. Vrisha-roha.2	Bengal.	Indigenous.	H. S.	H.&R.S.	
premorsum.		Sumatra.	Mr. Griffiths.	♃ P.	H. S.	
amabile.		Amboyna.	Mr. C. Smith.	♃ P.	H. S.	
pendulum, R.		India.	Mr. M. R. Smith, 1811.	♃ P.	H. S.	
imbricatum, R.		Chittagong.	Mr. J. R. 1810.	♃ P.	9---10	
iridifolium, R.		Chittagong.	Mr. J. R. 1810.	♃ P.	10---11	
aloifolium.		Garrow-hills.	Mr. M. R. Smith, 1812.	♃ P.	4---5	
sinense.		China.	Mr. W. Kerr, 1812.	♃ T.	11---1	
striatum.		China.	Mr. W. Kerr, 1812.	♃ T.		
LIMODORUM.						
virens, R.	H. Gorooma.3	India.	Indigenous.	H. E.	H. S.	
ramentaceum, R.		Bengal.	R. Smith, Esq. 1810.	H. E.	2---3	
nutans, R.	H. Pankula.	Bengal.	Indigenous.	H. E.	H.&R.S.	
recurvum, R.		India.	B. Smith, Esq. 1810.	H.	6---7	
bicolor, R.	B. Budbuhuri.	Bengal.	--- Smith, Esq. 1810.	H. E.	H. S.	
Tankervilliæ.	B. Louralee.	China, Silhet.	Kew Garden. 1806.	H. E.	H. S.	
plicatum, R.		Sumatra.	Mr. W. R. 1803.	♃ E.	H. S.	
candidum, R.	B. Miarmee.	Silhet.	Mr. M. R. Smith, 1811.	♃ E.	H. S.	
bracteatum, R.		Garrow-hills.	Mr. M. R. Smith, 1811.	♃ E.	7---8	
longifolium, R.	B. Telungshoora.	Garrow-hills.	Mr. M. R. Smith, 1811.	H. E.	12---1	
AERIDES.						
radiatum, R.		Sunderbunds.	Dr. W. Carey, 1812.	♃ P.	6---7	
suaveolens, R.		Chittagong.	Mr. J. R. 1810.	♃ P.	H. S.	
guttatum, R.		Dacca.	Hon'ble C. A. Bruce.	♃ P.	H. S.	
cornutum, R.		Dacca.	Hon'ble C. A. Bruce.	♃ P.	H. S.	
rostratum, R.		Silhet.	Mr. M. R. Smith, 1811.	♃ P.	H. S.	
uncinatum, R.		Silhet.	Mr. M. R. Smith, 1811.	♃ P.	H. S.	
ampullaceum, R.		Garrow-hills.	Mr. M. R. Smith, 1812.	♃ P.	5---6	
pallidum, R.		Chittagong.	Mr. J. R. 1810.	♃ P.	H---S	
DENDROBIUM.						
aggregatum, R.		India.	--- Pierard, Esq.	♃ P.	H. S.	
tripetaloides, R.		Pegu.	Mr. F. Carey, 1809.	♃ P.	C. S.	
flavum, R.		Chittagong.	Mr. J. R. 1810.	♃ P.	2---3	
pumilum, R.		Chittagong.	Mr. J. R. 1810.	♃ P.	R. S.	
Pierardii, R.			Mr. Pierard.	♃ P.	H. S.	
purpureum, R.		Amboyna.	Mr. C. Smith.	♃ P.	H. S.	
cruminatum.		Amboyna.	Mr. C. Smith, 1798.	♃ P.	H. S.	
formosum, R.		Silhet.	Mr. M. R. Smith, 1811.	♃ P.	H. S.	
pulchellum, R.		Silhet.	Mr. M. R. Smith, 1811.	♃ P.	H S.	
anceps.		Bengal.	W. Egerton, Esq. 1811.	♃ P.	H. S.	
acinaciforme, R.		Amboyna.	Mr. C. Smith.	♃ P.	R. S.	
calceolum, R.		Amboyna.	Mr. C. Smith.	♃ P.	R. S.	
aphyllum.		Chittagong.	Mr. J. R. 1810.	♃ P.	H. S.	
teres, R.		Chittagong.	Mr. J. R. 1810.	♃ P.	H. S.	
crispatum.	[Jhoopee.	Chittagong.	Mr. J. R. 1810.	♃ P.	H. S.	
veratrifolium, R.	B. Kathaeea-bun-	Garrow-hills.	Mr. M. R. Smith, 1812.	♃ P.	4---5	
clavatum, R.	B. Titur-keoo.			♃ P.	4---5	
pendulum, R.		Chittagong.	Mr. J. R. 1810.	♃ P.	4---5	

1 Two varieties. 2 H. Bunda and Algooch. B. Alugnuta. 3 B. Goruk-phuna.

HEXANDRIA.

ARISTOLOCHIA.						
indica.	B. & H. *Ishurmool*.1	India.	Indigenous.	♄ T.	R. S.	C. S.
anguicida.		N. America.	W. Hamilton, Esq.	♄ T.	R. S.	
bracteata.	T. Gardi-garapa.	Coromandel.	Dr. A. Berry, 1799.	H. D.	O	O
odoratissima.		N. America.	W. Hamilton, Esq.	♄ T.		
acuminata, R.		Chittagong.	Mr. J. R. 1810.	♄ T.	H.&R.S.	C. S.

1 S. *Urka-moola*.

CLASSIS XXI.

MONOECIA MONANDRIA.

AMBROSINIA.						
spirale, R.		Tranquebar.	Mr. Rottler.	H. E.	R. S.	R. S.
retrospirale, R.		Bengal.	Dr. W. Carey.	H. E.	R. S.	
ciliata, R.	B. Keralee.	Bengal.	Indigenous.	H. E.	R. S.	R. S.
ARUM.						
I. *Leaves simple.*						
Colocasia.	S. Kuchoo.1					
1 var.	B. Goori-kuchoo.2	India, &c.	Before 1794.	H.	R. S.	
2 var.	B. Ashoo-kuchoo.3	India, &c.	Before 1794.	H.	R. S.	
1 var.	B. Kala-kuchoo.4	Bengal.	Indigenous.	H.	R. S.	
2 var.	B. Chhar-kuchoo.	Bengal.	Indigenous.	H.	R. S.	
3 var.	B. Bun-kuchoo.5	Bengal.	Indigenous.	H.	R. S.	
nymphæafolia.	B. Sar-kuchoo.6	India.	Indigenous.	H.	R. S.	
viviparium, R.7		Napaul.	Dr. F. Buchanan.	H. E.		
rapiforme, R.		Pegu.	Mr. F. Carey, 1809.	H.	R. S.	
montanum, R.	T. Konda-rakis.	Circars.	Mr. F. Carey, 1808.	H.	R. S.	
indicum.	B. & H. Man.8	India.	Indigenous.	♄ E.		
odorum, R.		Pegu.	Dr. W. Carey, 1809.	♄ E.		
cucullatum.	B. Bish-man.	India.	Before 1794.	♄ E.	R. S.	
fornicatum, R.	B. Kuchiree.9	Dacca, &c.		♄ E.		
flagelliforme, R.		Bengal.	Indigenous.	H.	R. S.	
divaricatum.		China.	Mr. W. Kerr, 1809.	H.	R. S.	
orixense, R.	B. Ghetkool.10	India.	Indigenous.	H.	R. S.	
trilobatum.		Moluccas.	Accident.	H.	R. S.	
gracile, R.	H. Harpung.11	Silhet.	Mr. M. R. Smith, 1811.	H. E.	H.&R.S.	
undetermined.12		S. Sea Islands.	Mrs. Blaxland, 1812.			
II. *Leaves compound.*						
cuspidatum, R.		Pinang.	Mr. W. R. 1802.	H.	R. S.	
curvatum, R.	H. Beer-buka.	Napaul.	King of Napaul, 1804.	H.	R. S.	
sessiliflorum, R.	H. Binda-gooea.	Cawnpore.	Col. Hardwicke, 1802.	H.	H. S.	
campanulatum, R.	H. & B. Ol.13	India.	Before 1794.	H.	H. S.	
bulbiferum, R.	B. Umbla-Bela.	Bengal.	Indigenous.	H.	R. S.	
margaretifer, R.		Hindoostan.		H.	H. S.	R. S.
CALLA.						
virosa, R.	H. Bish-kuchoo.	Chittagong.	Mr. J. R. 1810.	♄ E.	9---10	
rubescens, R.		Chittagong.	Dr. F. Buchanan.	♄ E.	R. S.	
oblongifolia, R.14		Moluccas.	Mr. C. Smith, 1798.	♄ E.	R. S.	
calyptrata, R.	[bee.	Amboyna.		H.	H. S.	
aromatica, R.	H. Kuchoo-gundhu-	Chittagong.	Mr. J. R. 1810.	H.	C. S.	
picta, R.		Sumatra.	Mr. W. R.	♄ E.		
æthiopica.		Cape of G. H.	Dr. W. Carey.			
FICUS.						
I. *Fruit solitary or paired.*						
carica.	B. Doomoor.15		Before 1794.	L. ♄	3---5	6---8
hirsuta, R.	B. Dungura.	Silhet.	Mr. M. R. Smith, 1810.	T.	H. S.	R. S.
caricoides, R.		Rohilcund.	General Martyn, 1796.	S. T.	O	O
virgata, R.			Mr. A. Gott, 1803.	♄ E.	O	O
hirta, R.	B. Bun Doomoor.	Silhet.	Mr. M. R. Smith, 1810.	T.	H. S.	R. S.
heterophylla.	B. Gooti sheora.16	India.	Indigenous.	♄ Sc.	O	O
scandens, R.		Chittagong.	Mr. J. R. 1810.	♄ Sc.	O	O
radicans, R.		Silhet.	Mr. M. R. Smith, 1811.	♄ R.	O	O
quercifolia, R.		Sumatra.	Dr. C. Campbell, 1803.	♄ D.	O	O
humilis, R.		Sumatra.	Mr. W. R. 1803.	♄ Cr.	O	O
repens.	H. Bhooi-doomoor.	Bengal.	Indigenous.	♄ D.	O	O
Goolereea, R.	H. Kuth Goolureea.	Hindoostan.		S. T.		
Wassa, R.	Mn. Wassa.	Moluccas.	H. C. 1801.	♄ E	R. S.	R. S.
elastica, R.	B. Kusmeer.17	Garrow hills.	Mr. M. R. Smith, 1810.	L. T.	H. S.	R. S.
indica.	B. Bur, or But.18	South of Asia.	Indigenous.	L. T.	H. S.	H. S.
religiosa.	B. Ashwuttha.19	South of Asia.	Indigenous.	L. T.	H. S.	R. S.

1 S. Kuchwee. Two varieties cultivated. 2 T. Shama-kura. 3 Besides those, there are three wild varieties. 4 H. A. 5. t. 110. p. 1. 5 H. A. 5. p. 318. 6 T. Tshama. 7 H. M. 12. t. 9. 8 Man-kuchoo and Man-giri, two varieties. 9 T. Balrakis, 10 Or Ghet-kuchoo. 11 H. M. 11. t. 21. 12 Tarra root of New Zealand. 13 S. Kunda. T. Muncha-kunda. 14 H. A. 5. t. 108. 15 P. Unjeer. A. Teen. 16 T. Burong. H. M. 3. t. 62. 17 Abounds in Caoutchouc. 18 S. Vuta, Nyugrodha, &c. T. Murri. H. M. 1. t. 28. 19 S. Pippula. T. Ray. Cin. Bogaha.

cordifolia, R.	B. Guya ushwuttha.	Bengal.		L. T.	H. S.	R. S.
infectoria.	B. Pakur.1	India.		L. T.	H. S.	R. S.
rapiformis, R.		Moluccas.	H. C. 1800.	T.	8	
excelsa.2		Moluccas.	H. C. 1800.	T.	8	
Tsiela, R.3		India.		L. T.	H. S.	R. S.
laccifera, R.	B. Kutul But.	Silhet.	Mr. M. R. Smith, 1811.	L. T.	H. S.	R. S.
vagans, R.	B. Luta-Doomoor.	Silhet.	Mr. M. R. Smith, 1811.	♄ Sc.	H. S.	
lucida,			Mons. Jannet, 1810.	T.		
II. *Fruit fascicled, racemed, or panicled.*						
exasperata, R.4		Moluccas.	H. C. 1800.	S. T.	9	12
polycarpa, R.		Moluccas.	H. C.	♄ E.	R. S.	R. S.
macrophylla, R.		Chittagong.	Before 1793.	T.	○	○
lanceolata, R.		Chittagong.	Dr. Buchanan, 1798.	S. T.	○	○
glomerata.	B. Jugya-doomoor.5	India.	Indigenous.	T.	○	○
conglomerata, R.		Chittagong.	H. Harris, Esq.	S. T.	○	○
congesta, R.		Moluccas.	Captain Anderson.	S. T.	○	○
racemifera, R.6		Sumatra.	Dr. C. Campbell, 1796.	S. T.	○	○
cunea, R.		Napaul.	Dr. Buchanan.	S. T.	○	○
oppositifolia, R.	B. Kaka-doomoor.7	India.	Indigenous.	S. T.	○	○
dæmona.		Tanjore.	Dr. Rottler, 1802.	♄ E.	○	○
ARTOCARPUS.						
incisa.8		S. Sea Island.	Before 1794.	T.		
integrifolia.	B. Kantal.9	India.	Indigenous.	T.	1---2	5---7
Lacucha, R.	B. Dephul.10	Bengal.		T.	3---5	R. S.
Chaplasha, R.	B. Chapulasha.	Chittagong.	Dr. F. Buchanan, 1798.	T.	2---5	R. S.
echinata, R.	M-n. Tampoene.	Malay Islands.	Mr. W. R. 1803.	T.	H. S.	
serrata, R.		Travancore.	G. Young, Esq. 1808.	T.		
angustifolia, R.		Malay Islands.	J. Scott, Esq. 1800.	T.		
GNETUM.						
Gnemon.11		Moluccas.	Lieut. M'Kenzie, 1810.	T.		
scandens, R.	B. Nana-wita.12	Chtttagong.	Mr. J. R. 1810.	L. Sc. ♄	3---4	9---10
CASUARINA.						
muricata, R.	H. Huri.	Chittagong.	Dr. F. Buchanan, 1798.	T.	R. S.	
equisetifolia.		S. S. Islands.	Colonel Paterson.	T.		
nodiflora.		S. S. Islands.	Mons. Jannet, 1810.	T.		
distyla.		New Holland.	Dr. Wallich, 1811.			
torulosa.		New Holland.	Captain Murray, 1811.			
rugosa.			Sir A. Hume, 1812.			
CHARA.						
verticillata, R.		Bengal.	Indigenous.	⊙ A.	C.&H.S.	
involucrata, R.	B. Jhanji.	Bengal.	Indigenous.	⊙ A.	C.&H.S.	
furcata, R.	B. Jhanjhi.	Bengal.	Indigenous.	⊙ A.	C.&H.S.	

DIANDRIA.

LEMNA.						
orbiculata, R.		Bengal.	Indigenous.	⊙ A.	R.&C.S.	
globosa, R.		Bengal.	Indigenous.	⊙ A.	R.&C.S.	
cruciata, R.		Bengal.	Indigenous.	⊙ A.	R.&C.S.	

TRIANDRIA.

TYPHA.						
elephantina, R.	B. Hogla.13	India.	Indigenous.	♃ E.	R. S.	
angustifolia.	B. Kam-Hogla.	Bengal.	Indigenous.	♃ E.	R. S.	
ZEA.						
Mays.	B. Mokka.14	America.	Before 1794.	⊙	R.&C.S.	R.&C.S
COIX.						
pumila, R.		Mauritius.		♃ E	R. S.	C. S.
lacryma.	B.&H. Jul gurgur.15	India.	Indigenous.	♃ E.	R. S.	C. S.
barbata, R.	H. Gurgur.16			♃ E.	R. S.	C. S.
gigantea, Kœn.	B. Danga-gurgur.	Bengal.		♃ E.	R. S.	C. S.
aquatica, R.			Kerum, 1804.	♃ A.	C. S.	H. S.
heteroclita, R.			Mr. R. Kyd, 1810.	⊙ E.	R. S.	C. S.
SCLERIA.						
biflora, R.		Bengal.	Indigenous.	⊙ ♂ E.	R. S.	C. S.
TRAGIA.						
involucrata, R.	B. Bichitee.17	India.	Indigenous.	♄ Tw.	R. S.	C. S.
mercurialis.		India.	Indigenous.	⊙ E.	R.&C.S.	C. S.
HERNANDIA.						
ovigera.		Moluccas.	Mr. C. Smith.	T.	8---10	H. S.
SERPICULA.						
verticillata.	T. Poo-natsoo.	India.	Indigenous.	⊙ A.	R. S.	R. S.

1 S. *Plucsha.* H. M. 3. t. 64. 2 H. M. 3. t. 58. 3 H. M. 3. t. 63. 4 H. A. 4. t. 63. 5 S. *Oodoombura.* H. Doombur. T. Bruma-madi. 6 H. A. 3. t. 93. 7 S. Kako-doomborica. H. Koriya-doombar. T. Boda-mamadi. 8 Bread-fruit tree. 9 The Jack. S. & T. Punusa. 10 S. Lukoocha. H. Barul. 11 H. A. 1. t. 71. 12 H. M. 7. t. 22. H. A. 5. t. 7 & 8. 13 T. Enuga-junum. 14 H. Bhutta. T. Moka-juna. 15 Or Jeel gurgur or Kanch. S. Guberoo. H. M. 12. t. 70. 16 T. Ghella-gaddy. 17 S. Vrishchi-putree.

ALNUS.						
dioica, R.	**Kokra, H.**	Bengal.	Mr. M. R. Smith, 1811.	L. T.	3---4	9---10
URTICA.						
scabrella, R.		Chittagong.	Mr. J. R. 1811.	♄ D.	8---9	10---11
dioica.		Europe.	Col. Garstin.		C. S.	
urens.		Europe.	1799.	☉ E.	C. S.	H. S.
alienata.		Ceylon.	General M'Dowal.	H. E.	R. S.	C. S.
parviflora.		Rohilcund.	Mr. A. Gott, 1803.	H. E.	H. S.	H. S.
tuberosa, R.	T. Pilla-dumpa.	India.	Indigenous.	H. D.	R. S.	C. S.
suffruticosa, R.		Sumatra.	Mr. W. Ewer, 1808.	♃ E.	○	○
pentandra, R.	B. Pipeera-saree.	India.	Indigenous.	♃ D.	○	○
interrupta.	B.&H.Lall-bichitee.	India.	Indigenous.	☉ E.	R. S.	C. S.
heterophylla.1		Mysore.	Dr. Buchanan, 1800.	☉ E.	R. S.	C. S.
decumana.	Mn. Daun gattal.	Amboyna.	Mr. C. Smith.	♄ E.	R.&C.S.	
bicolor, R.		Rohilcund.	A. Gott, Esq. 1803.	♄ E.	R. S.	
pulcherrima, R.		Chittagong.	Mr. J. R. 1810.	♄ E.	R. S.	
crenulata, R.		India.	Dr. F. Buchanan.	♄ E.	R. S.	
involucrata, R.		Moluccas.	Accident.	S. T.	○	
tenacissima, R.	Mn. Ramy.2	Sumatra.	W. Ewer, Esq. 1803.	♄ E.	10---11	
naucliflora, R.	H. Luta-Kudum.	Chittagong.	Mr. J. R. 1810.	♄ Sc.	H.&R.S.	5---6
globulifera, R.	H. Lal-Luta-Kudum.	Chittagong.	Mr. M. R. Smith, 1811.	♄ Sc.	H.&R.S.	5---6
salicifolia, R.		Moluccas.	H. C. 1798.	♄ E.		
MORUS.						
alba.		Hindoostan.	Mr. Maclew.	S. T.	C. S.	C. S.
nigra.		Europe.	H. C. 1795.	T.		
atropurpurea, R.		China.	Before 1794.	T.	C. S.	C. S.
rubra.		America.	W. Hamilton, Esq.	S. T.	10---11	
indica.	B. & H. Toot.3	India.	Indigenous.	S. T.	2	4
tatarica.		In Gardens.		S. T.	2	
paniculata, R.4		Moluccas.	Accident, 1798.	S. T.	R. S.	
mauritiana.		Mauritius.	Mons. Cere, 1810.			

PENTANDRIA.

AMARANTHUS.						
I. *Triandrous.*						
albus.		America.	W. Hamilton, Esq.	☉		
polygonoides.	B. Cheloo-nuteeya.5	India.	Indigenous.	☉ D.	R. S.	R. S.
polygamus.	B. Champa-nuteeya.6	India.	Before 1794.	☉ D.	R. S.	R. S.
tristis.	T. Quoi-tota-kura.7	India.	Before 1794.	☉	H.&R.S.	R. S.
Blitum.		Europe.	1807.	☉ D.	R. S.	
viridis.		India.	Indigenous.	☉ D.	R. S.	R. S.
lividus.	B. Gobura-nuteeya.8	Bengal.	Before 1794.	☉ E.	R. S.	R. S.
oleraceus.	B. Shada-nuteeya.9	India.	Before 1794.	☉ E.	R. S.	R. S.
gangeticus.	H. Lal-sag.10	India.	Before 1794.	☉ E.	R. S.	R. S.
lanceofolius, R.	B. Bans-puta-nuteeya.	Bengal.	Indigenous.	☉ E.	R. S.	R. S.
atropurpureus, R.	B. & H. Lal-nuteeya.11	Bengal.	Dr. W. Carey.	☉ E.	R. S.	
tricolor.		China.	Mr. Honeycomb, 1797.	☉ E.	○	○
melancholicus.	[teeya.12	India.	1800.	☉ E.	R. S.	R. S.
fasciatus, R.	B. Toon-toonee-nu-	India.	Indigenous.	☉ E.	R. S.	R. S.
tenuifolius.	B. Geel-chumli.	Bengal.	Indigenous.	☉ E.	R. S.	R. S.
II. *Pentandrous.*						
hybridus.		Persia.	Major Gen. Malcolm.	☉ E.	R. S.	
frumentaceus.13	Tam. Kiery.	Mysore.	Dr. F. Buchanan, 1801.	☉ E.	R.&C.S.	C. S.
cruentus.		India.	Mr. Atkinson.	☉ E.	R. S.	R. S.
retroflexus.		N. America.	Colonel Garstin.	☉	R. S.	
caudatus.		Persia, &c.	Mr. Lambert.	☉ E.	C. S.	H. S.
paniculatus.		America.	Dr. Klien.	☉ E.	R. S.	R. S.
hypochondriacus.						
spinosus.	H. Kanta-nuteeya.	India.	Indigenous.	☉ E.	R.&C.S.	C. S.
AMBROSIA.						
trifida.		N. America.	W. Hamilton, Esq.		R. S.	C. S.
elatior.		N. America.	W. Hamilton, Esq.		R. S.	C. S.
XANTHIUM.						
indicum.	H. Okura.14	India.	Indigenous.	☉ E.	D. S.	H. S.

1 H. M. 2. t. 41. 2 Poolas, &c. Calooee. Marsd. 75. Ch. Pak-ma. 3 S. Tula. 4 H. A. 4. t. 56. 5 S. Tundooleeya. T. Chirra-kura. 6 T. Dayla-kura. H. Chumli-sag. Of this there are two or more varieties in cultivation. 7 H. A. 5. t. 82. f. 2. Tam. Kuppei-kirei. Several varieties are very generally cultivated over India, one of which I consider Willdenows A. campestris, and another his polystachyus, as Kuppei-kirei, the Tamul name, implies that it may be frequently cut. 8 Or Gobura-dengooa. 9 T. Tota-kura. Of this there are several varieties cultivated as pot-herbs; one very large was called A. giganteus, by Kœnig. 10 Or Dengooa. B. Rukta Shaka. The varieties of this species cultivated by the natives are endless, and in more general use over Bengal than any other. They, as well as oleraceus, and its varieties, differ most conspicuously from the other two extensive tribes of cultivated esculents amaranths, (viz. polygamous and tristis, and their varieties) in not admitting of being cut with the hope of succeeding crops from the same root, which those two tribes do, but are pulled up by the root, and carried to market in that state. 11 H. Bans-putta. 12 Or Bun-nuteeya. T. Chilaka-tola-kura. 13 A very large luxuriant species, extensively cultivated in Mysore, and Coimbetore, for its seed chiefly, the flour of which forms a chief article in the diet of the natives. 14 S. Urishta. T. Veri-talli-napi.

HEXANDRIA.

LEUCACEPHALA.						
graminifolia, R.	B. Gooree.1	India.	Indigenous.	⊙ E.	R. S.	C. S.
spathacea, R.2		India.	Indigenous.	⊙ E.	R. S.	C. S.
COCOS.						
nucifera.	B. Narikel.3	India.	Before 1794.	T.	H. S.	C. S.
BENTINCKIA.						
condapana.	Tam. Kunda-pana.	Malabar.	Dr. A. Berry, 1808.	T.	6	3
ARECA.						
Catechu.	B. Goowa.4	India.	Before 1794.	T.	H. S.	8---9
triandra, R.	B. Ran-goowa.	Chittagong.	Dr. F. Buchanan, 1800.	S. T.	R.&C.S.	H. S.
gracilis, R.	B. Ran-goowa-Sooparee.	Silhet.	Mr. R. Smith, 1810.	♄ E.	D. S.	R. S.
oleracea.		W. Indies.	Hon'ble C. Greville, 1807.			
WRIGHTEA.						
caryotoides, R.	B. Chilputa.	Chittagong.	--- Pierard, Esq. 1800.	♄ E.	R. S.	C. S.
SAGUS.						
spinosus.5		Malay Islands.	H. C. 1798.	T.		
inermis.6		Malay Islands.	H. C. 1798.	T.		

POLYANDRIA.

CERATOPHYLLUM						
verticillatum, R.	H. Jhanji.	Bengal, &c.	Indigenous.	⊙ A.		
SAGITTARIA.						
cordifolia, R.	B. Dal-kuchoo.	India.	Gardiners.	♂ A.	R. S.	C. S.
obtusifolia.7			Kherum.	♂ A.	R.&C.S.	C. S.
sagittifolia.		Bengal.	Indigenous.	♂ A.	3---4	4---5
BEGONIA.						
malabarica.8		Bengal.	Dr. F. Buchanan, 1808.	H.	H.&R.S.	C. S.
laciniata, R.	B. Hoowurjo.	Garrow-hills.	Mr. M. R. Smith, 1812.	♂ or H.	R. S.	C. S.
POTERIUM.						
sanguisorba.		Europe.	1808.		H. S.	
ALEURITES.						
triloba.	C. Kokuna.9	Malay Islands.	1795.	T.	C. S.	7—8
QUERCUS.						
serrata.	B. Gol-shingra.	Garrow hills.	Mr. R. Kyd, 1810.	T.	9---10	
semiserrata, R.	B. Ramkatha.	Garrow hills.	Mr. M. R. Smith, 1812.	L. T.	1---3	8---9
Robur.		Europe.	1798.			
Phellos.		N. America.	W. Hamilton, Esq. 1800.			
paludosus.			Viscount Valentia, 1807.			
lanceæfolia, R.	B. Shingra.	Silhet.	Mr. M. R. Smith, 1811.	L. T.	12	10
squamata, R.	B. Bura-chukma.		Mr. M. R. Smith, 1811.	L. T.	2	10
lappacea, R.	B. Ooloo-chukma.	Garrow-hills.	Mr. M. R. Smith, 1812.	L. T.	12---1	8---9
turbinata, R.	B. Bansooa-batana.	Chittagong.	Mr. J. R. 1810.	L. T.	4---5	9---10
fenestrata, R.	B. Kala-chukma.	Silhet, &c.	Mr. M. R. Smith, 1812.	L. T.	4---5	9---10
castanicarpa, R.	B. Lumba-kanta-hu- [rina-batana.	Chittagong.	Mr. J. R. 1811.	L. T.	7---8	11---12
JUGLANS.						
regia.	H. Akrot.10	N. of India.	1795.	T.		
pterococca, R.	B. Bolus.11	Garrow-hills.	Mr. M. R. Smith, 1812.	T.		7---8
CASTANEA.						
indica, R.	H. Badam-gota.	Silhet.	Mr. M. R. Smith, 1812.	T.	12	9
vesca.12		Europe.	1799.	L. T.		
pumila.	Ch. Fung-lot.	China.	Mr. W. Kerr, 1807.	T.		
SALISBURIA.						
adiantifolia.	Ch. Pac-quo.13	China.	Mr. W. Kerr, 1806--7.	T.		
CARYOTA.						
urens.	T. Teerooga.14	India.	Before 1794.	L. T.	H. S.	
SAGUERUS.						
Rumphii, R.15		Malay Islands.	Before 1794.	T.	H.&R.S.	D. S.

MONADELPHIA.

NIPA.						
fruticans.	B. Goolga.16	Bengal, &c.	Mr. J. R. 1810.	♄		
PINUS.						
sylvestris.		Scotland.	Colonel Beatson, 1810.			
Mughus.		Europe.	1799.			
Pinaster.		Europe.	1799.			
Pinea.		Europe.				
longifolia, R.	H. Cheer.17	Sewalick.	Before 1794.	L. T.		H. S.

1 Eriocaulon Linn. 2 Eriocaulon Linn. 3 S. Narikela. H. Naryel. T. Nari-kadam. A. Narjeel. Of this there are many varieties, chiefly natives of Ceylon, which do not thrive in this latitude. 4 S. Goovaka. H. &. B. Sooparee. T. Poka A. Fooful. 5 Sagus lævis. H. A. 1. p. 76. Willd. 4. 404--5. 6 Sagus genuina. H. A. 1. p. 75. t. 17 & 18. 7 H. M. 11. t. 45. 8 H. M. 9. t. 86. 9 H. A. 2. t. 58. 10 S. Unkotha. A. Jawz, or Shejir-jown, and Akrot. 11 H. Silapoma. 12 Common Chesnut. 13 Gingko biloba, Hort. Kew. 14 H. M. 1. t. 11. H. A. 1. t. 14. 15 H. A. 1. t. 13 Arenga. Labillard. 16 Guana, or Gubna. 17 And its wood Surul.

chinensi .1		China.	Mr. W. Kerr, 1811.			
Dammara.2		Amboyna .	H. C. 1798.	T.		
Deodara, R.	H. Deva-daroo.3	Almora.	T. Brooke, Esq. 1811.	L. T.		8---10
THUJA.						
orientalis.	Ch. Piem-pha.	China.	Before 1794.	S. T.	H. S.	R. S.
occidentalis.	Avbor vitæ.	N. America.	W. Hamilton, Esq. 1809.	L. T.		
CUPRESSUS.						
sempervirens.	H. Sarass.4	India, &c.	Before 1794.	S. T.	H. S.	R. S.
PHYLLANTHUS.						
I. *Leaves simple.*						
madraspatensis.	T. Nalla-usereki.	Coromandel.	Indigenous.	☉ E.	R. S.	C. S.
simplex.	T. *Oochi*-usereki.	Bengal.		☿ D.	R. S.	C. S.
gracilis, R.		Circars.	Nalla-mandu.	☿ E.	R. S.	C. S.
obcordatus, R.		Bengal.	Accident.	☿ E.	○	○
retusus, R.		India.	Colonel Kyd.	☿ E.	○	○
Leucopyrus, Kœn.5			Mr. P. Heshusius.	♄ E.	○	○
virosus.		Circars.	Before 1794.	♄ E.	H.&R.S.	R. S.
II. *Leaves pinnate.*						
Niruri.	B. Sada-hazar-muni.6	India.	Indigenous.	☉ E.	○	○
Urinaria.	B. Hazar-muni.7			☉ E.	○	○
bacciformis.	T. Rawa-salaburta-kanda.	Coromandel.	1796.	♂ or ☿ D.	○	○
pendulus, R.	T. Tella-usereki.	India.	Indigenous.	H. E.	R. S.	C. S.
multiflorus.	B. Pankooshee.8	India.	Indigenous.	♄ Sc.	○	○
Vitis-idæa.	B. Humkata-joolee.9	India.	Indigenous.	♄ E.	○	○
tenellus, R.		Mauritius.	Captain Blake, 1802.	☿ E.	○	○
Kirganelia.		Mauritius.	Captain Tennent, 1803.	♄ E.	○	○
patens, R.		Chittagong.	Mr. W. R. 1801.	♄ E.	○	○
strictus, R.		Moluccas.	Mr. Smith.	♄ E.	○	○
Emblica.	B. Amla.10	India.	Indigenous.	T.	H.&R.S.	C. S.
longifolius.	B. Lodh.11		Before 1794.	T.	H. S.	R.&C.S.
reclinatus, R.		Sumatra.	Mr. W. R. 1804.	♄ D.	R. S.	
tetrandrus, R.	B. Arjura.	Silhet.	Mr. M. R. Smith, 1812.	♄	4---5	9---10
ACALYPHA.						
indica.	B. Shwet-*busunta*.12	India.	Indigenous.	☉ E.	R. S.	C. S.
ciliata.		India.	Indigenous.	☉ E.	H. S.	C. S.
chinensis, R.		China.	Mr. W. Kerr, 1809.	☉ E.	○	○
conferta, R.		China.	Mr. W. Kerr, 1800.	☉ E.	○	○
cylindrica, R.		Mauritius.	Accident, 1811.	☉ D.	○	○
CROTON.						
variegatum. Three var. viz.						
1 var.	medium.13	Moluccas.	G. Westcott, Esq. 1807.	♄ E.	H.&R.S.	R. S.
2 var.	crispum.14	Moluccas.	Mr. C. Smith, 1798.	♄ E.	R. S.	
3 var.	latifolium.	Moluccas.	Mr. C. Smith, 1798.	♄ E.	R. S.	
bicolor, R.		Sumatra.	Mr. W. R. 1803.	L. ♄	3---4	
Tiglium.	H.&B. Jumalgota.15	India.	Before 1794.	L. T.	H. S.	R.&C.S.
oblongifolium, R.	H. Bura-gachh.	Bengal.	Indigenous.	S. T.	H. S.	R. S.
polyandrum, R.	H. Hakooee.16	India.	Before 1794.	♄ E.	R.&C.S.	C. S.
plicatum.	H.&B. Khoodi-Okra.		Indigenous.	☉	C. S.	H. S.
drupaceum, R.		Chittagong.	H. Harris, Esq.	♄ Sc.	3---4	8---9
JATROPHA.						
Curcas.	B.&H. Bagh-bherenda.17	India.	Indigenous.	S. T.	○	○
Manihot.		America.	Before 1794.	L. ♄		
multifida.		S. America.	W. Hamilton, Esq. 1804.	L. ♄	○	○
glandulifera, R.		India.	Mr. Farquaharson, 1796.	♄ E.	○	○
RICINUS.						
communis.	B. Bherenda.18	India.	Before 1794.	S. T.	D. S.	H. S.
Mappa.		Amboyna.	Mr. C. Smith, 1798.	T.	R.&C.S.	
dicoccus, R.		Amboyna.	Mr. C. Smith, 1798.	S. T.	4, 6	R. S.
SAPIUM.						
indicum.	B.&H. Hoorooa.19	Bengal.	Dr. F. Buchanan.	T.	○	○
sebiferum 20		China.	Before 1794.	T.	H. S.	C. S.
baccatum, R.	H. Bela.	Silhet.	Mr. M. R. Smith, 1811.	L. T.	3---4	8---9
BRADLEIA.						
multilocularis, R.21		Bengal.	Dr. F. Buchanan.	S. T.	H.&R.S.	C. S.
nitida, R.	B. Ungootee.	India.		S. T.	C. S.	H. S.
pinnata, R.22		Pulo Penang.	Mr. C. Smith, 1796.	T.	○	
lanceolaria, R.	B. Ungootee.	Silhet.	Mr. M. R. Smith, 1811.	S. T.	3---4	R. S.
BRIEDELIA.						
spinosa, R.	T. Kora-maun.	India.	S. Harris, Esq. 1797.	T.	R. S.	C. S.

1 Ch. Tsam-shew. 2 H. A. 2. t. 57 3 S. Deva-daroo. 4 C. Yeun-pha. A. Sircæ. 5 Fluggea Willd. 6 T. Nella-*U*sereki. 7 T. Yerra-*U*sereki. 8 Or Bun-joolee. T. Nella-pooroogoodoo. 9 T. Yerra-pooroogoodoo. 10 S. Amulukee, and *Umrita*. H. Amlaki. T. *U*sereki. A. Amla. 11 T. Rasa-*U*sereki. H. M. 3. 47. 8. 12 And Moocta-juree. 13 H. A. 4. t. 25. 14 H. A. 4. t. 26. f. 2. 15 S. Jipala. 16 T. Kunda-amadoo. 17 T. Nœpalam. 18 S. Erunda. H. Arundi. T. Amadum. Many varieties in cultivation over the various parts of India. 19 H. M. 4. t. 51. 20 Croton, Linn. Stillingia. Willd. 21 Agyneja. Willd. 22 Phyllanthus obscurus, Willd.

crenulata, R.		Chittagong.	S. Harris, Esq. 1798.	T.	5	10
scandens, R.	T. Doonki-boora.	India.	Indigenous.	L. Sc. ♄	♀	♀
lanceofolia, R.		Bengal.	1808.	T.	R. S.	C. S.
CLUYTIA.						
collina, R.	T. Kodisha.1	Coromandel.	Dr. A. Berry.	T.	H. S.	R. S.
patula, R.		India.		T.	H. S.	R. S.
oblongifolia, R.	B. Dookesswa.	Silhet.	Mr. M. R. Smith, 1812.	T.	H. S.	8—9
HURA.						
crepitans.		Mexico.	W. Hamilton, Esq. 1800.	L. T.	R. S.	C. S.
ANDRACHNE.						
trifoliata, R.		Chittagong.	Dr. F. Buchanan.	T.	H. S.	R. S.

POLYADELPHIA.

TRICHOSANTHES.						
anguina.	B. Chichinga.	India.	Before 1794.	⊙ Sc.	R. S.	C. S.
dioeca, R.	B. Putol.2	Bengal.	Indigenous.	H. Sc.	R. S.	R. S.
cucumerina.	T. Adavi-putla.3	India.	Indigenous.	⊙ Sc.	R. S.	C. S.
lobata, R.	H. Bun-chichinga.	India.	Indigenous.	⊙ Sc.	R.&C.S.	C. S.
cordata, R.	H. Bhooee-koomra.	Bengal.	Mr. W. Jackson, 1802.	H. Sc.	R. S.	
palmata, R.	B.&H. Makhal.4	India.	Indigenous.	♄ Sc.	R. S.	C. S.
heteroclita, R.	H. Goolal.	Silhet.	R. K. Dick, Esq. 1805.	♄ Sc.	H. S.	R. S.
MOMORDICA.						
carantia.	B. & T. Kurilla.5	India.	Before 1794.	⊙ Sc.	R.&C.S.	C. S.
muricata.	B. Oochhya.6	India.	Before 1794.	⊙ Sc.	R.&C.S.	C. S.
monadelpha, R.	B. Tela-koocha.7		Indigenous.	♂ Sc.	♀	♀
dioeca, R.	T. Agokara.8	Circars.	Mr. Heshusius.	H. Sc.	R. S.	C. S.
mixta, R.	H. Kakrol.9	Bengal.	Indigenous.	H. Sc.	R. S.	C. S.
umbellata, R.	B. Kosderi.10	Bengal.	Indigenous.	H. Sc.	R.&C.S.	C. S.
tubiflora, R.		Dacca.	Colonel P. Murray, 1801.	⊙ Cr.	R. S.	R. S.
LUFFA.						
pentandra, R.	B. Dhoondool.11	India.	Indigenous.	⊙ Sc.	R.&C.S.	C. S.
acutangula, R.	B. Jinga.12	India.	Indigenous.	⊙ Sc.	R.&C.S.	C. S.
clavata, R.		Hindoostan.	Colonel A. Kyd, 1802.	♀ E.	R.&C.S.	C. S.
amara, R.	H. Kerula.13	Coromandel.	1802.	⊙ Sc.	R. S.	C. S.
graveolens, R.		Bengal.	Mr. W. R. 1801.	⊙ Sc.	R. S.	C. S.
CUCURBITA.						
lagenaria.	B. Laoo.14	India.	Before 1794.	⊙ Sc.	R.&C.S.	C. S.
Pepo.	H. & B. Koomra.15	India.	Indigenous.	⊙ Sc.	R.&C.S.	C. S.
Melopepo.	H.&B.Suphuree-koomra.	India.	Indigenous.	⊙ Sc.	R.&C.S.	C. S.
citrullus.	H. & B. Turbooz.16	India.	Indigenous.	⊙ D.	D. S.	H. S.
A bitter or wild var.	B. Titu-Laoo.17	India.	Indigenons.	⊙ Sc.	R.&C.S.	C. S.
CUCUMIS.						
Colocynthis.	B. Makhal.18	India.	Dr. A. Berry, 1808.	⊙ D.	R. S.	C. S.
Melo.	B.&H.Khurbooja.19	Persia.	Before 1794.	⊙ D.	H. S.	H. S.
sativus.	B. Susha.20	India.	Before 1794.	⊙ D.	♀	♀
Momordica, R.	B. & H. Footi.21	India.	Before 1794.	⊙ D.	D. S.	H. S.
utilitissimus, R.	B. Kankoor.22	India.	Before 1794.	⊙ D.	H.&R.S.	C. S.
trigonus.	T. Pam-budinga.	India.	Indigenous.	⊙	R.&C.S.	C. S.
turbinatus, R.	T. Nella-budinga.	India.	Nalla-mandu.	⊙	R.&C.S.	C. S.
madraspatanus.	H. Bun-gumuk.23	India.	Indigenous.	⊙	R.&C.S.	C. S.
integrifolius, R.		Bengal.		⊙ Sc.	R. S.	R. S.
BRYONIA.						
scabrella.	H. Agumukee.24	India.	Indigenous.	H. Sc.	♀	♀
Garcini.	T. Gheedi-moraloo.	Coromandel.	1802.	H. Sc.		
tenella, R.		China.	Mr. W. Kerr, 1808.	H. Sc.	C. S.	H. S.
filiformis, R.		Moluccas.	Accident.	H. Sc.	C. S.	
SECHIUM.		Napaul.	Dr. Buchanan, 1802.			

1 Tam. Wodachœ-marum. 2 S. Putolika. H. Pulwul. 3 H. M. S. t. 15. 4 T. Kaki-dunda.
5 Also Oochi, and Poti-kakar. 6 S. Soosuvi. H. Bura kurella. 7 S. Vimbika. T. Kay-dunda.
8 The ♀ and Pooagakarathe ♂. H. M. S. t. 18. 9 Or Gol-kakara. S. Korkotee. 10 H.
Turulli, amoot-mool. T. Teea-dunda. 11 T. Nuni-Beera. H. Purula. N. Pulo. 12 H.
Turi. 13 T. Shetti-beera. 14 Or Kudoo. S. Ulavoo. 15 S. Kurkaroo. T.
Boorda-goomoodoo. 16 Or Turmooj. Water-melon. 17 S. Kutoo-toombee. 18 S.
Indra-varoonee. H. Indraini. T. Sheti-putsa. 19 P. Cherboz. 20 S. Sookhasa.
H. Keera. 21 T. Pedda-dos-kay. 22 H. Kakni. T. Dos-kay. 23 T. Kodi-
budinga. 24 T. Putri-budinga.

CLASSIS XXII.

DIOECIA MONANDRIA.

PANDANUS.						
inermis, R.		Moluccas.	Lieut. M'Kenzie, 1810.	♄		
odoratissimus.	B. Kea.1	India.	Indigenous.	T.	R. S.	C. S.
Millore.2						
fœtidus, R.	B. Kea-kanta.3	Bengal.		♄ D.	C. S.	R. S.
furcatus, R.4		Chittagong.	Mr. J. R. 1811.	♄		
marginatus, R.		Mauritius.	Colonel Hardwicke, 1811.			
amaryllifolius, R.		Amboyna.	H. C. 1796.	♄ Cr.		
inermis, R.			H. C. 1810.	♄ Cr.		
Three more species.5						
NAJAS.						
dichotoma, R.		Bengal.	Indigenous.	☉ A.		
BROSIMUM.						
Alicastrum.6		Jamaica.	W. Hamilton, Esq. 1804.	T.		
CYCAS.						
circinalis.7		Moluccas.	Dr. C. Campbell, 1798.	S. T.	H.S.&C.	C. S.
revoluta.		India.	Before 1794.	S T.	H. S.	C. S.
sphœrica, R.		Moluccas.	H. C. 1800.	S. T.	R.	C. S.
ZAMIA.8		New Holland.	Dr. C. Campbell, 1801.	♄		

DIANDRIA.

VALISNERIA.						
spiraloides, R.		India.	Indigenous.	☉ A.		
octandra.	B. Sheyala.9	India.	Indigenous.	☉ A.		
alternifolia.	B. *Rusna-jaji*.10	India.	Indigenous.	☉ A.		
verticillata.11		India.	Indigenous.			
SALIX.						
tetrasperma, R.	B. Baishi.12	India.	Indigenous.	T.	H.&R.S.	
babylonica.		W. of India.	Before 1794.	T.	H. S.	

TRIANDRIA.

♀ **OSYRIS.**						
peltata, R.	T. Bodie.	Circars.	Nalla-mandu, 1803.	T.	H. S.	R. S.
EXEOECARIA.						
Agallocha.	H. Gewa.13	India.	1800.		H. S.	R. S.
STILAGO.						
Bunias.14		India.	Rajah of Napaul.	T.	3---4	
diandra.		India.	Indigenous.	S. T.	3---4	8
lanceolaria, R.		Chittagong.	Mr. J. R. 1810.	♄	6---7	
tomentosa, R.	B. Tengurec.	Silhet.	Mr. M. R. Smith, 1812.	♄ E.	5---6	9---10

TETRANDRIA.

TROPHIS.						
aspera.	B. Sheora.15	India.	Indigenous.	T.		
spinosa, R.	T. Sukalli.	India.	Mr. J. R. 1810.	T.	C. S.	H. S.
BATIS.						
spinosa, R.	T. Koriti.16	Coromandel.	H. C. 1802.	Sc. ♄		
fruticosa, R.		Chittagong.	Mr. J. R. 1810.	♄ E.	10---5	
NAGEIA.						
Putranjiva, R.	T. *Roodra-joowee*.17	India.	General H. Macdowal, 1802.	T.	3	2
MONETIA.						
barleroides.	B. *Trikanta-jutee*.18	India.	Indigenous.	♄ D.	O	O
MYRICA.						
integrifolia, R.	B. Suphee.	Silhet.	Mr. M. R. Smith, 1812.	T.	12---1	5

1 S. Ketukee. H. Keura. T. Mugali. A. Kadhi. 2 Nicobar bread fruit tree, As. Res. 3. 161. seems only a variety of odoratissima. 3 H. Keur-kanta. 4 H. M. 2. t. 8. 5 Viz. two from Amboyna, and one from the mountains, N. E. of Bengal, have not yet blossomed in the B. G. 6 Jamaica Bread Nut Tree. 7 H. A. 1. t. 22. the ♀ and 23 the ♂. 8 Sp. uncertain, probably spiralis. 9 S. *Sivala*. H. Simar and halla. 10 H. *Janji*. 11 See Serpicula. 12 S. *Vuroona*. H. Pani-joma. 13 *Ugooroo*. 14 H. M. 4. t. 56. 15 S. Sakotuka. H. Suhoora. T. Barinka. 16 H. A. 5. t. 15. f. 2. 17 S. *Pootrunjiva*. 18 H. Kanta-goorkamace. T. Tella-upie.

PENTANDRIA.

PISTACIA.						
officinarum.		Levant.	H. C, 1806.			
ZANTHOXYLUM.						
alatum.	H. Doomur.	Hindoostan.	Colonel Hardwicke, 1800.	T.	H. S.	R. S.
ANTIDESMA.						
pubescens.	T. Polaree.	India.	1803.	S. T.	H. S.	R. S.
paniculata.	H. & B. Khoodi-jam.	Bengal.	Indigenous.	S. T.	H. S.	R. S.
SPINACEA.						
tetrandra, R.	B. Pinnees.1	India.		⊙ E.	C. S.	H. S.
oleracea.2				⊙	C. S.	H. S.
CANNABIS.						
sativa.	B. Ganja.3	India.	Before 1794.	⊙ E	○	○
FERRIOLA.						
buxifolia.	T. Pishanna.4	Coromandel.	Dr. A. Berry, 1802.	S. T.	H. S.	C. S.
SMILAX.						
glabra, R.	H. Hurina.5	Silhet.	Mr. M. R. Smith, 1811.			
herbacea.			W. Hamilton, Esq.			
macrophylla, R.		India.	1806.	H. & ♂ Sc.	H. & R. S.	9---10
pseudo-china.	H. Muheesa.	Silhet, &c.	Mr. M. R. Smith, 1810.	H. Sc.		
bona-nox.			W. Hamilton, Esq.			
maculata, R.		Napaul.	Dr. F. Buchanan.	♄ Sc.	C. S.	H. S.
prolifera, R.		India.	Indigenous.	♄ Sc.	2---3	R. S.
ovalifolia, R.	B. & H. Koomurka.6	India.	Indigenous.	♄ Sc.	H. S.	C. S.
laurifolia.	H. Koomaree.	Silhet.	Mr. M. R. Smith, 1811.	♄ Sc.		
lanceolata, R.	H. Goottia-shook-china.		Mr. M. R. Smith, 1811.	♄ Sc.		
retusa, R.		Bengal.	Indigenous.			
MENISPERMUM.						
cordifolium.	B. & H. Gorkha or Gool.7	India.	Indigenous.	♄ Tw.	H. & R. S.	C. S.
Columba, R.	Columba root, M. M.	Africa.	Dr. A. Berry, 1807.	H. Tw.	H. S.	
tomentosa, R.	B. Pudma-gooluncha.	Bengal.	Indigenous.	♄ Tw.	C. S	H. S.
megaspermum, R.		Silhet.	Mr. M. R. Smith, 1811.	♄ Tw.		H. S.
verrucosum, R.	Mn. Putra wali.	Sumatra.	Captain Wright.	♄ Sc.	2---3	
Cocculus.		Malabar.	B. Heyne, 1806.	♄ Tw.		
hexagynum, R.		China.	Mr. W. Kerr, 1806.	♄ Tw.	○	
hirsutum.	H. Di-er or Hier.8	India.	Indigenous.	♄ Tw.	R. S.	C. S.
polycarpon, R.	H. Bagha-luta.9	India.	Indigenous.	♄ Tw.	R. S.	C. S.
triandrum, R.		Malay Islands.	Mr. C. Smith, 1797.	♄ Tw.	R. S.	
carolinum.		Carolina.	W. Hamilton, Esq.			
canadensis.			W. Hamilton, Esq.			
laurifolium, R.	Pudma-gooluncha.	Napaul, &c.	Dr. F. Buchanan, 1802.	Tw.		
villosum, R.		Chittagong.	Mr. J. R. 1810.	♄ Tw.	R. S.	
DIOSCOREA.10						
globosa, R.	B. & H. Choopri-aloo.	Bengal.	Before 1794.	H. Tw.	R. S.	
alata.	H. & B. Kam-aloo.11	India.	Before 1794.	H. Tw.	R. S.	
rubella, R.	B. & H. Guranya-aloo.	Bengal.	Before 1794.	H. Tw.	R. S.	
purpurea, R.	B. & H. Lal-guranya-aloo.	Bengal.	Before 1794.	H. Tw.	R. S.	
aculeata.	H. & B. Mou-aloo.	Bengal.	Before 1804.	H. Tw.	R. S.	
fasciculata, R.	B. & H. Soosni-aloo.	Bengal.	Before 1794.	H. Tw.	R. S.	
atropurpurea, R.12		Malacca, &c.	Dr. W. Carey, 1808.	H. Tw.		
pulchella, R.		Chittagong.	Mr. J. R. 1810.	H. Tw.	R. S.	
anguina, R.	H. & B. Kookoor-aloo.	Bengal.		H. Tw.	R. S.	
crispata, R.			Dr. W. Carey.	H. Tw.	R. S.	
Nummularia.	B. & H. Shora-aloo.		Indigenous.	♃ Tw.	R. S.	
glabra, R.		Silhet.	Mr. M. R. Smith, 1812.	♄ Tw.		
heterophylla, R.		Amboyna.	Accidental, 1798.	H. Tw.	R. S.	
daemona, R.			Mr. C. Smith, 1798.	H. Tw.	R. S.	
pentaphylla.	B. & H. Kanta-aloo.	India.	Indigenous.	H. Tw.	R. S.	
CALAMUS.						
zalaca.13		Sumatra.	Dr. C. Campbell.	♄ E.		
humilis, R.		Chittagong.	Mr. W. R. 1801.	♄ E.		
erectus, R.	H. Sun-gootta.	Silhet.	Mr. M. R. Smith, 1811.	♄ E.		
Rotang.	H. & B. Beta.14	Bengal.	Indigenous.	♄ Sc.	R. S.	C. S.
rudentum.15		Amboyna.	H. C. 1798.	L. Sc. ♄	H. S.	
verus.16		Moluccas.	H. C. 1798.	L. Sc. ♄		
Draco.17		Sumatra.	Dr. C. Campbell.	♄ Sc.	3---4	
fasciculatus, R.	H. & B. Bura-beta.18	Bengal.	Indigenous.	♄ Sc.	R. S.	C. S.

1 P. Ispanaj. A. Ispanaj. 2 Seeds from Europe at various times. 3 S. Bhanga, Ganjika, &c. H. Bhang. 4 Tam. Irumbilli marum. 5 Shook-china. 6 T. Kondagaroo-tiga. 7 S. Goruksha. T. Tippa-tiga. 8 T. Doosera-tiga. 9 T. Tiga-mushadi. 10 The first eight or nine species are more or less cultivated in various parts of India, and the Isles, for their esculent roots, generally called Yams, and are arranged according to the estimation in which they are held; the first being very generally reckoned the best. Its roots are of a roundish-lobate form, the substance uncommonly white and delicate. 11 H. M. 7. t. 38. 12 Roundish deep purple Malacca Yam. 13 H. A. 5. t. 57. f. 2. 14 Or Sanchi beta. S. Vetra or vetusa. 15 H. A. 5. t. 53. 16 H. A. 5. t. 54. f. 2. 17 H. A. 5. t. 58. f. 1. 18 S. Umba-vetusa.

gracilis, R.	B. Mapoori-bet.	Chittagong.	Mr. W. R. 1801.	♄ Sc.	5---6	C. S.
monoicus, R.		India.	Accident.	♄ Sc.	R. S.	C. S.
latifolius, R.	B. Karak-bet.1	Chittagong.	Mr. W. R. 1801.	♄ Sc.		
polygamus, R.	B. Koodoom-bet.		Mr. W. R. 1801.	♄ Sc.		
tenius, R.	B. Bandhari-bet.		Mr. W. R. 1801.	♄ Sc.		
extensus, R.	H. Deu-gullar.	Silhet.	Mr. M. R. Smith, 1811.	♄ Sc.		
quinquenervius, R.	H. Hurmur-gullar.		Mr. M. R. Smith, 1811.			
PHŒNIX.						
dactylifera.		Arabia, &c.	Mr. Pleydell.	♄ T.	C. S.	H. S.
sylvestris, R.	H. & B. Khajoor.2	India.	Indigenous.			
paludosa, R.	H. & B. Hintal.3	Delta of the Ganges.	1796.	S. T.		
farinifera, R.	T. Chitta-eita.	Coromandel.	Nalla-mandu, 1803.	♄ T.	H. S.	R. S.
acaulis, R.		Bahar, &c.	Before 1794.			
BORASSUS.						
flabelliformis.	H. & B. Tal.4	India.	Indigenous.		3---5	7---9

ENNEANDRIA.

MERCURIALIS.						
annua.		Europe.	Accident.	☉ E.	D. S.	
ambigua.		Europe.	Accident.			
TETRANTHERA.						
nitida, R.	H. Kintoni-jam.	Silhet, &c.	Mr. M. R. Smith, 1812.	L. T.	7---8	4
apetala, R.	S. Guja-pippuli.5	India.	Indigenous.	T.	H. S.	R. S.
monopetala, R.	B. Kookoor-chitta.6	India.	Indigenous.	T.	H. S.	R. S.
quadriflora, R.	B. Sowaree.	Silhet.	Mr. M. R. Smith, 1812.	S. T.	2---3	R. S.
fruticosa, R.	B. Oaloa.	Silhet.	Mr. M. R. Smith, 1812.	♄ E.	6---7	C. S.
lanceofolia, R.	B. Bun-mool.			♄ E.	2---4	R. S.
macrophylla, R.	B. Soorea.7	Bengal.	Mr. M. R. Smith, 1810.	T.	H. S.	R. S.
laurifolia.		Bengal.	Indigenous.	S. T.	H. S.	R. S.
glabra, R.		Coromandel.	Dr. A. Berry.	S. T.	H. S.	R. S.
CARICA.						
Papaya.	H. & B. Papeya.	W. Indies.	Before 1794.	T.	○	○

POLYANDRIA.

FLACOURTIA.						
inermis, R.	M. Tomi-tomi.	Moluccas.	Dr. A. Berry.	T.	R.&C.S.	H.&R.S.
Ramontchi, R.		Madagascar.	Capt. Tennent.	♄ E.		
cataphracta, R.	B. & H. Paniyala.	India.		T.		
sepiaria.	T. Kanrew.8	Coromandel.		♄ E.		
sapida.	B. Bincha.9	India.	Indigenous.	S. T.		
cordifolia, R.		Silhet.	Mr. M. R. Smith, 1810.	♄		
rotundifolia, R.		Coromandel.	Nalla-mandu, 1804.	♄		
Campbelliana, R.		Sumatra.	Dr. C. Campbell, 1800.	S. T.		
ROTTLERA.						
tinctoria.	S. & B. Poonnag.10	India.	1797.	T.		
ferruginea, R.		Pullo-Pinang.	H. C. 1798.	S. T.	H. S.	H. S.
dicocca, R.	B. Akkoos.11	India.	Indigenous.	T.	H. S.	R. S.
alba, R.		P. Pinang.	Mr. W. R.	T.	9---10	
tetracocca, R.	B. Marlea.	Silhet.	Mr. M. R. Smith, 1811.	T.	H. S.	7---8
peltata, R.	B. Seergoolooli.		Mr. R. Kyd, 1810.	T.	H. S.	8
GELONIUM.						
fasciculatum, R.	H. Bun-naringa.12	India.	Indigenous.	S.T.	H. S.	R. S.
bifarium, R.		Uncertain.	Before 1794.	T.	H. S.	R. S.
lanceolatum, R.		Bengal.	Before 1804.	S.T.	3---5	C. S.
TREWIA.						
nudiflora.	H. & B. Pitoli.13	Bengal.	Indigenous.	L. T.	H. S.	R. S.

MONADELPHIA.

JUNIPERUS.						
elata, R.		P. Pinang.	H. C. 1798.	L. T.		
aquatica, R.		China.	Mr. W. Kerr, 1812.			
chinensis.		China.	Mr. W. Kerr, 1812.			
communis, R.		China.	Mr. W. Kerr, 1812.			
cæsia, R.		China.	Mr. W. Kerr, 1812.			
patens, R.		China.	Mr. W. Kerr, 1812.			
TAXUS.						
chinensis, R.	Ch. Lo-hon-tsong.	China.	Mr. W. Kerr, 1811.	♄ E.	C. S.	
elongata.		Cape of G. H.	1799.			
MYRISTICA.						
Moschata.14	B. Jaya-phula.	Moluccas.	H. C. 1797.	T.	R. S.	H.&R.S.
peltata, R.			H. C. 1798.	T.		
spicata, R.			H. C. 1798.	T.		

1 H. M. 12. t. 65. 2 S. Khurjjooree. T. Pedda-eita. 3 S. Hintala. 4 S. Tala.
T. Putoo Toadi ♂. Peati-Toadi ♀. 5 T. Narra-alhoghi. 6 T. Narra-mamadi.
7 On its leaves the Mugadooty silk worms feed. 8 H. M. 5. t. 29. 9 S. Swadoo-kuntuka.
T. Pudda-canrew. 10 T. Vassuntagunda. H. M. 5. t. 21. 11 T. Conda-vettoo. 12 T. Soora-gada. 13 H. M. 1. t. 42. 14 S. Jatee-phula, the nutmeg, and Jatee-putra, the mace.
H. Jayphul, the nutmeg, and Jawatree, the mace. A. & P. Jouz-bewa.

CISSAMPELOS.						
hexandra, R.	H. Nimookka.1	Bengal.	Indigenous.	♄ T.		
convulacca.	T. Pata.2	India.	Dr. Buchanan.	♄ T.		
glabra, R.	H. Peer-grooj.	Silhet.	Mr. M. R. Smith, 1810.	♄ T.	5	10—11
Caapeba.		Moluccas.	Accident.	♄ T.	H. S.	
hernandifolia.		Chittagong.	Mr. J. R. 1810.	♄ T.		
ADELIA.						
nereifolia, R.	H. Heabooka.3	India.	Mr. J. R. 1811.	♄ E.	H. S.	

1 S. Patha umbusta. 3 H. M, 11. t. 62. 3 Tam. Katt-alleræ. T. Janiki.

CLASSIS XXIII.

CRYPTOGAMIA MISCELLANEÆ.

EQUISETUM.						
debilis, R.		Bengal.	Indigenous.	♃ E.	R. S.	
MARSILEA.						
quadrifolia.	B. *Sooshuni*-shak.1	India.	Indigenous.	♃ Cr.	C. S.	C. S.
SALVINEA.						
imbricata.	B. Pana.2	Bengal.	Indigenous.	☉ A.	R. S.	
cuculata.	B. *Indoor*kanee-pana.3			☉ A.	R. S.	
verticillata.	B. *Oolki*pana.			☉ A.	R. S.	
LYCOPODIUM.						
imbricatum, R.	B. Hata-*jori*.	Bengal.	H. Colebrooke, Esq.	♃ Cr.	R. S.	
phlegmaria.	B. Sheetahar.4	Bengal, &c.	1797.	♃ P.	R. S.	
cernuum.		India.		♃ E.	R. S.	
plumosum.		India.		♃ P.	R. S.	
aristatum, R.5		China.	Mr. W. Kerr, 1812.	♃ P.		
ISOETES.						
capsularis, R.		Bengal.	Indigenous.	A.	C. S.	C. S.

FILICES.

OPHIOGLOSSUM.						
cordifolium, R.	B. Dang*a*-ghech*oo*.6	Bengal, &c.	Indigenous.		R. S.	
filiforme, R.7		Bengal.	Dr. F. Buchanan.	♃ Sc.	R. S.	
scandens.8		India.	Indigenous.	♃ Sc.	R. S.	
flexuosum.	B. Bhootraj.9	India.	Indigenous.	♃ Sc.	R. S.	
OSMUNDA.						
zeylanica.	B. Ekbeera.	Amboyna.	Mr. C. Smith.		R. S.	
ACROSTICUM.						
emarginatum.		Bengal.	1796.	♃ Sc.		
POLYPODIUM.						
glabrum, R.	B. Chit*ooa*-bora.	Bengal.	Indigenous.	♃ P.	R. S.	
Phyllitides.		Chittagong.	Dr. Buchanan.	♃		
pertusum, R.		Delta of the Ganges.		♃	R. S.	
quercifolium.	B. Gu*roor*.	India.	Indigenous.	♃ P.	R. S.	
excavatum, R.10		Moluccas.	Mr. Smith.	♃		
flagelliferum, R.		India.	Indigenous.	♃	R. S.	
unitum.	B. Pan-*kakrol*.	India.	Indigenous.			
sophoroides.		Moluccas.	1798.		R. S.	
proliferum, R.	B. Dap*oo*.11	Bengal.	Indigenous.	♃		
semisagittatum, R.		Delta of the Ganges.	Gardiners.			
mucronatum, R.		Silhet.	Mr. M. R. Smith, 1811.			
lucidum.		Napaul.	Dr. F. Buchanan.			
tenerum, R.		Silhet.	Mr. M. R. Smith, 1811.			
ASPLENIUM.						
heterophyllum.						
nidus.		Amboyna.	Mr. C. Smith, 1798.	♃ P.		
bipinnatum, R.			H. S. 1798.	♄		
HEMIONITIS.						
cordifolia, R.	B. Chak*oolya*.	Bengal.	Indigenous.	♃	R. S.	
PTERIS.						
graminifolia.		Silhet.	Mr. M. R. Smith, 1811.	P.		
angustifolia.		Delta of the Ganges.	1796.	♃ P.	R. S.	
vittata.		Delta of the Ganges.	1795.	♃	R. S.	
amplexicaule, R.		Bengal.	Indigenous.	♃	R. S.	
scandens, R.12		India.	Indigenous.	♄ Sc.	R. S.	
lunulata.	H. Kal*ka*-jhamp.13	India.	Indigenous.	♃		
succulenta, R.		India.	Indigenous.	☉ E.	R. S.	
piloselloides.		Chittagong.	Mr. J. R. 1810.	♃ P.		
dimidiata, R.		Silhet.	Mr. M. R. Smith, 1811.	♃	R. S.	
esculenta.14			Mrs. Blaxland, 1811.			
DAVALLIA.						
multiflora, R.		Bengal.	1799.	♃		
pilosa, R.		Bengal.	Indigenous.	♃		
angustifolia.		P. of W. Island.	Mr. W. R.			

1 S. *Soon*i-*shunnaka*. T. Chick-linta-kura. 2 S. Vari-purnee. 3 S. Akhoo-purnee. 4 H. M. 12. t. 14. 5 Dill musc, t. 66. f. 7. 6 Also Ehteera. 7 H. M. 12. t. 34. 8 H. M. 12. t. 33. 9 H. M. 12. t. 32. 10 H. A. 6. t. 35. f. 1. 11 H. Kulke-jamp. 12 H. M. 12. t. 35. 13 H. M. 12. t. 40. 14 Sent from New South Wales.

ADIANTUM. caudatum. LINDSÆA. odorata, R.	H. Pullee.	India. Garrow-hills.	H. T. Colebrooke. Esq. Mr. M. R. Smith, 1812.	℔.	R. S.	R. S.

A

CATALOGUE

OF

PLANTS,

DESCRIBED BY Dr. ROXBURGH,

IN HIS MSS. FLORA INDICA,

BUT NOT YET INTRODUCED INTO THE

Botanical Garden.

SERAMPORE.

1813.

CLASSIS I.

MONANDRIA MONOGYNIA.

ZINZIBER.				
roseum.	T. Bhooma kachkay.	Circars.		
marginatum.				
GLOBBA.				
Hura.1		Malacca.		
pendula.		P. of W. Island.		R. S.
SALICORNIA.				
brachiata.	T. Kiloo.	Coromandel.	♃ E.	○
indica.	B. Jidoo palung.2	Coromandel.	♃ D.	

1 Hura Siamensium. Kon in Rez. obs. 3. 49. 2 T. Kivia pippali.

CLASSIS II.

DIANDRIA MONOGYNIA.

JASMINUM.				
tubiflorum.		Malay Archipelago.	S. Sc.	
OLEA.				
paniculata.		CircarMounts.	S. T.	H. S.
CHIONANTHUS.				
zeylanica.		Ceylon.		
ERANTHEMUM.				
montanum.1		Circars.	♄ D.	C. S.
JUSTICIA.				
1 *Div.*				
bivalves.		Moluccas.	♄	
latebrosa.			H.	R.&C.S.
3 *Div.*				
orixcensis.		Coromandel.	H.	
quinquangularis, K.		Coromandel.	☿	
verticillata, R.	H. Koostoola.	Oojjein.	E. ♄	H. S.
GRATIOLA.				
alata, R.		Moluccas.	☉	
involucrata, R.		Moluccas.	☉ E.	R.&C.S.
rotundifolia, R.		India.	☉ E.	R. S.
racemosa, R.				
cuneifolia, R.		Coromandel.	☉ E.	
lobelioides, R.		Coromandel.	☉	R. S.
SALVIA.				
lanata, R.2		Sirinagur.	H.	

DIANDRIA TRIGYNIA.

PIPER.			
Cubeba.	P. of W. Isld.	♄ Cr.	
lanatum, R.	Malay Archipelago.	♄ Cr.	
methysticum.3	Malay Archipelago.		
arborescens, R.4	Moluccas.	Cr.	
lanceolatum, R.	Moluccas.	♄ Cr.	
Malmaris.5	Moluccas.	♄ Cr.	
rostratum, R.	Moluccas.	Cr.	
diffusum.6			
angustifolium, R.	Moluccas.	Cr.	

1 Justicia montana. Cor. Pl. A. 5. t. 27. 2 S. Integrifolia, A. Res. 3 Sirium decumanum. H 4 H. A. 5. t. 28. f. 1. 5 H. A. 5. t. 16. f. 2. 6 H. A. 5.

CLASSIS III.

TRIANDRIA MONOGYNIA.

COMMELINA.				
moluccana, R.		Moluccas.	⊙ Cr.	
FUIRENA.				
ciliaris, R.	T. Selago boro gaddi.	Coromandel.		
KYLLINGIA.				
cyperoides, R.				
TRINGA.				
triceps, R.		Coromandel.	E.	
levigata, R.			E.	
triandra, R.		Amboyna.	E.	
SCHŒNUS.				
articulatus, R.	T. Kimda-tunga.	Circars.	E.	
CYPERUS.				
setaceus.		Coromandel.		R. S.
mucronatus.		India.		
lateralis.		Coromandel.		
nudus, R.		India.		
diffusus, R.				
squarrosus.		Coromandel.		
aristatus.				
pectinatus, R.		Coromandel.		
niveus, R.				
jemenicus.	T. Poori-gaddi.	Coromandel.		
polystachion.				
castaneus.				
pumilus.				
strictus, R.				
flavidus.		Coromandel.		
umbellatus.	T. Pedda shoka.			
digitatus, R.		India.		
SCIRPUS.				
1. *Leafless.*				
spiralis.		India.		
acutangulus, R.				
spiraloides, R.				
subarticulatus, R.				
dubius, R.	T. Alikay.	Coromandel.		
capitatus.				
junceoides, R.				
triangulatus, R.		Chittagong.		
globosus.	T. Billapa.	Coromandel.		
pectinatus, R.				
2. *Leafy.*				
atropurpureus.				
minimus, R.				
scaber, R.		Coromandel.		
monander.				
antarcticus.				
argenteus, K.				
tenellus, R.		Coromandel.		
tridentatus, R.	T. Goraki shaka tunga.	Coromandel.		
strictus, R.				
brevifolius.				
palescens, R.				
maximus, R.	T. Booda tunga.			

TRIANDRIA DIGINIA.

PEROTIS.				
latifolia.		Coromandel.		
CENCHRUS.				
biflorus, R.		Coromandel.		
SACCHARUM.				
exaltatum, R.	T. Kunda kanumoo.			

ANDROPOGON.		
1. *Spikes terminal.*		
tridentatum, R.		Coromandel.
Ischœmum.		Coromandel.
2. *Long peduncled axillary spikes.*		
monandrum, R.		
strictum, R.		
polystachyum, R.		Coromandel.
3. *Panicles terminal, naked.*		
verticillatum, R.		
saccharoides.		Circar mount.
fascicularis, R.	T. Kunda jerri.	
montanum, R.	T. Aurooch hungul*i*.	
4. *Panicles foliaceous.*		
pumilum, R.		Coromandel.
parviflorum, R.		Circar mount.
LAPPAGO.		
biflora, R.		Coromandel.
PANICUM.		
1. *Spikes simple.*		
indicum.		Coromandel.
dimidiatum.		Coromandel.
2. *Spikes paired.*		
conjugatum, R.		Coromandel.
squarrosum.		
3. *Spikes alternate, secund.*		
grossarium.		
umbrosum.		
4. *Panicled.*		
trigonum.		
patens.		
brevifolium.		
montanum, R.		Circar mount.
AGROSTIS.		
coromandeliana.	T. Yelloo tungoo gaddi.	Coromandel.
Matrella.		Coromandel.
maxima.		
HOLCUS.		
nervosus, R.		Coromandel.
ciliatus, R.		Coromandel.
ISCHŒMUM.		
tenellum, R.		
repens, R.		Pr. of W. I.
MELICA.		
digitata, R.		
CHLORIS.		
tenella, K.		Penins. of India.
montana, R.		
polystachya, R.		Penins. of India.
POMMEREULA.		
cornucopia.		
POA.		
bifaria.	T. Wooda talum.	
procera, R.	T. Reewoona.	
ciliaris.		
cylindrica, R.		
ciliata, R.		
interrupta, K.	T. Nakurnarul.	
viscosa.		
tenella.		
punctata.		
elegans, R.		
flexuosa, R.		
paniculata, R.		
DACTYLIS.		
lagopoides.		
brevifolia.		
ARISTIDA.		
setacea.	T. Shippuroogaddi.	
hystrix.	T. Shippuroo kali.	
depressa.	T. Nali pootiki.	
MANISURUS.		
Myricus.	T. Nali poonookoo.	
ROTBŒLIA.		
corymbosa.	T. Peddoo poonookoo.	
setacea, R.		
Thomæa.		
biflora, R.		Penins. of India.

CLASSIS IV.

TETRANDIA MONOGYNIA.

RHOPALA.				
moluccana.		Pr. of W. I.		
serrata, R.		Malay Archipelago.		
excelsa, R.		Chittagong.	L. T.	
robusta, R.	B. Janeka.	Silhet.	L. T.	
HEDYOTIS.				
geniculata, R.		Malay Isld.		
SPERMACOCE.				
sumatrensis.		Circar mount.	♄	
glabra, R.		Pr. of W. I.	♄	
lineata, R.		Moluccas.		
nana, R.		Honumoa.		
stricta.	T. Loukada.		☉	
IXORA.				
tenuiflora, R.		Moluccas.		
congesta, R.		Moluccas.	♄	
PYROSTRIA.				
hexasperma, R.		Honumoa.		
SALVADORA.				
persica.	T. Pedda waraga winki.	Circars.	S. T.	
CALLICARPA.				
lanata.		Circars.	S. T.	
longifolia.		Pr. of W. I.	♄	R. S.
cuspidata, R.		Moluccas.	♄	
pentandra, R.		Moluccas.	♄	
EXACUM.				
bicolor, R.		Cuttak.	☉	
zeylanicum, R.		Ceylon.	☉	
PLADERA.				
sessiliflora, R.		Coromandel.	☉	
perfoliata, R.		India.	☉	
CISSUS.				
cordata, R.		Amboyna.	♄ Sc.	
setosa, R.				
SAMARA.				
paniculata, R.		Circar mountains.	T.	
polygama, R.		Moluccas.		
BLACKBURNIA.				
monadelpha, R.		Circar mountains.	L. T.	
OLDENLANDIA.				
umbellata.	T. Chesi veloo.1	Coromandel.		
AMMONIA.				
pentandra, R.			Cr.	
TRAPA.				
bicornis.	Ch. Lin-ko.	China.	A.	
POTHOS.				
pinnatifida,* R.		Sumatra.	♃	R. S.
caudata,* R.		Malayana.	♃	R. S.
gracilis, R.		Amboyna.	Sc.	
pertusa, R		Coromandel.	Sc. p.	
pinnata.		Malay Archipelago.		
CANSJERA.				
scandens, R.		Circar mount.	♄ Sc.	
SANTALINUS.				
myrtifolium, R.		Circars.	S. T.	

TETRANDRIA TETRAGYNIA.

MYRIOPHYLLUM.				
tetrandrum.	T. Neela pamarum.		A.	R. S.

1 Tam. Sayawer or Imboril.

* These two species are in the Botanical Garden but were left out of the Catalogue by mistake.

CLASSIS V.

PENTANDRIA MONOGYNIA.

SALSOLA.				
nudiflora.	T. Rawa kada.		♃ D.	☉
indica.	T. Ella koora.		♃ E.	
SPINACHIA.				
tetrandra, R.	B. Pinnis.1	Bengal.	☉ E.	C. S.
ACHYRANTHES.				
sericea, K.			☉	
diandra, R.		Ceylon.	☉	
nodiflora, R.2		Coromandel.	☉ D.	
CEOLOSIA.				
corymbosa.	T. Rajuma.		♃	
polysperma, R.		Moluccas.	☉ E.	
ANDROSACE.				
rotundifolia.		Sirinagur.		
CAPSICUM.				
cerasiforme.				
GENTIANA.				
Cherayti, R.	B. Cherayti.3	Silhet, Napaul.		
CHIRONIA.				
centauroides, R.	B. Girmi.	Bengal.	☉ E.	
CONVOLVOLUS.				
maximus.		Coromandel.	♃ Tw,	
LETTOSMIA.				
uniflora.	T. Boordi-tiga.	Tanjore.	♃ Tw.	
IPOMÆA.				
multiflora, R.			♃ Tw.	
EVOLVOLUS.				
alsinoides.		India.	♃ D.	☉
pilosus, R.		India.	♃ D.	☉
angustifolius, R.		Moluccas.	D.	
CRESSA.				
indica.			☉ E.	
NAMA.				
zeylanica.	S. Languli.4	Bengal.	☉ Cr.	
erecta.		North of Bengal.	☉ E.	
EHRETIA.				
uniflora.		Moluccas.	♃	
CORDIA.				
moluccana, R.		Moluccas.		
FAGROA.				
fragrans, R.		China.	T.	
elliptica, R.		Moluccas.		
CARISSA.				
axillaris, R.		Moluccas.	♄	
HUNTERIA.				
corymbosa, R.		Pr. of W. I.	T.	H. S.
ALYXIA.				
stellata, R.5				C. S.
CERBERA.				
maculata.		Mauritius.		
URCEOLA.				
elastica, R.		Sumatra.	♄ Sc.	
VINCA.				
parviflora.			☉ E.	R. S.
TABERNÆMONTANA.				
corymbosa, R.			♄	
persicariafolia.			S. T.	
NERIUM.				
tomentosum, R.		Circars.	S. T.	H. S.
chinensis.		China.		
caudatum, R.		Coromandel.	♄ Sc.	
ECHITES.				
grandiflora, R.		Silhet.	♄ Sc.	D. S.
marginata, R.	B. Dood luta.	Silhet.	♄ Sc.	
cymosa, R.	B. Ootaso braba.	Silhet.	♄	

1 A. Isfanajor Ispanaj. 2 See Ceolosia nodiflora. Cat. 3 S. Chirataka. Chirata tikta.
4 B. KankraEesh langula. 5 See Gynopogon Stellatum. Cat.

hircosa, R.		Bengal.	♄ Tw.		
parviflora, R.		Circars.	♄ Tw.		
clavata, R.		Moluccas.	♄ Tw.		
ASCLEPIAS.					
1. *Coral rotate.*					
sussuela, R.		Moluccas.	Tw. ☌		
microphylla, R.	T. Poola palla.	Circars.	Tw.	H.&R.S.	
pendula, R.			☌ Tw.		
annularia, R.	T. Poola gurgi.	Coromandel.	☌ Tw.	R.&C.S.	
suberosa, R.					
2. *Coral salver shaped.*					
montana, R.			♄ Tw.		
longistigma, R.			♄ Tw.		
CEROPEGIA.					
bulbosa.	T. Munchi goodi munda.		☌ Tw.		
acuminata.	T. Kumoo mundoo.				
STAPELIA.					
adsendens.	T. Kar alum.		☌	R. S.	
ARDISIA.					
villosa, R.		Pr. of W. Isl.	♄ E.	R. S.	
lanceolaria, R.		Pr. of W. Isl.	♄		
divergens, R.		Moluccas.	♄		
crenata, R.		Pr. of W. Isl.	♄ E.		
SIDEROXYLON.					
tomentosum.			S. T.		
LEEA.					
integrifolia, R.	T. Boora deepaki.	Circars.	♄ L.	R. S.	
ITEA.					
umbellata, R.		Malay Island.	♄		
COMPANULA.					
dehiscens, R.		Bengal.	☉	H. S.	
lanceofolia, R.		Chittagong.		C. S.	
LOBELIA.					
meotianofolia.		Bangalore.			
trigona.			☉		
SCÆVOLA.					
oppositifolia, R.		Moluccas.	♄		
BÆOBOTRYS.					
nemoralis, R.	B. Moolmuriya.	Silhet.	♄	H. S.	
glabra, R.		Chittagong.	S. T.	H. S.	
tetrandra, R.		Moluccas.			
PHYTEUMA.					
begonifolia, R.		Pr. of W. Isl.	Cr.	H. S.	H. S.
MACROCNEMUM.					
stipulaceum, R.		Moluccas.			
parvifloram, R.		Moluccas.	♄		
CINCHONA.					
excelsa, R.	T. Bundoroo.	Circars.	T.	R. S.	
WEBERA.					
macrophylla, R.		Pr. of W. Isl.	♄		
oppositifolia, R.		Chittagong.	♄	H. S.	
RONDELETIA.					
tetrandra, R.		Pr. of W. Isl.	♄		
scandens, R.		Mauritius.	♄ Sc.		H. S.
OPHIORHIZA.					
Mungos.					
villosa, R.		Chittagong.	♄ E.	H. S.	R. S.
GARDENIA.					
1. *Unarmed.*					
caliculata, R.		Coromandel.	♄		
gummifera.		Sirinagur.	♄		
tetrasperma, R.					
2. *Armed.*					
montana, R.	T. Tella kakisha.	Circars.	S. T.	H. S.	
fragrans, K.	T. Pedali.	Chittagong.	♄		
longiflora, R.			S. T.	H. S.	R. S.
CANTHIUM.					
angustifolium, R.	B. Katha mooli.	Chittagong.	♄	H. S.	
didymum.	T. Naum papala.	Coromandel.	S. T.		
moluccanum, R.		Moluccas.			
HAMILTONIA.					
pilosa, R.		Moluccas.	♄		
PSATHURA.					
borbonica.		Mauritius.	S. T.		
PSYCHOTRIA.					
asiatica.		Moluccas.	♄		
COFFEA.					
tetrandra, R.	B. Chilmuri.	Chittagong.	♄	H. S.	D. S

pedunculata, R.		Moluccas.	♄		
angustifolia, R.		Pigeon Island.			
GUETTARDA.					
speciosa.		Coromandel.	S. T.		
MORINDA.					
scandens, R.		Ceylon.	♄ Sc.		
NAUCLEA.					
purpurea.	T. Bagada.	Circars.	S. T.	H. S.	
sessilifolia, R.		Chittagong.	L. T.	C. S.	
glabra, R.		Moluccas.			
rotundifolia, R.		Chittagong.	L. T.	C. S.	H. S.
UNCARIA.					
Gambier, R.		Pr. of W. Isl.	♄ Sc.		
ovalifolia, R.		Pr. of W. Isl.	♄ Sc.		
pedicellata, R.		Moluccas.			
ciriflora, R.		Moluccas.			
acida.		Pr. of W. Isl.			
sclerophylla.		Pr. of W. Isl.			
sessilicarpa, R.		Chittagong.	♄ Sc.		
piloso, R.		Chittagong.	♄ Sc.		
TRIOSTEUM.					
hirsutum, R.		Chittagong.		H. S.	H.&R.S
LONICERA.					
quinquelocularis.	H. Loung phool.	Sirinagur.			
STROEMIA.					
tetrandra.	T. Chimoorda.		♄	☋	
trifoliata, R.		Cootullum.	♄		
BUTTNERA.					
herbacea.		Circars.	E.		
VARECA.					
lanceolata, R.		Pr. of W. Isl.	♄ E.		
VIOLA.					
enneasperma.		Coromandel.	☿ D.		
HARPULLIA.					
cuponioides, R.	B. Harpoolli.	Chittagong.	S. T.		
OPILIA.					
amentacea, R.	T. Balli koma.	Circars.	S. T.		
NEERIA.					
dichotoma, R.	T. Neeriya.	Coromandel.	L. T.	H. S.	
ELÆODENDRUM.					
indicum.	Bois d'olive.	Mauritius.	S. T.	H. S.	C. S.
EUONYMUS.					
glaber, R.		Chittagong.	S. T.	H. S.	
CELASTRUS.					
paniculatus.		Circars.	♄ Sc.	H. S.	
trigynus, R.		Moluccas.			
micranthus, R.		Moluccas.	♄ Sc.		
obtusifolius, R.		Mauritius.	S. T.		
RHAMNUS.					
lucidus, R.		Mauritius.	S. T.		
ZIZYPHUS.					
Oenoplia.		India.	♄	C. S.	

PENTANDRIA DIGYNIA.

ULMUS.					
lanceofolia, R.		Chittagong.	L. T.	H. S.	
CELTIS.					
trinervia, R.		Chittagong.	S. T.	H. S.	
tomentosa, R.		Chittagong.	♄	H. S.	
BOSEA.					
trinervia, R.		Circars.	L. T.		
VAHLIA.					
oldenlandioides, R.		Coromandel.	☉ E.		
viscosa, R.		Coromandel.	☉		
FERULA.					
assafœtida.	B. Hing.1	Persia.			
ANETHUM.					
trifoliatum, R.		Circars.	☉		

PENTANDRIA PENTAGYNIA.

ÆGELATIS.					
rotundifolia, R.		Soonderbunds.	♄		
DROSERA.					
indica.		Coromandel.		C. S.	

1 S. Hing. Arab. Pers. Unjudan.

CLASSIS VI.

HEXANDRIA MONOGYNIA.

BURMANNIA.				
triflora, R.		Pr. of W. Isl.	☉	
TRADESCANTIA.				
tuberosa, R.			☉ Cr.	
paniculata, R.	T. Kunda amadikada.		☉ S. Cr.	
ANTHERICUM.				
uniflorum, R.				
DRACÆNA.				
cernua.		Mauritius.	S. T.	R. S.
umbracaulifera.		Mauritius.	S. T.	R. S.
CANARINA.				
moluccana, R.		Moluccas.	H. E.	
BERBERIS.				
angustifolia, R.		Rohilcund.	♄	

HEXANDRIA DIGYNIA.

LORANTHUS.				
scurrula, R.	T. Wadimki.		P. ♄	
ferrugineus, R.		Pr. of W. Isl.	P. ♄	
involucratus, R.		Chittagong.	P. ♄	☿
ampullaceus, R.		Silhet.	P. ♄	☽ S.
clavatus, R.		Silhet.	P. ♄	
pentapetalus, R.		Silhet.	P. ♄	R. S.
ORIZA.				
coarctata, R.		Delta of the Ganges.	☿ s.	R. S.

HEXANDRIA TRIGYNIA.

APONOGETON.				
echinatum, R.		Circars.		
ANDERSONIA.				
cucullata, R.	B. Amoora.	Delta of the Ganges.	S. T.	R. S.
Rohitoka, R.	B. Tikta raj.1	Bengal.	S. T.	R. S.

1 S. Rohi-tuka. H. Hurinhura.

CLASSIS VIII.

OCTANDRIA MONOGYNIA.

XANTHOPHYLLUM.				
virens, R.	B. Gundhi.	Silhet.	L. T.	H. S.
flavescens, R.	B. Agyasak.	Chittagong.	L. T.	H. S.
OSBECKIA.				
zeylanica.		Ceylon.	☉ E.	
chinensis.		Cuttack.	☉ E.	C. S.
tetrandra, R.		Pr. of W. Isl.	♄ Sc.	
COMBRETUM.				
rotundifolium, R.	B. Kowa luta.	Silhet.	♄ Sc.	C. S.
macrophyllum, R.		Chittagong.	♄ Sc.	C. S.
squamosum, R.		Malay Archipelago.	♄ Sc.	
laxum.	T. Bandi koota.	Circars.	♄ Sc.	
CYMINOSMA.				
pedunculata,		Ceylon.		
MOLINÆA.				
lævis.		Mauritius.	S. T.	
AMYRIS.				
simplicifolia, R.		Pr. of W. Isl.	S. T.	
Commiphora, R.1		Silhet.		
suffruticosa, R.2				
MELICOPE.				
tetrandra, R.		Pr. of W. Isl.	T.	H. S.
MEMECYLON.				
edule, R.		Coromandel.	♄	S. T.
amplexicaule, R.		Pr. of W. Isl.		
ALLOPHYLLUS.				
lanatus.		Pr. of W. Isl. and Silhet.	S. T.	H. S.
ORNITROPE.				
Cobbe.			♄	
integrifolia.		Moluccas.		
repanda, R.		Moluccas.		
SCYTALIA.				
parviflora, R.		Moluccas.		
oppositifolia, R.		Moluccas.		
SAPINDUS.				
squamosus, R.		Moluccas.		
longifolius.		Moluccas.		
serratus, R.		Moluccas.		

OCTANDRIA TRIGYNIA.

1. *style, 2-cleft, seeds without angles.*

POLYGONUM.			
chinensis.	Bengal.	Sc.	H. S.
cymosum, R.	Chittagong.	♄ Sc.	H. S.

1 See A. Agallocha. Cat. 2 See A. Fruticosa. Cat.

CLASSIS IX.

ENNEANDRIA MONOGYNIA.

LAURUS.			
1. *leaves opposite.*			
multiflora.	Ceylon.		
2. *leaves alternate.*			
Camphora.	Sumatra or Java.	T.	
sylvestris.	Circars.		
villosa, R.	Chittagong.	L. T.	C. S.
ANACARDIUM.			
dubium, R.	Sumatra.	L. T.	

ENNEANDRIA HEXAGYNIA.

BUTOMUS.			
anceolatus, R.	Bengal.		

CLASSIS X.

DECANDRIA MONOGYNIA.

BAUHINIA.				
férruginea, R.		Malacca.	♄ Sc.	
integrifolia, R.		Pr. of W. Isl.	♄ Sc.	
cordifolia, R.		Moluccas.	♄ Sc.	
CÆSALPINIA.				
inermis, R.		Moluccas.		
lacerans, R.	T. *Wallikadooda.*		♄ Sc.	R. S.
ADENANTHERA.				
aculeata, R.	T. *Shumi.*	Coromandel.	L. T.	C.&H.S.
PETALOMA.				
alternifolia, R.		Sunderbunds.	L. T.	H. S.
LIMONIA.				
arborea, R.	T. *Kunda gulloonga.*	Circars.	T.	
BOSWELLIA.				
glaber, R.	T. *Googoolapoo chittoo.*	Coromandel.		D. S.
WALSURA.				
ternata, R.	T. *Chinni wallursi.*1		S. T.	H. S.
HEYNIA.				
quinquijuga, R.		Moluccas.	T.	
MELIA.				
tomentosa, R.	Mal. *Burani babi.*	Pr. of W. Isl.	L. T.]	
MELASTOMA.				
ferruginea, R.		Pr. of W. Isl.	♄	
crinita, R.		Chittagong.	♄	
pulchella, R.		Chittagong.	♄	
geniculata, R.		India.	♄	
impuber, R.		Moluccas.		
cordifolia, R.		Chittagong.	♄	
decemfida, R.		Pr. of W. Isl.	♄	
curva, R.		Chittagong.	♄	
furcata, R.		Moluccas.	♄	
dodecandra, R.		Moluccas.		
GASTONIA.				
saruroides, R.		Moluccas.		
ARBUTUS.				
herpeticus, R.		Sirinagur.	T.	
CASEARIA.				
ovata.			L. T.	H. S.
tomentosa, R.	T. *Garugoodoo.*	Circars.	S. T.	
esculenta, R.	T. *Kunda jungira.*	Circars.	L. ♄	
NECTANDRA.				
decandra, R.	B. Hurindur.	Silhet.	L. ♄	D. S.
QUISQUALIS.				
villosa, R.		Pegu.	♄ Sc.	
indica.		Amboyna.	♄ Sc.	
GETONIA.				
floribunda, R.	T. *Bandi mooroodoodoo.*	Circars.		D. S.
TERMINALIA.				
biticaria, R.		Silhet.	L. T.	
Benzoin.		Mauritius.		

DECANDRIA TRIGYNIA.

HIRÆA.				
indica, R.	T. *Rigrak tiga.*	Circars.	♄ Sc.	
rotundifolia, R.		Chittagong.	♄ Sc.	H. S.
ERYTHROXYLON.				
laurifolium.		Mauritius.	T.	R. S.

DECANDRIA PENTAGYNIA.

ROBEIGIA.				
hirsuta, R.		Chittagong.	♄ Sc.	H. S.

1 Tam. Kaka walsura.

CLASSIS XI.

DODECANDRIA MONOGYNIA.

PORTULACA.		
tuberosa, R.	Circars.	R. S.
PEMPHIS.		
angustifolia, R.	Mauritius.	

DODECANDRIA TRIGYNIA.

EUPHORBIA.		
dichotoma, R.	Oojjein.	C. S.
cuneifolia, R.	Hindoostan.	H. S.

CLASSIS XII.

ICOSANDRIA MONOGYNIA.

FABRICIA.					
bracteata, R.		Moluccas.			
METROSIDEROS.					
comosa, R.		Moluccas.			
suberosa, R.		Moluccas.			
CORALLIA.					
lucida, R.	T. Koralli.	Circars and Chittagong.		H. S.	
EUGENIA.					
cerasoides, R.		Chittagong.	L. T.	H. S.	R. S.
præcox, R.		Chittagong.	L. T.	C. S.	
Panialla, R.	B. Panialla jamba.	Chittagong.	L. T.	H. S.	H. S.
laurifolia, R.			L. T.	H. S.	R. S.
angustifolia, R.		Chittagong.	S. T.	H. S.	R. S.
Zeylonica.	B. Nagkeshura jamba.	Silhet.	S. T.	H. S.	
polygama, R.		Malay Archipelago.			
leptosperma, R.		Hunumoa.			
Thumna, R.		Pegu.			
rubens, R.		Chittagong.	L. T.	H.S.	R. S.
EUCALYPTUS.					
moluccana, R.		Moluccas.			
FŒTIDIA.					
mauritiana.		Mauritius.	L. T.		R. S.
PRUNUS.					
sylvatica, R.		Hurdwar.	L. T.		
CHRYSOBALANUS.					
racemosus, R.		Moluccas.			

ICOSANDRIA DIGYNIA.

SESUVIUM.					
portulacastrum.	T. Wangaredokoodoo.	Coromandel.		☉	○

ICOSANDRIA POLYGYNIA.

ROSA.			
pubescens, R.		Rohilcund.	♄
diffusa, R.		China.	♄
RUBUS.			
moluccanus.		Malay Archipelago.	♄
racemosus, R.		Mysore.	♄

CLASSIS XIII.

POLYANDRIA MONOGYNIA.

TERNSTRŒMIA.					
trilocularis, R.		Moluccas.			
BASSIA.					
butyracea, R.	H. Phul-wara.				
DIOSPYROS.					
lanceæfolia, R.	B. Goolal.	Silhet.	L. T.	H. S.	
stricta, R.		Tipperah.	L. T.	H. S.	
bracteata, R.		Dooab.			
1. *Unarmed, spikes globular.*					
MIMOSA.					
Djiringa, R.		Malacca.	L. T.		
monadelpha, R.					
trapezifolia, R.		Moluccas.			
procera, R.	T. Pedda patsiroo.	Coromandel.	L. T.		
2. *Thorny, spikes globular.*					
eburnea, R.		Coromandel.	♄	C. S.	
3. *Thorny, spikes cylindric.*					
obovata, R.		Rohilcund.	T.	H. S.	C. S.
catechuoides, R.			T.	H. S.	C. S.
4. *Prickly, spikes globular.*					
rubicaulis.	B. Kuch kanta.	India.	♄	R. S.	
CAPPARIS.					
subspinosa, R.		Moluccas.	♄		
bisperma, R.	T. Gooli.	Coromandel.	L. T.		
corymbosa, R.	T. Aguba.	Coromandel.	♄ Sc.		
GORDONIA.					
integrifolia, R.	B. Makri-shala.	Silhet.	L. T.		
oblata, R.		Prince of Wales's Island.	T.		
GREWIA.					
pedicellata, R.		Amboyna.			
lanceæfolia, R.		Chittagong.	♄		
excelsa.		Chittagong.	♄		
salvifolia, R.	T. Pastiki.	Coromandel.	♄	R. S.	
hirsuta.	T. Javelliki.	Coromandel.	♄		
trifolia.		Circars.	T.	H. S.	
obliqua, R.		India.	T.		
heteroclita, R.		Moluccas.			
didyma, R.	T. Alpear.	Circars.	T.	R. S.	
paniculata, R.		Prince of Wales's Island.			
begonifolia, R.		Amboyna.	T.	R. S.	
ELÆOCARPUS.					
tuberculatus, R.		Travancore.	T.	H. S.	
fruticosus, R.		Moluccas.	♄		
CALLOPHYLLUM.					
tetrapetalum, R.		Moluccas.			
Suriga.	Mal. Soorgiha.	Malacca.	T.		
HOPEA.					
scaphula, R.	Mal. Bilshor.	Mascal Island.	T.	C. S.	
DIPTEROCARPUS.					
tuberculatus, R.		Chittagong.	T.	H. S.	H. S.
pilosus, R.		Mascal Island.	L. T.		
SHOREA.					
camphorifera, R.	Camphor tree.	Sumatra.	L. T.		
Talura, R.	Tam. Talura.	Balla ghat. M.	L. T.		
longisperma, R.		Pr. of W. Island.			
GARCINIA.					
zeylanica, R.	Cing. Goorka.	Ceylon.			

POLYANDRIA TETRAGYNIA.

TETRACERA.					
sarmentosa.		Pr. of W. Island.	♄ Sc.		
trigyna, R.			♄ Sc.		R. S.
Euryandra.		Moluccas.	♄ Sc.		

POLYANDRIA POLYGYNIA.

UVARIA.					
dicœca, R.	B. Tushee.	Silhet.	♄		
fornicata, R.	B. Luta mal.	Silhet.	♄ Sc.	H. S.	
bicolor, R.	B. Luta mal.	Silhet.	♄ Sc.	H. S.	
cordifolia, R.		Chittagong.	♄	H. S.	C. S.
lutea.	T. Mekari.	Circars.		H. S.	
tomentosa.		Circars.		H. S.	
axillaris, R.		Mauritius.		H. S.	
nitida, R.		Moluccas.			
UNONA.					
discolor.	T. Chilia durguga.	Circars.	L. T.		

CLASSIS XIV.

DIDYNAMIA GYMNOSPERMIA.

ELSHOLTZIA.					
villosa, R.	B. *Rujunta.*	Silhet.	♄	R. S.	
MENTHA.					
paniculata, R.		Chittagong.	☉	C. S.	
auricularia.		Moluccas.			
stellata.					
PHLOMIS.					
urticifolia.		Coromandel.	☉ E.		
moluccana, R.		Moluccas.	☉ E.		
montana, K.			♃		
OCYMUM.					
bullatum.			♂		
thyrsiflorum.					
herbaceum.		Napaul.	♃ H.	R. S.	D. S.
odoratum.		Napaul.	♄		
tuberosum, R.		Orissa.	♃		

DIDYNAMIA ANGIOSPERMIA.

AIGENETIA.					
indica.		Circars.	☉		
BUCHNERA.					
asiatica.	T. Woolisi.		☉ E.		
euphrasioides.		Prince of Wales's Island.	☉ E.		
BARLERIA.					
longiflora.			♄		
RUELLIA.					
racemosa, R.		Prince of Wales's Island.	H. E.		
flava, R.	Bur. Phetchee.	Pegu.			
bracteata, R.		Circars.	♄		
patula.		Nagapatam.	♃		
flagelliformis, R.		Moluccas.	H.		
Pavala, R.	T. Pavala.	Circars.	♄ D.		
suffruticosa, R.		Dinagepore.	♄ E.		
hirsuta, R.		Chittagong.	☉ E.		
uliginosa.		Bengal.	☉ D.	D. S.	
biflora, R.			☉ Cr.		
ROSCOEA.					
villosa, R.		Pegu.	♄ Sc.		
tomentosa.		Chittagong.	♄ Sc.	H. S.	
VOLKAMERIA.					
obovata, R.		Moluccas.			
HASTINGIA.					
scandens, R.	B. Dadmari.	Silhet.	♄ Sc.	C. S.	
VITEX.					
paniculata.		Moluccas.	T.		
PREMNA.					
tomentosa.	T. Naooroo.	Circars.	T.	C. S.	
cordifolia, R.		Prince of Wales's Island.	♄	H. S.	
mucronata, R.		Silhet.	S. T.	H. S.	R. S.
scandens, R.		Silhet.	♄ Sc.		
GMELINA.					
oblongifolia, R.		Eastern border.	L. T.	H. S.	R. S.
TORENIA.					
vagans, R.		Chittagong.	♄	C. S.	
GERARDIA.					
delphinifolia, R.		Circars.	☉ E.	R. & C. S.	
BIGNONIA.					
comosa, R.		Moluccas.			
quadrilocularis, R.		Circars.	L. T.	H. S.	
INCERVILLIA.					
oblongifolio, R.		Chittagong.	♄	C. S.	
CYRILLA.					
aquatica, R.	T. Baggapatti.	Coromandel.	☉	R. & C. S.	

CLASSIS XV.

TETRADYNAMIA SILIQUOSA.

CLEOME.				
diffusa, R.	India.	⊙ D.	C. S.	C. S.
macrophylla.	Coromandel.	⊙ E.	R.&C.S.	C. S.

CLASSIS XVI.

MONADELPHIA DECANDRIA.

MELOCHIA.				
corchorifolio.	B. Beel paat.	India.	☿	R.&C.S. R.&C.S
HELICTERES.				
spicata. Colebrooke.				
SIDA.				
glutinosa, R.			H. E.	
alnifolia.	T. Shya nella benda.	Circars.	♄	R.&C.S. R.&C.S.
chinensis.			♄	
MALVA.				
mauritiana.			☉	C. S.
URENA.				
palmata, R.		Rohilcund.		R. S.
KYDIA.				
paterna, R.	T. Potri.	Circars.	L. T.	R. S.
HIBISCUS.				
1 *Leaves entire, or but slightly lobed, or angular.*				
setosus, R.		Prince of Wales's Island.		
fragrans, R.	B. Kurundur.	Silhet.	S. T.	R. S.
rhombifolius.				
tetralocularis, R.	T. Kunda putta.	Coromandel.	S. T.	
2. *Leaves deeply divided.*				
pumilus, R.		Coromandel.	☉ E.	R.&C.S.
heterophyllus.		N. S. Wales.	S. T.	
PAVONIA.				
zeylanica.		Coromandel.		R. S.

CLASSIS XVII.

DIADELPHIA OCTANDRA.

POLYGALA.				
telephioides.				
undulata, R.		Prince of Wales's Island.	⊙	

DIADELPHIA DECANDRIA.

DALBERGIA.				
alata, R.		Prince of Wales's Island.	S. T.	
parviflora, R.		Moluccas.		
ferruginea, R.		Malay Island.		
Crowey, R.	B. Kroee.	Silhet.	L. T.	
rubiginosa.	T. Tella tiga.	Circars.	♄ Cl.	
volubilis, R.	T. Bandi girgin.	Coromandel.	♄ Tw.	
spinosa, R.		Chittagong.	♄ E.	
GALEDUPA.				
indica.	See G. Arborea, Cat.			
megasperma.				
ERYTHRINA.				
sublobata, R.	T. Badadoomoo.	Circars.	L. T.	
CROTOLORIA.				
1. *Leaves simple.*				
rubiginosa.			♃ D.	
cespitosa, R.	T. Nella guri galli gutsa.	Coromandel.	♃ C.	
uniflora, K.	T. Nella benda.		H. D.	
nummularia.	T. Pili wata tiga.		H.	R.&C.S.
alata, R.		Hindoostan.	H.	R. S.
2. *Leaves compound.*				
orixcensis, R.	T. Koona galli gitsa.		♃ D.	
procumbens, R.			♃ D.	
CARPOPOGON.				
capitatum, R.	T. Surootoo.		⊙ Tw.	
PHASEOLUS.				
maximus.				
DOLICHOS.				
prostratus, K.	T. Kanchi chikurkai.		H. Tw.	C. S.
GLYCINE.				
tenuiflora.		Coromandel.	♃ Tw.	
ROBINIA.				
sennoides, R.		Circars.	♄	H. S.
ferruginea.		Silhet.	♄ Se.	
FLEMINGIA.				
angustifolia, R.		Hardwar.	♄ E.	
1. *Leaves simple.*				
HEDYSARUM.				
Sororium.	T. Nala sara kada.		H. D.	
2. *Leaves ternate.*				
quinquangulatum, R.			⊙ D.	
biarticulatum.	T. Nalla Kavani.		♃ D.	R.&C.S.
umbellatum.	T. Chitkinta.		S. T.	H. S.
arboreum, R.		Moluccas.	S. T.	
DOODIA.				
lagopodioides, R.	See Hedysarum l. Cat.			
hamosa, R.	See Hedysarum hamosum. Cat.			
alopecuroides, R.	See Hedysarum, A. Cat.			
picta, R.	See Hedysarum pictum, Cat.			
crinita, R.	See Hedysarum crinitum, Cat.			
INDIGOFERA.				
echinata.	T. Nalla paleroo.		♃ D.	
aspalathifolia, R.		Coromandel.	♄	R. & C. S.
cinerea.	T. Veri nalli.		♄	R. & C. S.
prostrata.			♄	
flaccida, K.			♄	
violacea, R.		Hardwar.	♄	
GALEGA.				
spinosa.	T. Mulla yampali.	Coromandel.	♄	

PSORALEA. obovata, R.	Hurdwar.	♄	
TRIGONELLA. indica.	Coromandel.	H. D.	☉

CLASSIS XVIII.

POLYADELPHIA POLYANDRIA.

MELALEUCA.		
viridiflora.	Prince of Wales's Island.	L. T.
rigida.	Moluccas.	T.

CLASSIS XIX.

SYNGENESIA ÆQUALIS.

CARDUUS.			
ramosus, R.		☉	C. S.
CACALIA.			
reclinata. R.	Malacca.	♄	
EUPATORIUM.			
flexuosum.	Mauritius.	♄ E.	

SYNGENESIA POLYGAMIA SUPERFLUA.

ARTIMISIA.			
moluccana, R.	Moluccas.	♄ E.	
GNAPHALIUM.			
orixcensis, R.		☉ E.	
CONYZA.			
pubigera.	Moluccas.	♄ Sc.	
fistulosa, R.	Coromandel.	☉ E.	
salicifolia.	Mauritius.	♄	R. S.
SENECIO.			
moluccana, R.	Moluccas.	♄ E.	
ZINNIA.			
bidens.	Coromandel.	♃ E.	R. S.
CHRYSANTHEMUM.			
cuneatum, R.	Moluccas.	♃ D.	
SIEGESBECKIA.			
orientalis.	Moluccas.	☉ E.	
VERBESINIA.			
Boswellia.	Coromandel.	☉ C.	

CLASSIS XX.

GYNANDRIA MONANDRA.

ORCHIS.
platyphyllus.
plantiginœa, R.1 ♃
EPIPACTIS.
trinervia, R. Moluccas.
CYMBIDUM.
aphyllum.
tesselatum.

HEXANDRA.

ARISTOLOCHIA.
longifolia. Moluccas. ♄ Tw.

1 Corom. plants.

CLASSIS XXI.

MONŒCIA MONANDRIA.

AMBROSINIA.				
unilocularis, R.	T. Pecheka boora gaddi.	Coromandel.		R. S.
ARUM.				
lyratum, R.	T. Adivi kunda.	Circars.		
silvaticum, R.	T. Adivi chama.	Circars.		R. S.
ARTOCARPUS.				
hirsuta.	M. Anjeli.	Tillecherry.	L. T.	
lanceolata, R.		Prince of Wales's Island.		
FICUS.				
1 *Sect. Fruit solitary, or paired.*				
palmata, R.		Prince of Wales's Island.		
squamosa, R.		Rohilcund.	♄	
laminosa, R.	N. Chun cheri.	Chinoolee.	♄ D.	
scabrella, R.	B. Bullum dumoor.	Chittagong.	♄ Sc.	
fruticosa, R.	B. Purkutta dumoor.	Chittagong.	♄ L.	
Luducca, R.	H. Looduka.1	Dosa.	♄ L.	
Chincha, R.	H. Chincha.	Dosa.	♄ L.	
vagans, R.		Chittagong.	♄ Sc.	
acuminata, R.		Chittagong.	♄ Sc.	
hederacea, R.	B. Luta purkutti dumoor.	Chittagong.	♄ Sc.	
obtusifolia, R.		Chittagong.	L. T.	
ramentacea, R.		Chittagong.	S. T.	
benjamina.	T. Yerra yerra.	Circars.	L. T.	
tomentosa.	T. Petta mari.	Circars.	L. F.	
comosa.	T. Pootra jivee.	Circars.	L. T.	
urticifolia, R.		North of Bengal.		
ampelos, K. Mss.	T. Tella barinha.	Circars.	L. T.	
asperrima, R.	T. Karakar booda.		L. T.	
tuberculata, R.	T. Kooda juvee.	Coromandel.	S. T.	
angustifolia, R.		Circars.	L. T.	
2 *Sect. Fruit fascicled, racemed, or panicled.*				
rotundifolia, R.		North of Bengal.		

MONŒCIA TRIANDRIA.

CAREX.				
bengalensis, R.		Silhet.		
SCLERIA.				
lithosperma.		Coromandel.		
tenuis.		Moluccas.		
levis.		Malay Archipelago.		
corymbosa, R.		Chittagong.		
setigera, R.		Chittagong.		
TRAGIA.				
cannabina.	T. Doolya gunda.	Coromandel.	♄ Tw.	☌
Camolia.		Coromandel.		

MONŒCIA TETRANDIA.

ALNUS.					
glutinosa.		North of Hindoostan.			
URTICA.					
trinervia, R.		Chittagong.	♄	H. S.	
minima, R.		Homaoon.	☉		
vesicaria, R.		Circars.	♄ E.	R. S.	R. S.
frutescens.		Northern Mountains.	♄		
spherocephala, R.			♄		
paniculata, R.		Moluccas.	♄		
lineata, R.		Moluccas.	L. ♄		
acuminata, R.		Chittagong.	S. T.		
inequalifolia, R.		Chittagong.	♄	H. S.	
MORUS.					
serrata, R.	H. Heemoo.	Dosa.			

MONŒCIA HEXANDRIA.

PHARUS.					
urceolatus, R.		Penang.			
ARECA.					
Dicksonii, R.		Malabar.	S. T.	R. S.	R. S.
disticha, R.		Prince of Wales's Island.	S. T.		
SAGUS.					
Rumphii.		Malay Island.			

MONŒCIA POLYANDRIA.

QUERCUS.					
depressa, R.		Prince of Wales's Island.	L. T.		
lamellata, R.		Prince of Wales's Island.			
glomerata, R.		Prince of Wales's Island.	L. T.		
muricata, R.		Prince of Wales's Island.	T.		
lucida, R.		Prince of Wales's Island.			
ferox, R.	B. Hurineea batana.	Chittagong.	T.		
acuminata, R.	B. Kanta batana.	Chittagong.	L. T		
armata, R.	B. Kanta lal batana.1	Chittagong.	L. T.		
incana, R.		Humaoon.	L. T.		
BEGONIA.					
aptera, R.		Mauritius.	♄		

MONŒCIA MONADELPHIA.

PHYLLANTHUS.					
rhamnoides.		Coromandel.	♄		
peticlaris, R.		Circars.	♄		
turbinatus, Kœn. Mss.	T. Dava dari.	Circars.	T.		
Cheramela, R.2					
ACALYPHA.					
pilosa.		Moluccas.	☉ E.		
amentacea, R.		Moluccas.	♄		
CROTON.					
bractiferum, R.		Moluccas.			
dicœcum, R.		Moluccas.			
asperum, Kœn. Mss.		India.			
Halecum, R.		Moluccas.	T.		
Jouffiah, R.	B. Joffia.	Silhet.	L. T.	H. S.	
SAPIUM.					
cordifolium, R.	T. Badeki.	Circars.	T.		
BRADLEIA.					
impuber, R.		Moluccas.	T.		
pubera, R.		Moluccas.			
hirsuta, R.		Prince of Wales's Island.	♄		
sinica, R.		Prince of Wales's Island.	♄ E.		

MONŒCIA POLYADELPHIA.

LUFFA.					
racemosa, R.		Bengal.	☉ Sc.		
echinata, R.		Coromandel.	Sc.	C. S.	C. S.
Bendaul, R.		Hindoostan.	Sc.		
tuberosa, R.			♃ Sc.		
BRYONIA.					
glabra, R.	T. Naga dunda.		☉ Sc.		
pilosa, R.	T. Kukuma dunda.		☉ Sc.		
laciniosa.	H. Gurga naroo.	India.	Sc.		

MONŒCIA GYNANDRIA.

CLUYTIA.					
semperflorens, R.		Silhet.	♄ E.		
diversifolia, R.		Moluccas.			
BRIEDELIA.					
montana.	T. Patunga.	Circars.	S. T.		

1 Also *Singa-hura*, and Kanta *singa*. 2 Also *Huri-phul*.

CLASSIS XXII.

DIŒCIA TRIANDRIA.

EXEOCARIA.					
integrifolia, R.		Moluccas.			
CATURUS.					
speciflorus.			♄		

DIŒCIA TETRANDRIA.

VISCUM.					
monoecum, R.		Sunderbund.	P. ♄		
verticillatum.	T. Soondra woodiniki.	Coromandel.	P. ♄		
opuntioides.		Coromandel.	P. ♄		
confertum, R.		Silhet.	P. ♄		

DIŒCIA HEXANDRIA.

CALAMUS.					
paniculatus, R.		Prince of Wales's Island.	♄ Sc.		
DIOSCOREA.					
oppositifolia.	T. Ava tenga tiga.	Coromandel.	H. Tw.	R. S.	
tomentosa, Kœn. Mss.		Circars.	H. Tw.		
MENISPERMUM.					
fenestratum, R.	Cing. Wuriwal.	Ceylon.	♄ Sc.	C. S.	H. S.
heteroclytum, R.		Circars.	♄ Tw.		

DIŒCIA POLYANDRIA.

FLACOURTIA.					
obcordata, R.		Chittagong.	♄ E.		
CHAULMOOGRA.					
odorata, R.	B. Chaoolmoogra.1	Silhet.	T.	H. S.	C. S.
ROTTLERA.					
hexandra, R.		Moluccas.			

DIŒCIA MONADELPHIA.

JUNIPERUS.					
cernua, R.	Ch. Ying-lok.	China.	♄		
dimorpha, R.	Ch. Kung-nom-toong.	China.	♄		
CISSAMPELOS.					
tetrandra, R.		Amboyna.	♄ Tw.		
MYRISTICA.					
macrophylla, R.					
montana, R.		Moluccas.			
salicifolia.		Moluccas.			
angustifolia, R.		Arakan.	L. T.		
punicifolia, R.		Moluccas.	S. T.		
linifolia, R.		Chittagong.	L. T.		
ADELIA.					
castanicarpa, R.	Booi Kukra.	Silhet.	L. T.		
cordifolia, R.		Moluccas.			

1 Also Piturkurra.